

1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	

# Name:

## LINEAR ALGEBRA AND VECTOR ANALYSIS

MATH 22B

Total:

### Unit 39: Final Exam Practice A

#### PROBLEMS

**Problem 39A.1) (10 points):**

On the graph  $G$  in Figure 1 we are given a 1-form  $F$  on a graph  $G = (V, E)$ .

- a) (3 points) Write the values of the curl  $dF$ . As a 2-form it is a function on the set  $T$  of triangles.
- b) (3 points) Compute the “discrete divergence”  $d^*F$ , which is a 0-form, a function on the vertices.
- c) (4 points) Find the value of the Laplacian  $d^*dF + dd^*F$  and enter the values near the edges in Figure 2.




FIGURE 1. A graph with a 1-Form  $F$ . Enter here the result for a) and b).




FIGURE 2. Enter here the result for c).

**Problem 39A.2 (10 points) Each question is one point:**

- Who formulated the law of gravity in the form the partial differential equation  $\operatorname{div}(F) = 4\pi\sigma$ ?
- The expression  $5xdxdzdx + 77dydzdy + 3dxdy + 6dydx$  simplifies to ...
- What value is  $\iint_S [x, y, z] \cdot dS$  if  $S$  is the unit sphere oriented outwards?
- What is the distance between the point  $(0, 0, 3)$  and the  $xy$ -plane?
- Is it true that if  $|r'(t)| = 1$  everywhere, then  $r''(t)$  is perpendicular to the velocity  $r'(t)$ ?
- What is the distortion factor  $|dr|$  for the change of coordinates  $r(u, v) = [-2v, 3u]$ ?
- If  $r(u, v)$  parametrizes a surface in  $\mathbb{R}^3$ , is it true that  $r_u \times (r_u \times r_v)$  tangent to the surface?
- Yes or no: if  $(0, 0, 0)$  is a maximum of  $f(x, y, z)$  then  $f_{xx}(0, 0, 0) < 0$ .
- Write down the quadratic approximation of  $1 + x + y + \sin(x^2 - y^2)$ ?
- If  $S : f(x, y, z) = x^2 + y^2 + z^2 = 1$  is oriented outwards, then the flux of  $\nabla f$  through  $S$  is either negative, zero or positive. Which of the three cases is it?

**Problem 39A.3 (10 points) Each problem is 1 point:**

- Which of the triangles in Figure 3 is integrated over in  $\int_0^1 \int_y^1 f(x, y) dx dy$ ?
- We have seen a counter example for Clairaut's theorem. This function  $f(x, y)$  was in  $C^k$  but not in  $C^{k+1}$ . The integer  $k$  indicated how many times we could differentiate  $f$  continuously. What was the  $k$ ?
- To what group of partial differential equations belongs  $\operatorname{div}(E) = 4\pi j + E_t$ ?
- Write down the Cauchy-Schwarz inequality.
- Let  $G$  be the first stage of the Menger sponge (with 20 cubes from 27 cubes present). Is it simply connected?
- Take a exterior derivative of the differential form  $F = \sin(xz)dxdy$ .
- Parametrize the surface  $x = z^2 - y^3$ .
- Parametrize the curve obtained by intersecting of the ellipsoid  $x^2/4 + y^2 + z^2/9 = 1$  with the plane  $y = 0$ .
- What surface is given in spherical coordinates as  $\sin(\phi) \cos(\theta) = \cos(\phi)$ ?
- Write down the general formula for the area of a triangle with vertices  $(0, 0, 0), (a, b, c), (u, v, w)$ .

**Problem 39A.4 (10 points):**

- (6 points) Find the equation of the plane which contains the line  $r(t) = [1+t, 2+t, 3-t]$  and which is perpendicular to the plane  $\Sigma : x+2y-z = 4$ .
- (4 points) What is the angle between the normal vectors of  $\Sigma$  and the plane you just found?




FIGURE 3. Four triangles

**Problem 39A.5) (10 points):**

- a) (8 points) Find the critical points of the function  $f(x, y) = \cos(x) + y^5 - 5y$  and classify them using the second derivative test. You can assume that  $0 \leq x < 2\pi$ .
- b) (2 points) Does the function  $f$  have a global maximum or a global minimum?

**Problem 39A.6) (10 points):**

- a) (5 points) Use the Lagrange method to find the maximum of  $f(x, y) = y^2 - x$  under the constraint  $g(x, y) = x + x^3 - y^2 = 2$ .
- b) (5 points) The Lagrange equations fail to find the maximum of  $f(x, y) = y^2 - x$  under the constraint  $g(x, y) = x^3 - y^2 = 0$ . Still, the Lagrange theorem still allows you to find the maximum. How?

**Problem 39A.7) (10 points):**

- a) (6 points) Find the tangent plane at the point  $P = (4, 2, 1, 1)$  of the surface  $x^2 - 2y^2 + z^3 + w^2 = 2$ .
- b) (4 points) Parametrize the line  $r(t)$  which passes through  $P$  which is perpendicular to the hyper surface at that point. Then find  $(r(1) + r(-1))/2$ .

**Problem 39A.8) (10 points):**

- a) Estimate  $f(0.012, 0.023)$  for  $f(x, y) = \log(1 + x + 3xy)$  using linear approximation.
- b) Estimate  $f(0.012, 0.023)$  for  $f(x, y) = \log(1 + x + 3xy)$  using quadratic approximation.

**Problem 39A.9) (10 points):**

- a) Lets look at the curve which satisfies the acceleration  $r''(t) = [-2 \cos(t), -2 \sin(t), -2 \cos(t), -2 \sin(t)]$ , has the initial position  $[2, 0, 2, 0]$  and initial velocity  $[0, 2, 0, 2]$ . Find  $r(t)$ .
- b) What is the curvature  $|T'(t)|/|r'(t)|$  of  $r(t)$  at  $t = 0$ ?

**Problem 39A.10) (10 points):**

a) Integrate the function  $f(x, y) = x + x^2 - y^2$  over the region  $1 < x^2 + y^2 < 4, xy > 0$ .

b) Find the **surface area** of

$$r(t, s) = [\cos(t) \sin(s), \sin(t) \sin(s), \cos(s)]$$

$$0 \leq t \leq 2\pi, 0 \leq s \leq t/2.$$

**Problem 39A.11) (10 points):**

Let  $E$  be the solid

$$x^2 + y^2 \geq z^2, x^2 + y^2 + z^2 \leq 9, y \geq |x|.$$

a) (7 points) Integrate

$$\iiint_E x^2 + y^2 + z^2 \, dx dy dz.$$

b) (3 points) Let  $F$  be a vector field

$$F = [x^3, y^3, z^3]$$

Find the flux of  $F$  through the boundary surface of  $E$ , oriented outwards.




FIGURE 4. The solid in Problem 10.

**Problem 39A.12) (10 points):**

What is the line integral of the force field  $F(x, y, z, w) = [1, 5y^4 + z, 6z^5 + y, 7w^6]^T + [y - w, 0, 0, 0]^T$  along the path  $r(t) = [t^3, \sin(6t), \cos(8t), \sin(6t)]$  from  $t = 0$  to  $t = 2\pi$ . Hint. We have written the field by purpose as the sum of two vector fields.

**Problem 39A.13) (10 points):**

Find the area of the region  $|x|^{2/5} + |y|^{2/5} \leq 1$ . Use an integral theorem.

**Problem 39A.14) (10 points):**

What is the flux of the vector field  $F(x, y, z, w) = [x + \cos(y), y + z^2, 2z, 3w]$  through the boundary of the solid  $E : 1 \leq x \leq 3, 3 \leq y \leq 5, 0 \leq z \leq 1, 4 \leq w \leq 8$  oriented outwards?

**Problem 39A.15) (10 points):**

Find the **flux** of the curl of the vector field

$$F(x, y, z) = [-z, z + \sin(xyz), x - 3]^T$$

through the **twisted surface** seen in Figure 3 is oriented inwards and parametrized by

$$r(t, s) = [(3 + 2 \cos(t)) \cos(s), (3 + 2 \cos(t)) \sin(s), s + 2 \sin(t)],$$

where  $0 \leq s \leq 7\pi/2$  and  $0 \leq t \leq 2\pi$ .




FIGURE 5. The boundary of the surface is made of two circles  $r(t, 0)$  and  $r(t, 7\pi/2)$ . The picture gives the direction of the velocity vectors of these curves (which in each case might or might not be compatible with the orientation of the surface).