

# Math 191 Notes, 2003 September 23

Given an increasing sequence of sets  $A_1 \subseteq A_2 \subseteq A_3 \dots$ ,

$$A = \bigcup_{i=1}^{\infty} A_i$$

## Inclusion-Exclusion

$$\begin{aligned}\mathbb{P}(A \cup B \cup C) &= \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cap B) \\ &\quad - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C) + \mathbb{P}(A \cap B \cap C)\end{aligned}$$

## Poker

All hands  $\binom{52}{5} = 2568960$

4 of a kind  $13 \cdot 48 = 624$

Full house  $13 \cdot 12 \cdot 4 \cdot 6 = 3744$

3 of a kind  $13 \cdot 4 \cdot 48 \cdot 44 = 109824$

## Bridge

See outline and Durango Bill's website.

## Conditional Probability

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

This is sometimes known as Bayes' rule, given  $\mathbb{P}(B) > 0$ .

Example: Bearded Ben, explosives in shoes.  $A$  is "explosive shoes",  $B$  is "bearded",  $\mathbb{P}(B|A) = 0.6$ ,  $\mathbb{P}(B|A^c) = 0.05$ ,  $\mathbb{P}(A) = 0.2$ .

Lemma:  $A = (A \cap B) \cup (A \cap B^c)$  disjoint,  $\mathbb{P}(A) = \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c)$ ,  $\mathbb{P}(A) = \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^c)\mathbb{P}(B^c)$ .

Continued:  $\mathbb{P}(B) = \mathbb{P}(B|A)\mathbb{P}(A) + \mathbb{P}(B|A^c)\mathbb{P}(A^c) = 0.16$ .

Example: Probability of two-child family which has at least one boy of having the other child being a boy is  $1/3$ .

Example: Monty Hall problem.