

Name:

MWF 9 Chung-Jun John Tsai
MWF 10 Ivana Bozic
MWF 10 Peter Garfield
MWF 10 Oliver Knill
MWF 11 Peter Garfield
MWF 11 Stefan Hornet
MWF 12 Aleksander Subotic
TTH 10 Ana Caraiani
TTH 10 Toby Gee
TTH 10 Valentino Tosatti
TTH 11:30 Ming-Tao Chuan
TTH 11:30 Valentino Tosatti

- Start by printing your name in the above box and check your section in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.
- The hourly exam itself will have space for work on each page. This space is excluded here in order to save printing resources.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) TF questions (20 points)

Mark for each of the 20 questions the correct letter. No justifications are needed.

- | | | |
|-----|---|---|
| 1) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The point $(-1, -1, -1)$ is in spherical coordinate described as $(\rho, \theta, \phi) = (\sqrt{3}, 5\pi, 3\pi/4)$ |
| 2) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | If $ \vec{v} \times \vec{w} = 0$ then $\vec{v} = \vec{0}$ or $\vec{w} = \vec{0}$. |
| 3) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The surface $z^2 + 4y^2 = x^2 + 1$ is a two sheeted hyperboloid. |
| 4) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The surface $4x^2 - 4x + y^2 - 2y - 120 = -z^2$ is an ellipsoid. |
| 5) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The parametrized lines $\vec{u}(t) = \langle 1 + 2t, 2 - 5t, 1 + t \rangle$ and $\vec{v}(t) = \langle 3 - 4t, -3 + 10t, 2 - 2t \rangle$ are the same line. |
| 6) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The surface $\sin(x) = z$ contains lines which are parallel to the y-axis. |
| 7) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | If $\vec{u} \cdot \vec{v} = 0$, $\vec{v} \cdot \vec{w} = 0$ and $\vec{v}$ is not the zero vector, then $\vec{u} \cdot \vec{w} = 0$. |
| 8) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The curvature of a curve depends upon the speed at which one travels upon it. |
| 9) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | Two lines in space that do not intersect must be parallel. |
| 10) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | A line in space can intersect an elliptic paraboloid in 4 points. |
| 11) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | If $\vec{u} \times \vec{v} = 0$ and $\vec{u} \cdot \vec{v} = 0$, then one of the vectors $\vec{u}$ and $\vec{v}$ is zero. |
| 12) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | If the velocity vector $\vec{r}'(t)$ and the acceleration vector $\vec{r}''(t)$ of a curve are parallel at time $t = 1$, then the curvature $\kappa(t)$ of the curve is zero at time $t = 1$. |
| 13) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | If the speed of a parametrized curve is constant over time, then the curvature of the curve $\vec{r}(t)$ is zero. |
| 14) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The length of the vector projection of a vector $\vec{v}$ onto a vector $\vec{w}$ is always equal to the length of the vector projection of $\vec{w}$ onto $\vec{v}$. |
| 15) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | A quadric $ax^2 + by^2 + cz^2 = 1$ is contained in the interior of a sphere $x^2 + y^2 + z^2 < 100$, then the constants a, b, c are all positive and the quadric is an ellipsoid. |
| 16) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | There is a hyperboloid of the form $ax^2 + by^2 - cz^2 = 1$ which has a trace which is a parabola. |
| 17) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The set of points in space which have distance 1 from the line $x = y = z$ form a cylinder. |
| 18) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The velocity vector of a parametric curve $\vec{r}(t)$ always has constant length. |
| 19) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The volume of a parallelepiped spanned by $\vec{u}, \vec{v}, \vec{w}$ is $ (\vec{u} \times \vec{v}) \cdot \vec{w} $. |
| 20) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin: 0 5px;">F</div> | The equation $x^2 + y^2/4 = 1$ in space describes an ellipsoid. |


Match the equation with their graphs. No justifications are needed.



Enter I,II,III,IV here	Equation
	$z = \sin(5x) \cos(2y)$
	$z = \cos(y^2)$
	$z = e^{-x^2-y^2}$
	$z = e^x$

Problem 2b) (4 points)

Match the contour maps with the corresponding functions $f(x, y)$ of two variables. No justifications are needed.



Enter I,II,III,IV,V or VI here	Function $f(x, y)$
	$f(x, y) = \sin(x)$
	$f(x, y) = x^2 + 2y^2$
	$f(x, y) = x + y $
	$f(x, y) = \sin(x) \cos(y)$
	$f(x, y) = xe^{-x^2-y^2}$
	$f(x, y) = x^2/(x^2 + y^2)$

Match the following points in cartesian coordinates with the points in spherical coordinates:

a) $(x, y, z) = (\sqrt{2}, 0, 0)$

b) $(x, y, z) = (0, \sqrt{2}, 0)$

c) $(x, y, z) = (0, 0, \sqrt{2})$

d) $(x, y, z) = (1, 1, 0)$

e) $(x, y, z) = (1, 0, 1)$

f) $(x, y, z) = (0, 1, 1)$

1) $(\rho, \phi, \theta) = (\sqrt{2}, 0, 0)$.

2) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, \pi/4)$.

3) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, 0)$.

4) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, \pi/2)$.

5) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/4, \pi/2)$.

6) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/4, 0)$.

Problem 3) (10 points)

- a) (7 points) Find a parametric equation for the line which is the intersection of the two planes $2x - y + 3z = 9$ and $x + 2y + 3z = -7$.
- b) (3 points) Find a plane perpendicular to both planes and which passes through the point $P = (1, 1, 1)$.

Problem 4) (10 points)


Given the vectors $\vec{v} = \langle 1, 1, 0 \rangle$ and $\vec{w} = \langle 0, 0, 1 \rangle$ and the point $P = (2, 4, -2)$. Let Σ be the plane which goes through the origin $(0, 0, 0)$ and which contains the vectors $\vec{v}$ and $\vec{w}$. Let S be the unit sphere $x^2 + y^2 + z^2 = 1$.

- a) (6 points) Compute the distance from P to the plane Σ .
- b) (4 points) Find the shortest distance from P to the sphere S .

Problem 5) (10 points)

- a) (6 points) Find an equation for the plane through the points $A = (0, 1, 0)$, $B = (1, 2, 1)$ and $C = (2, 4, 5)$.
- b) (4 points) Given an additional point $P = (-1, 2, 3)$, what is the volume of the tetrahedron which has A, B, C, P among its vertices.

A useful fact which you can use without justification in b): the volume of the tetrahedron is $1/6$ of the volume of the parallelepiped which has AB , AC , and AP among its edges.



Problem 6) (10 points)

The parametrized curve $\vec{u}(t) = \langle t, t^2, t^3 \rangle$ (known as the "twisted cubic") intersects the parametrized line $\vec{v}(s) = \langle 1 + 3s, 1 - s, 1 + 2s \rangle$ at a point P . Find the angle of intersection.

Problem 7) (10 points)


Let $\vec{r}(t)$ be the space curve $\vec{r}(t) = (\log(t), 2t, t^2)$, where $\log(t)$ is the natural logarithm (denoted by $\ln(t)$ in some textbooks).

- What is the velocity and what is the acceleration at time $t = 1$?
- Find the length of the curve from $t = 1$ to $t = 2$.

Problem 8) (10 points)

A planar mirror in space contains the point $P = (4, 1, 5)$ and is perpendicular to the vector $\vec{n} = \langle 1, 2, -3 \rangle$. The light ray $\vec{QP} = \vec{v} = \langle -3, 1, -2 \rangle$ with source $Q = (7, 0, 7)$ hits the mirror plane at the point P .

- (4 points) Compute the projection $\vec{u} = \vec{P}_{\vec{n}}(\vec{v})$ of $\vec{v}$ onto $\vec{n}$.
- (6 points) Identify $\vec{u}$ in the figure and use it to find a vector parallel to the reflected ray.



Problem 9) (10 points)

We know the acceleration $\vec{r}''(t) = \langle 2, 1, 3 \rangle + t\langle 1, -1, 1 \rangle$ and the initial position $\vec{r}(0) = \langle 0, 0, 0 \rangle$ and initial velocity $\vec{r}'(0) = \langle 11, 7, 0 \rangle$ of an unknown curve $\vec{r}(t)$. Find $\vec{r}(6)$.

Problem 10) (10 points)

Intersecting the elliptic cylinder $x^2 + y^2/4 = 1$ with the plane $z = \sqrt{3}x$ gives a curve in space.

- (3 points) Find the parametrization of the curve.
- (3 points) Compute the unit tangent vector $\vec{T}$ to the curve at the point $(0, 2, 0)$.
- (4 points) Write down the arc length integral and evaluate the arc length of the curve.