

Name:

MWF 9 Oliver Knill
MWF 10 Hansheng Diao
MWF 10 Joe Rabinoff
MWF 11 John Hall
MWF 11 Meredith Hegg
MWF 12 Charmaine Sia
TTH 10 Bence Béky
TTH 10 Gijs Heuts
TTH 11:30 Francesco Cavazzani
TTH 11:30 Andrew Cotton-Clay

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, and 5, we need to see **details** of your computation.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) True/False questions (20 points), no justifications needed

- 1)  T  F      There is a function  $f(x, y)$  for which the linearization at  $(0, 0)$  is  $L(x, y) = x^2 + y^2$ .
- 2)  T  F      For any two functions  $f, g$  and unit vector  $\vec{u}$  we have  $D_{\vec{u}}(f + g) = D_{\vec{u}}f + D_{\vec{u}}g$ .
- 3)  T  F       $\int_0^2 \int_0^{\sqrt{4-x^2}} (x^2 + y^2) dydx = \int_0^2 \int_0^{\pi/2} r^2 d\theta dr$ .
- 4)  T  F      If we solve  $\sin(y) - xy^2 = 0$  for  $y$ , then  $y' = -y^2/(\cos(y) - 2xy)$ .
- 5)  T  F      If  $f(x, 0) = 0$  for all  $x$  and  $f(0, y) = 0$  for all  $y$ , then  $g(x, y) = \int_0^x \int_0^y f(s, t) dt ds$  solves  $g_{xy}(x, y) = f(x, y)$ .
- 6)  T  F      If  $|\nabla f| = 1$  at  $(0, 0)$ , then there exists a direction in which the slope of the graph of  $f$  at  $(0, 0)$  is 1.
- 7)  T  F      The function  $f(x, y) = x^2 + y^2$  satisfies the partial differential equation  $f_{xx}f_{yy} - f_{xy}^2 = 4$ .
- 8)  T  F      The height of Mount Wachusett is  $f(x, y) = 4 - 2x^2 - y^2$ . On the trail  $x^2 + y^2 = 1$ , the point  $(1, 0)$  is a maximum.
- 9)  T  F      Mount Wachusett has height  $f(x, y) = 4 - 2x^2 - y^2$ . Except at the maximum  $(0, 0)$ , the gradient vector is perpendicular to the graph of the function.
- 10)  T  F      If  $f_x(a, b) > 0$  and  $f_y(a, b) > 0$  then for any unit vector  $\vec{u}$  we must have  $D_{\vec{u}}f(a, b) > 0$ .
- 11)  T  F      If  $f(x, y)$  has two local minima, then  $f$  must have at least one local maximum.
- 12)  T  F      If  $\vec{r}(t)$  is a curve on the surface  $g(x, y, z) = x^2 + y^2 - z^2 = 6$  then  $\nabla g(\vec{r}(t)) \cdot \vec{r}'(t) = 0$ .
- 13)  T  F      If  $f$  and  $g$  have the same trace  $\{x = 5\}$  then  $f_x(5, y) = g_x(5, y)$  for all  $y$ .
- 14)  T  F      If  $f$  and  $g$  have the same trace  $\{x = 5\}$  then  $f_y(5, y) = g_y(5, y)$  for all  $y$ .
- 15)  T  F      The surface area of  $\vec{r}_1(u, v) = \langle u \cos(v), u \sin(v), u^2 \rangle$  and  $\vec{r}_2(u, v) = \langle \sqrt{u} \cos(v), \sqrt{u} \sin(v), u \rangle$  defined on  $\{0 \leq u, v \leq 1\}$  are the same.
- 16)  T  F      If  $\vec{r}(t)$  is a curve on a graph of a function  $f(x, y)$ , then the velocity vector of  $r$  is perpendicular to the vector  $\langle f_x, f_y, -1 \rangle$ .
- 17)  T  F      A continuous function  $f(x, y)$  on the closed disc  $R = \{x^2 + y^2 \leq 51^2\}$  (of course,  $R$  is called “**area**  $51\pi$ ”) has a global maximum on  $R$ .
- 18)  T  F      Any continuous function  $f(x, y)$  has a global minimum and maximum on the curve  $y = x^2$ .
- 19)  T  F      Fubini’s theorem assures that  $\int_a^b \int_c^d f(x, y) dydx = \int_a^b \int_c^d f(x, y) dx dy$ .
- 20)  T  F       $\iint_R \sin(x + y) dx dy = 0$  for  $R = \{-\pi \leq x \leq \pi, -\pi \leq y \leq \pi\}$ .

Problem 2) (10 points)

a) (6 points) Match the integration regions with the integrals. Each integral matches exactly one region  $A - F$ .



Enter A-F	Integral
	$\int_{-1}^1 \int_{-x^2}^{x^2} f(x, y) dydx.$
	$\int_{-1}^1 \int_{-y^2}^{y^2} f(x, y) dx dy.$
	$\int_{-1}^1 \int_{y^2-1}^{1-y^2} f(x, y) dx dy.$
	$\int_{-1}^1 \int_0^{\sqrt{1-y^2}} f(x, y) dx dy.$
	$\int_{-1}^1 \int_{x^2-1}^{1-x^2} f(x, y) dy dx.$
	$\int_{-1}^1 \int_{-\sqrt{1-x^2}}^0 f(x, y) dy dx.$


b) (4 points) Fill in one word names (like “Heat”, “Wave” etc) for the partial differential equations:

Enter one word	PDE
	$g_x = g_y$
	$g_{xx} = g_{yy}$
	$g_{xx} = -g_{yy}$
	$g_x = g_{yy}$

Problem 3) (10 points)


(10 points) A function  $f(x, y)$  of two variables has level curves as shown in the picture. We also see a constraint in the form of a curve  $g(x, y) = 0$  which has the shape of the graph of the cos function. The arrows show the gradient. In this problem, each of the 10 letters  $A, B, C, D, E, F, G, H, K, M$  appears exactly once.

Enter A-P	Description
	a local maximum of $f(x, y)$ .
	a local minimum of $f(x, y)$ .
	a saddle point of $f(x, y)$ where $f_{xx} < 0$ .
	a saddle point of $f(x, y)$ where $f_{xx} > 0$ .
	a saddle point of $f(x, y)$ where $f_{xx}$ is close to zero
	a point, where $f_x = 0$ and $f_y \neq 0$
	a point, where $f_y = 0$ and $f_x \neq 0$
	the point, where $ \nabla f $ is largest
	a local maximum of $f(x, y)$ under the constraint $g(x, y) = 0$ .
	a local minimum of $f(x, y)$ under the constraint $g(x, y) = 0$ .



Problem 4) (10 points)


Find and classify all the extrema of the function  $f(x, y) = x^5 + y^3 - 5x - 3y$ . This function measures “eat temptation” in the  $x$ =Easy- $y$ =Tasty plane. Is there a global minimum or global maximum?



The “Easy-Tasty plane” was introduced in the XKCD cartoon titled “F&#% Grapefruits”.

Problem 5) (10 points)

After having watched the latest Disney movie “Tangled”, we want to build a hot air balloon with a cuboid mesh of dimension  $x, y, z$  which together with the top and bottom fortifications uses wires of total length  $g(x, y, z) = 6x + 6y + 4z = 32$ . Find the balloon with maximal volume  $f(x, y, z) = xyz$ .



Problem 6) (10 points)

a) (8 points) Find the tangent plane to the surface  $f(x, y, z) = x^2 - y^2 + z = 6$  at the point  $(2, 1, 3)$ .

b) (2 points) A curve  $\vec{r}(t)$  on that tangent plane of the function  $f(x, y, z)$  in a) has constant speed  $|\vec{r}'| = 1$  and passes through the point  $(2, 1, 3)$  at  $t = 0$ . What is  $\frac{d}{dt}f(\vec{r}(t))$  at  $t = 0$ ?

Problem 7) (10 points)

a) (5 points) Estimate  $\sqrt{\sin(0.0004) + 1.001^2}$  using linear approximation.

b) (5 points) We know  $f(0, 0) = 1$ ,  $D_{\langle \frac{3}{5}, \frac{4}{5} \rangle}f(0, 0) = 2$  and  $D_{\langle -\frac{4}{5}, \frac{3}{5} \rangle}f(0, 0) = -1$ . If  $L(x, y)$  is the linear approximation to  $f(x, y)$  at the point  $(0, 0)$ , find  $L(0.06, 0.08)$ .

Problem 8) (10 points)

a) (5 points) Find the following double integral

$$\int_0^1 \int_{x^2}^{\sqrt{x}} \frac{\pi \sin(\pi y)}{y^2 - \sqrt{y}} dy dx .$$

b) (5 points) Evaluate the following double integral

$$\int \int_R \frac{\sin(\pi \sqrt{x^2 + y^2})}{\sqrt{x^2 + y^2}} dx dy$$

over the region

$$R = \{x^2 + y^2 \leq 1, x > 0\} .$$

Problem 9) (10 points)

a) (8 points) Find the surface area of the surface parametrized as

$$\vec{r}(u, v) = \langle u - v, u + v, (u^2 - v^2)/2 \rangle ,$$

where  $(u, v)$  is in the unit disc  $R = \{u^2 + v^2 \leq 1\}$ .

b) (2 points) Give a nonzero vector  $\vec{n}$  normal to the surface at  $\vec{r}(4, 2) = \langle 2, 6, 6 \rangle$ .

Problem 10) (10 points)

a) (6 points) Integrate

$$\int_0^{\pi/2} \int_x^{\pi/2} \frac{\cos(y)}{y} dy dx$$

b) (4 points) Find the moment of inertia

$$\iint_R (x^2 + y^2) dy dx ,$$

where  $R$  is the ring  $1 \leq x^2 + y^2 \leq 9$ .