

# Math 213A F23 Homework 1 Solutions

Dhruv Goel

August 2023

If you find any errors, or have any comments or questions, please send them to me via e-mail at this e-mail address.

---

**Q1.** Let  $T \subset \mathbb{R}^3$  be the spherical triangle defined by  $x^2 + y^2 + z^2 = 1$  and  $x, y, z \geq 0$ . Let  $\alpha = z \, dx \, dz$ .

- (a) Find a smooth 1-form  $\beta$  on  $\mathbb{R}^3$  such that  $\alpha = d\beta$ .
- (b) Define consistent orientations for  $T$  and  $\partial T$ .
- (c) Using your choices in [(b)], compute  $\int_T \alpha$  and  $\int_{\partial T} \beta$  directly, and check that they agree. (Why should they agree?)

*Solution.* Many solutions are possible.

- (a) One solution is  $\beta = -\frac{1}{2}z^2 \, dx$ . All possible solutions are obtained by setting  $\beta = f \, dx + g \, dy + h \, dz$ , and solving the triple of PDE's in the smooth functions  $f, g, h$  obtained by requiring  $d\beta = \alpha$ .
- (b) One approach:  $S^2 \subset \mathbb{R}^3$  obtains an orientation as the boundary of the closed disc

$$\mathbb{D}^3 = \{x \in \mathbb{R}^3 : |x| \leq 1\}$$

(which has an orientation as a codimension-0 submanifold-with-boundary of the oriented  $\mathbb{R}^3$ ); namely, if  $u$  is any outward normal vector at a point  $p \in S^2$ , then a basis given by  $v, w \in T_p S^2$  is an oriented basis iff  $u, v, w$  is an oriented basis of  $T_p \mathbb{R}^3$  in that order. The spherical triangle  $T$  obtains an orientation as a codimension-0 submanifold-with-corners of  $S^2$ . Given this, to give  $\partial T$  the boundary orientation (away from the corners), we do the same as before: if  $v$  is any outward normal vector at a boundary point  $p$  (so for instance on the arc  $T \cap \{z = 0\}$  by the vector  $(0, 0, -1)^t$ ), then  $w \in T_p(\partial T)$  is an oriented basis iff  $v, w \in T_p(T)$  give an oriented basis in that order. For the solution, it suffices to provide a picture that looks like Figure 1.

- (c) Let  $D \subset \mathbb{R}^2$  be the region given by

$$D = \{(x, y) : 0 \leq x, y \leq 1 \text{ and } x^2 + y^2 \leq 1\}$$

with the orientation on  $D$  induced by the orientation on  $\mathbb{R}^2$ , and the boundary  $\partial D$  given the boundary orientation. Then the map  $\varphi : D \rightarrow T$  given by  $(x, y) \mapsto (x, y, \sqrt{1 - x^2 - y^2})$  is an oriented parametrization of  $D$  (and its restriction to  $\partial D$  an oriented parametrization of  $\partial T$ ), as one can check explicitly. Therefore, this can be used to compute the integrals above. Namely, we have

$$\varphi^*(dz) = -\frac{x \, dx + y \, dy}{\sqrt{1 - x^2 - y^2}}$$

and so

$$\int_T \alpha = \int_D \varphi^*(\alpha) = \int_D \sqrt{1 - x^2 - y^2} \, dx \wedge \left(-\frac{x \, dx + y \, dy}{\sqrt{1 - x^2 - y^2}}\right) = -\int_D y \, dx \wedge dy = -\int_D y \, dx \, dy.$$

This can be computed by various ways (e.g. a parametrization by polar coordinates); we just use Fubini's Theorem to write this as

$$-\int_0^1 \left( \int_0^{\sqrt{1-x^2}} y \, dy \right) dx = -\int_0^1 \frac{1-x^2}{2} dx = -\frac{1}{3}.$$




Figure 1: Consistent Orientations on  $T$  and  $\partial T$ .

On the other hand, pulling  $\beta$  back to  $\partial D$  via  $\varphi|_{\partial D}$ , note that the only term of the three that survives is the integral along the interval  $[0, 1]$  with its usual orientation (why?), and so

$$\int_{\partial T} \beta = \int_{\partial D} \varphi|_{\partial D}^*(\beta) = \int_0^1 -\left(\frac{1-x^2}{2}\right) dx = -\frac{1}{3}.$$

These are the same, as they should be, because of Stokes' Theorem. ■

*Remark 1.* Note that you are only allowed to compute integrals of forms via oriented parameterizations. A common oversight in the solutions was the failing to check or justify that your parameterization was orientation-preserving.

---

**Q2.** Let  $f(z) = (az + b)/(cz + d)$  be a Möbius transformation. Show [that] the number of rational maps  $g : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  such that

$$g(g(g(g(z)))) = f(z)$$

is 1, 5 or  $\infty$ . Explain how to determine which alternative holds for a given  $f$ .

*Solution.* Many solutions are possible. Clearly there is nothing special about 5; we work with a general  $n \geq 1$ , which is to say, given a Möbius transformation  $f(z)$ , we find the number of rational maps  $g(z)$  such that  $g^{\circ n}(z) = f(z)$ , where  $g^{\circ n}(z)$  means the  $n$ -fold iteration of  $g$ .

- Any non-identity Möbius transformation has either exactly 1 or 2 fixed points. Indeed, write  $f(z) = (az + b)/(cz + d)$  with  $ad - bc \neq 0$  (this is part of the definition of a Möbius transformation). If  $c \neq 0$ , then solving  $f(z) = z$  amounts to solving a quadratic equation (which can have exactly one or two distinct roots), and  $\infty$  is not a fixed point because  $f(\infty) = ac^{-1}$ . If  $c = 0$ , then  $ad \neq 0$ , and so replacing  $a, b$  by  $ad^{-1}, bd^{-1}$  respectively, we can write  $f(z) = az + b$  with  $a \neq 0$ ; in this case,  $f(\infty) = \infty$ , so  $\infty$  is at least one fixed point. Then  $f(z)$  has no more fixed points if  $a = 1$  and  $b \neq 0$ , exactly one more fixed point if  $a \neq 1$ , and in the remaining case  $a = 1, b = 0$ , it is in fact the identity map. The above reasoning shows that, in fact, Möbius transformation  $f$ , we have that  $f$  fixes infinity (i.e.  $f(\infty) = \infty$ ) iff  $f(z) = az + b$  for some  $a, b \in \mathbb{C}$  with  $a \neq 0$ . In this case,  $\infty$  is the only fixed point iff  $a = 1$  and  $b \neq 0$ , and  $\infty$  and 0 are the only fixed points iff  $a \neq 1$  and  $b = 0$ .<sup>1</sup>
- Now, let's move on to the problem. First note that any such  $g$  must be a Möbius transformation itself, by the fact that

$$1 = \deg f = \deg g^{\circ n} = (\deg g)^n.$$

This degree can be the topological degree or the degree as a rational function—we showed in class that these are equivalent. Note also that any fixed point of  $g$  is a fixed point of  $f$ .

- Suppose  $f(z) = z$  is first the identity. Then for any  $c \in \mathbb{C}$ , the Möbius transformation  $g(z) = e^{2\pi i/n}z + c$  is an  $n^{\text{th}}$  functional root of  $f$ . In particular, there are infinitely many such  $g$ .
- Next, suppose  $f$  has exactly one fixed point. Then  $g$  cannot have 2 or more fixed points, and hence  $g$  has exactly one fixed point, namely that of  $f$ . By a change of coordinates (i.e. conjugating by a Möbius transformation that sends this fixed point to  $\infty$ ), we can assume that  $f$  and  $g$  fix only  $\infty$ . By our observation above, this implies that  $f(z) = z + \lambda$  and  $g(z) = z + \mu$  for some  $0 \neq \lambda, \mu \in \mathbb{C}$ . Then  $g^{\circ n}(z) = z + n\mu$ , and so we have  $g^{\circ n}(z) = f(z)$  iff  $n\mu = \lambda$ , and this tells us that there is exactly one such  $g$ .
- Finally, suppose that  $f$  has exactly two fixed points. Then  $g$  must have exactly two fixed points as well; indeed, we showed above that if  $g$  has only one fixed point, then so does every iterate  $g^{\circ n}$  for  $n \geq 1$ . Again, by a change of coordinates, we can assume that both  $f$  and  $g$  fix only  $\infty$  and 0, and then  $f(z) = \lambda z, g(z) = \mu z$  with  $\lambda, \mu \in \mathbb{C} \setminus \{0, 1\}$ . Then  $g^{\circ n}(z) = \mu^n z$ , and so we have  $g^{\circ n}(z) = f(z)$  iff  $\mu^n = \lambda$ , and this tells us there are exactly  $n$  such  $g$ .

This proves:

**Theorem 0.0.1.** Let  $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  be a Möbius transformation. Given any integer  $n \geq 1$ , the number of rational maps  $g : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$  such that  $g^{\circ n}(z) = f(z)$  is exactly 1,  $n$ , or  $\infty$ . More precisely,

- it is 1 iff  $f$  is in some coordinate system a nonzero translation  $z \mapsto z + \lambda$  with  $\lambda \in \mathbb{C}^*$  (or equivalently iff any matrix representing  $f$  is non-diagonalizable, or equivalently iff  $f$  has only one fixed point),
- it is  $n$  iff  $f$  is in some coordinate system a non-identity similarity  $z \mapsto \lambda z$  with  $\lambda \in \mathbb{C} \setminus \{0, 1\}$  (or equivalently iff  $f$  is not the identity but any matrix representing  $f$  is diagonalizable, or equivalently iff  $f$  has two distinct fixed points), and
- it is  $\infty$  iff  $f = \text{id}$ .

■

---

<sup>1</sup>The same result can also be obtained by looking at eigenvalues of any matrix representing  $f$  or considering its Jordan canonical form.

---

**Q3.** Let  $\sum_n a_n z^n$  be the Taylor series for  $\tanh(z)$  at  $z = 0$ .

- (a) What is the radius of convergence of this power series?
- (b) Show that  $a_5 = 2/15$ .
- (c) Give an explicit value of  $N$  such that  $\tanh(1)$  and  $\sum_0^N a_n$  agree to 1000 decimal places. Justify your answer. [Correction: we only ask for you to given an  $N$  such that

$$\left| \tanh 1 - \sum_0^N a_n \right| < 10^{-1000}.$$

In fact, this suffices, because  $\tanh 1$  doesn't have tons of consecutive 9's or 0's at the end of its first 1000 decimal digits, but you don't need to prove that.]

*Solution.*

- (a) The meromorphic function

$$\tanh(z) = \frac{e^z - e^{-z}}{e^z + e^{-z}}$$

on  $\mathbb{C}$  has poles exactly at  $\{(2n+1)\pi i/2\}_{n \in \mathbb{Z}}$ . In particular, the poles closest to zero are  $\pm\pi i/2$ , so the radius of convergence of this series must be  $\pi/2$ , since the power series of a holomorphic function around a point in a region in the complex plane is valid (i.e. converges absolutely and locally uniformly) in the largest open disc contained in that region (this follows from the Cauchy integral formula; see [1, Thm. 1.1.2.3]).

- (b) One possible solution: differentiate a few times and plug in 0 (and divide by 5!). Alternatively, compute a suitable integral; namely

$$a_5 = \operatorname{Res}_{z=0} z^{-6} \tanh(z) = \frac{1}{2\pi i} \oint_{S^1} z^{-6} \tanh(z) dz.$$

Alternatively, you could expand

$$\begin{aligned} \tanh(z) &= \frac{\sinh(z)}{\cosh(z)} \\ &= \frac{\left(z + \frac{z^3}{6} + \frac{z^5}{120} + [z^6]\right)}{\left(1 + \frac{z^2}{2} + \frac{z^4}{24} + [z^6]\right)} \\ &= \left(z + \frac{z^3}{6} + \frac{z^5}{120}\right) \left(1 - \left(\frac{z^2}{2} + \frac{z^4}{24}\right) + \left(\frac{z^2}{2} + \frac{z^4}{24}\right)^2\right) + [z^6] \\ &= \left(z + \frac{z^3}{6} + \frac{z^5}{120}\right) \left(1 - \frac{z^2}{2} + \frac{5z^4}{24}\right) + [z^6], \end{aligned}$$

where we use the notation of [1, §5.1.2]. At this point, collecting coefficients of  $z^5$  yields

$$a_5 = \frac{5}{24} - \frac{1}{12} + \frac{1}{120} = \frac{2}{15}.$$

- (c) In general, we use the following version of Taylor's Theorem with remainder:

**Theorem 0.0.2.** Let  $\Omega \subset \mathbb{C}$  be an open subset and let  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. If  $\sum_n a_n (z-a)^n$  is the Taylor series for  $f(z)$  at  $z = a$  for some  $a \in \Omega$ , i.e. if  $a_n = \frac{f^{(n)}(a)}{n!}$  for  $n \geq 0$ , then for any integer  $N \geq 0$ , we can write

$$f(z) = \sum_{n=0}^N a_n (z-a)^n + f_{N+1}(z)(z-a)^N$$

---

for some holomorphic function  $f_{N+1} : \Omega \rightarrow \mathbb{C}$ . If  $r > 0$  is chosen such that the open disk  $\Delta(a; r) = \{z \in \mathbb{C} : |z - a| < r\}$  is contained in  $\Omega$  and is  $S^1(a; r) = \partial\Delta(a; r)$  is the circle of radius  $r$  at  $a$  oriented counterclockwise, then for any  $z \in \Delta(a; r)$ , we have the representation

$$f_{N+1}(z) = \frac{1}{2\pi i} \oint_{S^1(a; r)} \frac{f(\zeta) d\zeta}{(\zeta - a)^{N+1}(\zeta - z)}.$$

In particular, if  $M_r := \sup_{\zeta \in S^1(a; r)} |f(\zeta)|$ , then for all  $z \in \Delta(a; r)$  we have

$$|f_{N+1}(z)| \leq \frac{M_r}{r^N(r - |z - a|)}.$$

*Proof.* See [1, §4.3.1]. ■

Let's apply this to our case, with  $\Omega = \Delta(\pi/2)$  and  $f(z) = \tanh(z)$  around  $a = 0$ . Then for any  $r$  with  $1 < r < \pi/2$ , we have for all  $z \in \Delta(r) = \Delta(0; r)$  that

$$|\sinh(z)| \leq \frac{e^{\operatorname{Re} z} + e^{-\operatorname{Re} z}}{2} \leq e^{\pi/2}$$

and

$$|\cosh(z)| = |\cosh x \cos y + i \sinh x \sin y| \geq |\cosh x \cos y| \geq |\cos y|$$

if  $z = x + iy$  with  $x, y \in \mathbb{R}$  because  $\cosh x \geq 1$  for all  $x \in \mathbb{R}$ . It follows that for  $1 < r < \pi/2$  we have

$$M_r \leq e^{\pi/2} \sec r.$$

Applying the above bound for  $z = 1$  gives us that for any  $N \geq 0$ , we have

$$\left| \tanh(1) - \sum_{n=0}^N a_n \right| = |f_{N+1}(1)| \leq \frac{e^{\pi/2} \sec r}{r^N(r-1)}.$$

Therefore, for any  $r$  with  $1 < r < \pi/2$ , given any  $\varepsilon > 0$ , we can take

$$N > \frac{\pi/2 + \ln \sec r - \ln(r-1) - \ln \varepsilon}{\ln r}$$

to give us an error

$$\left| \tanh(1) - \sum_{n=0}^N a_n \right| < \varepsilon.$$

Now take any  $r \in (1, \pi/2)$  and  $\varepsilon = 10^{-1000}$  to get your  $N$ , e.g. taking  $r = 3/2$  tells us that  $N = 5691$  suffices. ■

*Remark 2.* For  $\varepsilon = 10^{-1000}$ , the expression on the right as a function of  $r$  achieves a minimum at around  $r \approx 1.5705$  of around 5123.773, so  $N = 5124$  certainly suffices and is the best bound we get from this method. A few lines of Sage code tell us that the least  $N$  for which the problem statement is true is  $N = 5099$ , so our estimate using complex analysis is not bad at all!

*Remark 3.* A lot of solutions either did not bound  $\tanh(z)$  at all, or bounded it incorrectly, claiming for instance that  $|\tanh(z)| \leq 1$  for all  $|z| < \pi/2$ . This statement is not true; indeed,  $|\tanh x| \leq 1$  for all  $x \in \mathbb{R}$ , but  $\tanh(z)$  has a pole, and is in particular unbounded in absolute value, as  $z \rightarrow \pm i\pi/2$ .

---

**Q4.** Let  $f : U \rightarrow V$  be a proper local homeomorphism between a pair of open sets  $U, V \subset \mathbb{C}$ . Prove that  $f$  is a covering map.

*Solution.* Here's the definition of covering space that I use:

**Definition 0.0.3.** A map  $p : X \rightarrow Y$  of topological spaces is a **covering space** if for each  $y \in Y$  there is an open neighborhood  $W$  of  $y$  in  $Y$ , a discrete space  $\Gamma$ , and a homeomorphism  $\Phi : p^{-1}(W) \rightarrow W \times \Gamma$  that commutes with projection to  $W$ , i.e. satisfies  $p(x) = \text{pr}_W(\Phi(x))$  for all  $x \in p^{-1}(W)$ , where  $\text{pr}_W : W \times \Gamma \rightarrow W$  is the first projection.

*Remark 4.* Note that  $\Gamma = \emptyset$  is allowed. We call any such  $W$  an evenly covered neighborhood of  $y$ . When our spaces locally path-connected Hausdorff spaces, we can assume by shrinking  $W$  that  $W$  is connected, and then the second condition above is equivalent to saying that  $p$  maps each connected component of  $p^{-1}(W)$  homeomorphically onto  $W$ . The cardinality of any such  $\Gamma$  is called the local degree of this covering map at  $y$ ; this is a locally constant function of  $y \in Y$ . Usually, we require that the local covering degree be positive or at most countable (the former is equivalent to surjectivity, and the latter is automatic if  $X$  and  $Y$  are manifolds by the second countability/paracompactness hypothesis), but neither is strictly logically necessary.

The only property of open subsets of  $\mathbb{C}$  we'll use is that they are locally compact Hausdorff. We show:

**Theorem 0.0.4.** Let  $p : X \rightarrow Y$  be a proper local homeomorphism between locally compact Hausdorff spaces. Then  $p$  is a covering map.

*Proof.* For any  $y \in Y$ , the fiber  $p^{-1}(y)$  is discrete (because  $p$  is a local homeomorphism) and compact (because  $p$  is proper), and hence a finite set. Fix a  $y \in Y$ , and let  $p^{-1}(y) = \{x_1, \dots, x_n\}$  for some  $n \geq 0$  and  $x_i$  distinct. For each  $i = 1, \dots, n$ , there is an open neighborhood  $U_i \subset X$  such that  $p(U_i) \subset Y$  is open and  $p|_{U_i} : U_i \rightarrow p(U_i)$  is a homeomorphism. Since  $X$  is Hausdorff and  $n$  is finite, we may shrink the  $U_i$  to be pairwise disjoint. It is not true that any open neighborhood  $W$  of  $y$  contained in  $O := \bigcap_{i=1}^n p(U_i)$  is evenly covered, because there is no guarantee that  $p^{-1}(W) \subset \bigcup_{i=1}^n U_i$ . However, when  $p$  is proper, there is some neighborhood  $W$  of  $y$  contained in  $O$  which satisfies  $p^{-1}(W) \subset \bigcup_{i=1}^n U_i$ , and then this neighborhood is evenly covered, finishing the proof.

There are various ways to produce such a  $W$ . We mention three:

- Take any precompact open neighborhood  $V \subset Y$  of  $y$  (i.e. with  $\bar{V}$  compact) satisfying  $\bar{V} \subset O$ , and define

$$W := V \setminus p \left( p^{-1}(\bar{V}) \setminus \bigcup_{i=1}^n U_i \right).$$

Note that  $p^{-1}(\bar{V})$  is compact because  $p$  is proper, and hence so is  $p^{-1}(\bar{V}) \setminus \bigcup_{i=1}^n U_i$ , and hence so is its image in  $p$ . It follows that this image is closed (since  $Y$  is Hausdorff), and hence  $W$  is an open neighborhood of  $y$  in  $Y$  with the required property.

- Another, similar, way to proceed is to use the following helpful lemma:

**Lemma 0.0.5.** Let  $p : X \rightarrow Y$  be a proper map between topological spaces with  $Y$  locally compact and Hausdorff. Then  $p$  is a closed map.

*Proof.* Let  $C \subset X$  be closed, and let  $y \in Y \setminus p(C)$ . Pick a precompact neighborhood  $U$  of  $y$  (i.e. an open neighborhood with compact closure  $\bar{U}$ ). Since  $p^{-1}(\bar{U})$  is compact, so is its closed subset  $K := C \cap p^{-1}(\bar{U})$ , and therefore  $p(K)$ . Since  $Y$  is Hausdorff,  $p(K)$  is closed in  $Y$ , and then  $y \in U \setminus p(K) \subset Y \setminus p(C)$ , proving  $Y \setminus p(C)$  is open. ■

Having done this, look at  $p(X \setminus \bigcup_{i=1}^n U_i)$ , which is a closed subset of  $Y$  because  $p$  is closed, and note that taking

$$W := O \setminus p \left( X \setminus \bigcup_{i=1}^n U_i \right)$$

works.

- 
- This third method works only in our setting of open subsets of the plane<sup>2</sup> Note that for any  $y' \in O$ , the fiber  $p^{-1}(y')$  has cardinality at least  $n$  (why?). It suffices to show that there is some neighborhood  $W$  of  $y$  in  $Y$  contained in  $O$  such that for all  $y' \in W$ , the fiber  $|p^{-1}(y')|$  has cardinality exactly  $n$ , because this would prove that  $p^{-1}(W) \subset \bigcup_{i=1}^n U_i$  (why?). Suppose this is not the case, so there is a sequence of points  $\{y_k\}_{k \geq 1}$  in  $O$  converging to  $y$ , where each  $y_k$  has at least  $n + 1$  preimages. Let  $K := \{y_k\}_{k \geq 1} \cup \{y\}$ , and note that this is a compact subset of  $Y$ . We claim that the preimage  $p^{-1}(K) \subset X$  is not compact, and this is the required contradiction. Indeed, for each  $y_k$ , pick a preimage  $x_k$  that is not contained in  $\bigcup_{i=1}^n U_i$ ; this exists by hypothesis (think about this). Then  $\{x_k\}_{k \geq 1} \subset f^{-1}(K)$  is a sequence with no convergent subsequence: if a subsequence  $\{x_{k_j}\}$  converged to say  $x' \in X$ , then by continuity we would have  $p(x') = y$  and so  $x' = x_i$  for some  $i$ , which tells us that all but finitely many (in particular at least one)  $\{x_{k_j}\}$  lies in  $U_i$ , which is a contradiction.<sup>3</sup>

■

*Remark 5.* Note that if we assume that  $U$  is nonempty and  $V$  is connected, then the map  $f : U \rightarrow V$  is necessarily surjective (and so  $n \geq 1$  in the above). This follows because then the image  $f(U)$  is open (it's the image of a local homeomorphism), closed (by Lemma 0.0.5), and nonempty (since  $U$  is nonempty); therefore, by connectedness, we would conclude that  $f(U) = V$ . In general, surjectivity of  $f$  is neither a necessary nor sufficient condition for it to be a covering map.

*Remark 6.* A lot of the submissions failed to do the final step, and ended up showing the false statement that a local homeomorphism with finite fibers (even between open subsets of the complex plane) is a covering map. A counterexample to this statement would be the inclusion  $\mathbb{C}^\times \hookrightarrow \mathbb{C}$ . It is instructive to see what goes wrong in the above argument for this example. This is false also if we add the word “surjective”; here's a surjective continuous map between open subsets of the plane with finite fibers that is not a covering map: let  $\Delta(a; r) = \{z \in \mathbb{C} : |z - a| < r\}$  for  $a \in \mathbb{C}, r \in \mathbb{R}_{>0}$ , and take  $U = \Delta(0; 1) \setminus \{0\} \cup \Delta(3; 1)$  and  $V := \Delta(0; 1)$ , with the map  $f : U \rightarrow V$  given by the identity on the first (punctured) disc and  $z \mapsto z - 3$  on the second disc.

---

<sup>2</sup>Or at least needs additional hypotheses like first countability and the equivalence of compactness and sequential compactness, etc....

<sup>3</sup>This reasoning shows more generally that a local homeomorphism between Hausdorff spaces with fibers of locally constant finite cardinality is a covering map.

---

**Q5.** Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be given by a polynomial of degree 2 or more. Let

$$V_1 = \{f(z) : f'(z) = 0\} \subset \mathbb{C}$$

be the set of critical values of  $f$ , let  $V_0 = f^{-1}(V_1)$  and let  $U_i = \mathbb{C} \setminus V_i$  for  $i = 0, 1$ . Prove that  $f : U_0 \rightarrow U_1$  is a covering map.

*Proof.* We show that  $f$  is a proper local homeomorphism and use the previous result.

First note that if  $f : \mathbb{C} \rightarrow \mathbb{C}$  is a polynomial map, then it is proper because if  $\{z_n\}$  is any sequence escaping to infinity, then so is  $(f(z_n))$ , because  $f$  is eventually dominated by its leading term, for which this statement is clearly true (this needs to be more precise in your solution). Alternatively, we may use the Heine-Borel theorem to establish this fact. Next, note that if  $f : X \rightarrow Y$  is a proper map between topological spaces and  $V \subset Y$  any subspace, then

$$f|_{X \setminus f^{-1}(V)} : X \setminus f^{-1}(V) \rightarrow Y \setminus V$$

is also proper (why?). In our situation, we can take  $X = Y = \mathbb{C}$ ,  $V = V_1$ , and  $f$  the given polynomial map to conclude that  $f : U_0 \rightarrow U_1$  is proper.

Under the identification of  $\mathbb{C} \cong \mathbb{R}^2$ , the smooth map  $f$  at the point  $z$  has Jacobian matrix

$$J_z f = \begin{bmatrix} \partial_x u & \partial_x v \\ \partial_y u & \partial_y v \end{bmatrix} = \begin{bmatrix} \partial_x u & \partial_x v \\ -\partial_x v & \partial_x u \end{bmatrix},$$

where in the second equality we have used the Cauchy-Riemann equations. It follows that the Jacobian determinant

$$\det J_z f = (\partial_x u)^2 + (\partial_x v)^2 = |f'(z)|^2.$$

By definition, we have for each  $z \in U_0$  that  $f'(z) \neq 0$ , which tells us that the Jacobian matrix  $J_z f$  of  $f$  at  $z$  is invertible. It follows from the Inverse Function Theorem that  $f$  is a  $C^\infty$ -diffeomorphism in a neighborhood of  $z$ . This shows that  $f : U_0 \rightarrow U_1$  is a local homeomorphism.

Another way of showing that  $f$  is a local homeomorphism whenever  $f'(z) \neq 0$  (without resorting to  $x, y$  coordinates) is to note that for any smooth map  $g$  the total differential  $dg$  evaluated at a tangent vector  $v \in T_z U_0$  is given by

$$dg(v) = \frac{\partial g}{\partial z} v + \frac{\partial g}{\partial \bar{z}} \bar{v}.$$

In particular, if  $f$  is holomorphic and so  $\partial f / \partial \bar{z} = 0$ , this then reduces to

$$df(v) = f'(z)v,$$

so that if  $f'(z) \neq 0$ , then this is an isomorphism, and we may apply the Inverse Function Theorem as before. ■

*Remark 7.* In fact,  $f|_{U_0}$  is a local biholomorphism. This follows from the fact that the inverse of an invertible holomorphic function still satisfies the Cauchy-Riemann equations, and is hence holomorphic. This can be proved in one dimension by the Cauchy integral formula (how?).

*Remark 8.* A lot of students applied the Heine-Borel Theorem incorrectly. A metric space  $X$  is said to be a Heine-Borel space or have the Heine-Borel property if a subset  $K \subset X$  is compact iff it is closed and bounded. For any metric space, one implication is always true (which one? why?); it's the other implication that defines a Heine-Borel space. The Heine-Borel Theorem states that  $\mathbb{R}^n$  for any  $n \geq 0$  is a Heine-Borel space. It is, however, not true above (for any  $f$ ) that  $U_1$  is a Heine-Borel space (why?). Therefore, you cannot simply quote the Heine-Borel theorem and that the preimage of bounded subsets is bounded under  $f$  to conclude that  $f : U_0 \rightarrow U_1$  is proper; this needs, the additional argument of the sort made above, i.e. that properness descends when restricting the codomain.



---

**Q6.** Given an example where  $U_0/U_1$  is a normal (or Galois) covering, i.e. where  $f_*(\pi_1(U_0))$  is a normal subgroup of  $\pi_1(U_1)$ .

*Solution.* Many solutions are possible. One possible line of approach: this is automatically the case if  $\pi_1(U_1)$  is abelian, which happens iff  $V_1$  is a single point (proof: exercise). Therefore, it suffices to take  $f(z) = z^n$  for any  $n \geq 2$ .

*Remark 9.* For an example where the covering  $f : U_0 \rightarrow U_1$  is not a Galois cover, consider  $f(z) = z^3 - 3z$ . In this case,  $\pi_1(U_1)$  is the free group on two generators, say  $\alpha$  and  $\beta$ , and (by taking 0 as the basepoint for both domain and codomain), the subgroup  $f_*\pi_1(U_0)$  is  $\langle \alpha^2, \beta^2, \alpha\beta\alpha, \beta\alpha\beta \rangle$ , which is not a normal subgroup. (Exercise: justify all of these assertions. What cover of the doubly-punctured plane, or equivalently the figure-8, does this correspond to? We'll talk more about this cover in class.)

*Remark 10.* Can you classify all covers of the multiply-punctured complex plane (or equivalently Riemann sphere) that arise in this way?

## References

- [1] L. V. Ahlfors, *Complex Analysis*. International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., third ed., 1979.