

## Problem Set # 2

M392C: Bordism Old and New

Due: November 6, 2012

- (a) Recall from (14.4) in the lecture notes that a diffeomorphism  $f: Y \rightarrow Y$  of a closed manifold  $Y$  determines a bordism  $X_f$ . Let  $f_0, f_1$  be diffeomorphisms. Prove that  $X_{f_0}$  is diffeomorphic to  $X_{f_1}$  (as bordisms) if and only if  $f_0$  is pseudoisotopic to  $f_1$ .

(b) Find a manifold  $Y$  and diffeomorphisms  $f_0, f_1: Y \rightarrow Y$  which are pseudoisotopic but not isotopic.
- (a) Let  $S$  be a set with composition laws  $\circ_1, \circ_2: S \times S \rightarrow S$  and distinguished element  $1 \in S$ . Assume (i)  $1$  is an identity for both  $\circ_1$  and  $\circ_2$ ; and (ii) for all  $s_1, s_2, s_3, s_4 \in S$  we have

$$(s_1 \circ_1 s_2) \circ_2 (s_3 \circ_1 s_4) = (s_1 \circ_2 s_3) \circ_1 (s_2 \circ_2 s_4).$$

Prove that  $\circ_1 = \circ_2$  and that this common operation is commutative and associative.

- (b) Let  $C$  be a symmetric monoidal category. Apply (a) to  $C(1, 1)$ , where  $1 \in C$  is the tensor unit.
- Let  $y \in C$  be a dualizable object in a symmetric monoidal category, and suppose  $(y^\vee, c, e)$  and  $(\tilde{y}^\vee, \tilde{c}, \tilde{e})$  are two sets of duality data. Prove there is a unique map  $(y^\vee, c, e) \rightarrow (\tilde{y}^\vee, \tilde{c}, \tilde{e})$ .
- For each of the following symmetric monoidal categories determine all of the dualizable objects.

  - (Top,  $\amalg$ ), the category of topological spaces and continuous maps under disjoint union.
  - (Ab,  $\oplus$ ), the category of abelian groups and homomorphisms under direct sum.
  - (Mod $_R$ ,  $\otimes$ ), the category of  $R$ -modules and homomorphisms under tensor product, where  $R$  is a commutative ring.
  - (Set,  $\times$ ), the category of sets and functions under Cartesian product.
- Recall that every category  $C$  has an associated groupoid  $|C|$  obtained from  $C$  by inverting all of the arrows. What is  $|Bord_{\langle 1, 2 \rangle}|$ ?  $|Bord_{\langle 1, 2 \rangle}^{\text{Spin}}|$ ? What are all Vect $_C$ -valued invertible topological quantum field theories with domain  $Bord_{\langle 1, 2 \rangle}$ ?  $Bord_{\langle 1, 2 \rangle}^{\text{Spin}}$ ?
- Fix a finite group  $G$ . Let  $C$  denote the groupoid  $G//G$  of  $G$  acting on itself by conjugation. Let  $D$  denote the groupoid of principal  $G$ -bundles over  $S^1$ . (A principal  $G$ -bundle is a regular, or Galois, cover with group  $G$ .) Prove that  $C$  and  $D$  are equivalent groupoids. You should spell out precisely what these groupoids are.

7. Explain why each of the following fails to be a natural map  $\eta: F \rightarrow G$  of symmetric monoidal functors  $F, G: C \rightarrow D$ .

(a)  $F, G$  are the identity functor on  $\text{Vect}_k$  for some field  $k$ , and  $\eta(V): V \rightarrow V$  is multiplication by 2 for each vector space  $V$ .

(b)  $C, D$  are the category of algebras over a field  $k$ , the functor  $F$  maps  $A \mapsto A \otimes A$ , the functor  $G$  is the identity, and  $\eta(A): A \otimes A \rightarrow A$  is multiplication.

8. In this problem you construct a simple TQFT  $F: \text{Bord}_{\langle 0,1 \rangle} \rightarrow \text{Vect}_{\mathbb{Q}}$ . For any manifold  $M$  let  $\mathcal{C}(M)$  denote the groupoid of principal  $G$ -bundles over  $M$ , as in Problem 6.

(a) For a compact 0-manifold  $Y$ , define  $F(Y)$  as the vector space of functions  $\mathcal{C}(Y) \rightarrow \mathbb{Q}$ . Say what you mean by such functors on a groupoid.

(b) For a *closed* 1-manifold  $X$  define

$$F(X) = \sum_{[P] \in \pi_0 \mathcal{C}(X)} \frac{1}{\# \text{Aut}(P)},$$

where the sum is over equivalence classes of principal  $G$ -bundles. Extend this to all bordisms  $X: Y_0 \rightarrow Y_1$ .

(c) Check that  $F$  is a symmetric monoidal functor.

(d) Calculate  $F$  on a set of duality data for the point  $\text{pt} \in \text{Bord}_{\langle 0,1 \rangle}$ . Use it to compute  $F(S^1)$ .

9. Fix a nonzero number  $\lambda \in \mathbb{C}$ . Construct an invertible TQFT  $F: \text{Bord}_{\langle 1,2 \rangle}^{SO} \rightarrow \text{Vect}_{\mathbb{C}}$  such that for a closed 2-manifold  $X$  we have  $F(X) = \lambda^{\chi(X)}$ , where  $\chi(X)$  is the Euler characteristic. Can you extend to the bordism category  $\text{Bord}_{\langle 1,2 \rangle}$ ?