

Math 155: Designs and Groups

Homework Assignment #3 (10 February 2010): Inclusion-Exclusion, designs, and a bit about regular graphs

This problem set is due Friday, Feb. 19 in class.

- [CvL p.26 #10] Use the method of “intersection triangles” to prove that if Y is a set of 5 of the points of a 3-(10,4,1) design then Y contains no block if and only if the complement of Y contains no block.
- [CvL p.25 #5] Let x_1, x_2, \dots, x_{t+1} be points of a t -(v, k, λ) design $\mathcal{D} = (X, \mathcal{B})$, and let μ be the number of blocks containing those $t + 1$ points. Use the Principle of Inclusion and Exclusion to show that there exists a number F , depending only on the parameters t, v, k, λ of \mathcal{D} , such that the number of blocks containing none of x_1, x_2, \dots, x_{t+1} is $F + (-1)^{t+1}\mu$.
- Solve Problem #18 on p.27–28 of the textbook. [For part (a), if you know about finite differences then you may be able to recognize an alternative proof. Part (b) you may know already; some 20 years ago it also appeared on our Qualifying Exam for graduate students.]
- Solve Problem #16 on p.27 (this involves arguments similar to those we’ll use in analyzing arcs and ovals).
- Let k be the finite field of q elements where $q \equiv 3 \pmod{4}$, and let G be the group of $(q^2 - q)/2$ permutations of k of the form $x \mapsto a^2x + b$ where $a, b \in k$ and $a \neq 0$. Prove that while G is not doubly transitive, G does act transitively (indeed simply transitively) on *unordered* pairs of elements of k . Use this to give an alternative verification of Paley’s construction of a Hadamard 2-design.
- Let F be a perfect (but not necessarily finite) field¹ of characteristic 2, and $C \subset \mathbf{P}^2(F)$ the conic $xz = y^2$, i.e. $(x : y : z) = (r^2 : rs : s^2)$ for $(r : s) \in \mathbf{P}^1(F)$.² Determine for each point on C the tangent through C , and find the point $P \in \mathbf{P}^2(F)$ at which all the tangents meet. Check algebraically that any point $P' \neq P$ in the projective plane lies on a unique tangent. [Our combinatorial techniques don’t apply when F is infinite.]
- Now let F be a finite field of 2^n elements, and let d be an integer relatively prime to n . Prove that the subset of $\mathbf{P}^2(F)$ consisting of $(1 : 0 : 0)$, $(0 : 1 : 0)$, and all points of the form $(x : y : z) = (a^{2^d} : a : 1)$ ($a \in F$) is a hyperoval, and is an extended conic (a conic together with its center) if and only if $d \equiv \pm 1 \pmod{n}$. Conclude that if $n = 5$ or $n > 6$ then $\mathbf{P}^2(F)$ contains hyperovals that are not extended conics.
- Recall that the “girth” of a graph is the length of its shortest cycle. Prove that a regular graph of degree d and girth 6 has at least $2(d^2 - d + 1)$ vertices, with equality possible if and only if there is a finite projective plane of order $d - 1$. For instance there is up to isomorphism a unique cubic graph³ of girth 6 on 14 vertices (the “Heawood graph”); what is its automorphism group? Show that this is also the graph obtained by tiling the torus with seven pairwise adjacent hexagons.

¹Recall that a field k called is *perfect* if every finite extension of k is separable. This condition is automatic if k has characteristic zero, while in characteristic p it is equivalent to the condition that $k = k^p$, i.e. every field element is of the form c^p for some $c \in k$ (unique because $c^p - c'^p = (c - c')^p$). This is automatic for finite fields but may fail in general, e.g. $\mathbf{F}_p(X)$ is not perfect since X is not a p -th power.

²Remember that the notation $(x : y : z) = (r^2 : rs : s^2)$ means that there exists $c \in k^*$ such that $(x, y, z) = c(r^2, rs, s^2)$; in general we cannot assume that $(x, y, z) = (r^2, rs, s^2)$ (with $c = 1$).

³As you might remember from our first meeting, “cubic graph” is standard graph theory lingo for “regular graph of degree 3”.