

A “STRANGE” FUNCTIONAL EQUATION FOR EISENSTEIN SERIES AND MIRACULOUS DUALITY ON THE MODULI STACK OF BUNDLES

D. GAITSGORY

ABSTRACT. We show that the failure of the usual Verdier duality on Bun_G leads to a new duality functor on the category of D-modules, and we study its relation to the operation of Eisenstein series.

INTRODUCTION

0.1. Context for the present work.

0.1.1. This paper arose in the process of developing what V. Drinfeld calls *the geometric theory of automorphic functions*. I.e., we study *sheaves* on the moduli stack Bun_G of principal G -bundles on a curve X . Here and elsewhere in the paper, we fix an algebraically closed ground field k , and we let G be a reductive group and X a smooth and complete curve over k .

In the bulk of the paper we will take k to be of characteristic 0, and by a “sheaf” we will understand an object of the derived category of D-modules. However, with appropriate modifications, our results apply also to ℓ -adic sheaves, or any other reasonable sheaf-theoretic situation.

Much of the motivation for the study of sheaves on Bun_G comes from the so-called *geometric Langlands program*. In line with this, the main results of this paper have a transparent meaning in terms of this program, see Sect. 0.2. However, one can also view them from the perspective of the classical theory of automorphic functions (rather, we will see phenomena that so far have not been studied classically).

0.1.2. *Constant term and Eisenstein series functors.* To explain what is done in this paper we will first recall the main result of [DrGa3].

Let $P \subset G$ be a parabolic subgroup with Levi quotient M . The diagram of groups

$$G \leftarrow P \rightarrow M$$

gives rise to a diagram of stacks

$$(0.1) \quad \begin{array}{ccc} & \text{Bun}_P & \\ \swarrow p & & \searrow q \\ \text{Bun}_G & & \text{Bun}_M. \end{array}$$

Using this diagram as “pull-push”, one can write down several functors connecting the categories of D-modules on Bun_G and Bun_M , respectively. By analogy with the classical theory of automorphic functions, we call the functors going from Bun_M to Bun_G “Eisenstein series”, and the functors going from Bun_G to Bun_M “constant term”.

Date: May 9, 2016.

Namely, we have

$$\begin{aligned} \mathrm{Eis}_! &:= \mathfrak{p}_! \circ \mathfrak{q}^*, & \mathrm{D}\text{-mod}(\mathrm{Bun}_M) &\rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G), \\ \mathrm{Eis}_* &:= \mathfrak{p}_* \circ \mathfrak{q}^!, & \mathrm{D}\text{-mod}(\mathrm{Bun}_M) &\rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G), \\ \mathrm{CT}_! &:= \mathfrak{q}_! \circ \mathfrak{p}^*, & \mathrm{D}\text{-mod}(\mathrm{Bun}_G) &\rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M), \\ \mathrm{CT}_* &:= \mathfrak{q}_* \circ \mathfrak{p}^!, & \mathrm{D}\text{-mod}(\mathrm{Bun}_G) &\rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M). \end{aligned}$$

Note that unlike the classical theory, where there is only one pull-back and one push-forward for functions, for sheaves there are two options: $!$ and $*$, for both pull-back and push-forward. The interaction of these two options is one way to look at what this paper is about.

Among the above functors, there are some obvious adjoint pairs: $\mathrm{Eis}_!$ is the left adjoint of CT_* , and $\mathrm{CT}_!$ is the left adjoint of Eis_* .

In addition to this, the following, perhaps a little unexpected, result was proved in [DrGa3]:

Theorem 0.1.3. *The functors $\mathrm{CT}_!$ and CT_*^- are canonically isomorphic.*

In the statement of the theorem the superscript “ $-$ ” means the constant term functor taken with respect to the *opposite* parabolic P^- (note that the Levi quotients of P and P^- are canonically identified).

Our goal in the present paper is to understand what implication the above-mentioned isomorphism

$$\mathrm{CT}_! \simeq \mathrm{CT}_*^-$$

has for the Eisenstein series functors $\mathrm{Eis}_!$ and Eis_* . The conclusion will be what we will call a “strange” functional equation (0.9), explained below.

In order to explain what the “strange” functional equation does, we will need to go a little deeper into what one may call the “functional-analytic” aspects of the study of Bun_G .

0.1.4. *Verdier duality on stacks.* The starting point for the “analytic” issues that we will be dealing with is that the stack Bun_G is *not quasi-compact* (this is parallel to the fact that in the classical theory, the automorphic space is not compact, leading to a host of interesting analytic phenomena). The particular phenomenon that we will focus on is the absence of the usual Verdier duality functor, and what replaces it.

First off, it is well-known (see, e.g., [DrGa2, Sect. 2]) that if \mathcal{Y} is an arbitrary reasonable¹ quasi-compact algebraic stack, then the category $\mathrm{D}\text{-mod}(\mathcal{Y})$ is compactly generated and naturally self-dual.

Perhaps, the shortest way to understand the meaning of self-duality is that the subcategory $\mathrm{D}\text{-mod}(\mathcal{Y})^c \subset \mathrm{D}\text{-mod}(\mathcal{Y})$ consisting of compact objects carries a canonically defined contravariant self-equivalence, called Verdier duality. A more flexible way of interpreting the same phenomenon is an equivalence, denoted $\mathbf{D}_{\mathcal{Y}}$, between $\mathrm{D}\text{-mod}(\mathcal{Y})$ and its *dual* category $\mathrm{D}\text{-mod}(\mathcal{Y})^\vee$ (we refer the reader to [DrGa1, Sect. 1], where the basics of the notion of duality for DG categories are reviewed).

Let us now remove the assumption that \mathcal{Y} be quasi-compact. Then there is another geometric condition, called “truncatability” that ensures that $\mathrm{D}\text{-mod}(\mathcal{Y})$ is compactly generated (see [DrGa2, Definition 4.1.1], where this notion is introduced). We remark here that the goal of the paper [DrGa2] was to show that the stack Bun_G is truncatable. The reader who is not familiar with this notion is advised to ignore it on the first pass.

¹The word “reasonable” here does not have a technical meaning; the technical term is “QCA”, which means that the automorphism group of any field-valued point is affine.

Thus, let us assume that \mathcal{Y} is truncatable. However, there still is no obvious replacement for Verdier duality: extending the quasi-compact case, one can define a functor

$$(\mathrm{D}\text{-mod}(\mathcal{Y})^c)^{\mathrm{op}} \rightarrow \mathrm{D}\text{-mod}(\mathcal{Y}),$$

but it no longer lands in $\mathrm{D}\text{-mod}(\mathcal{Y})^c$ (unless \mathcal{Y} is a disjoint union of quasi-compact stacks). In the language of dual categories, we have a functor

$$\mathrm{Ps}\text{-Id}_{\mathcal{Y},\mathrm{naive}} : \mathrm{D}\text{-mod}(\mathcal{Y})^\vee \rightarrow \mathrm{D}\text{-mod}(\mathcal{Y}),$$

but it is no longer an equivalence.²

In particular, the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,\mathrm{naive}}$ is *not* an equivalence, unless G is a torus.

0.1.5. *The pseudo-identity functor.* To potentially remedy this, V. Drinfeld suggested another functor, denoted

$$\mathrm{Ps}\text{-Id}_{\mathcal{Y},!} : \mathrm{D}\text{-mod}(\mathcal{Y})^\vee \rightarrow \mathrm{D}\text{-mod}(\mathcal{Y}),$$

see [DrGa2, Sect. 4.4.8] or Sect. 3.1 of the present paper.

Now, it is not true that for all truncatable stacks \mathcal{Y} , the functor $\mathrm{Ps}\text{-Id}_{\mathcal{Y},!}$ is an equivalence. In [DrGa2] the stacks for which it is an equivalence are called “miraculous”.

We can now formulate the main result of this paper (conjectured by V. Drinfeld):

Theorem 0.1.6. *The stack Bun_G is miraculous.*

We repeat that the above theorem says that the canonically defined functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,!}$ defines an identification of $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ and its dual category. Equivalently, it gives rise to a (non-obvious!) contravariant self-equivalence on $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)^c$.

0.1.7. *The “strange” functional equation.* Finally, we can go back and state the “strange” functional equation, which is in fact an ingredient in the proof of Theorem 0.1.6:

Theorem 0.1.8. *We have a canonical isomorphism of functors*

$$\mathrm{Eis}_!^- \circ \mathrm{Ps}\text{-Id}_{\mathrm{Bun}_M,!} \simeq \mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,!} \circ (\mathrm{CT}_*)^\vee.$$

In the Theorem 0.1.8, the functor $(\mathrm{CT}_*)^\vee$ maps

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_M)^\vee \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee$$

and is the *dual* of the functor CT_* . As we shall see in Sect. 1.5, the functor $(\mathrm{CT}_*)^\vee$ is a close relative of the functor Eis_* , introduced earlier.

0.2. Motivation from geometric Langlands. We shall now proceed and describe how the results of this paper fit into the geometric Langlands program. The contents of this subsection play a motivational role only, and the reader not familiar with the objects discussed below can skip this subsection and proceed to Sect. 0.3.

²The category $\mathrm{D}\text{-mod}(\mathcal{Y})^\vee$ and the functor $\mathrm{Ps}\text{-Id}_{\mathcal{Y},\mathrm{naive}}$ will be described explicitly in Sect. 1.2.

0.2.1. *Statement of GLC.* Let us recall the statement of the categorical geometric Langlands conjecture (GLC), according to [AG, Conjecture 10.2.2].

The left-hand (i.e., geometric) side of GLC is the DG category $\mathbf{D}\text{-mod}(\text{Bun}_G)$ of D-modules on the stack Bun_G .

Let \check{G} denote the Langlands dual group of G , and let $\text{LocSys}_{\check{G}}$ denote the (derived) stack of \check{G} -local systems on X . The right-hand (i.e., spectral) side of GLC has to do with (quasi)-coherent sheaves on $\text{LocSys}_{\check{G}}$.

More precisely, in [AG], a certain modification of the DG category $\text{QCoh}(\text{LocSys}_{\check{G}})$ was introduced; we denote it by $\text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}})$. This category is what appears on the spectral side of GLC.

Thus, GLC states the existence of an equivalence

$$(0.2) \quad \mathbb{L}_G : \mathbf{D}\text{-mod}(\text{Bun}_G) \rightarrow \text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}),$$

that satisfies a number of properties that (conjecturally) determine \mathbb{L}_G uniquely.

The property of \mathbb{L}_G , relevant for this paper, is the compatibility of (0.2) with the functor of Eisenstein series, see Sect. 0.2.5 below.

0.2.2. *Interaction of GLC with duality.* A feature of the spectral side crucial for this paper is that the Serre duality functor of [AG, Proposition 3.7.2] gives rise to an equivalence:

$$\mathbf{D}_{\text{LocSys}_{\check{G}}}^{\text{Serre}} : (\text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}))^{\vee} \rightarrow \text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}).$$

(Here, as in Sect. 0.1.4, for a compactly generated category \mathbf{C} , we denote by \mathbf{C}^{\vee} the dual category.)

Hence, if we believe in the existence of an equivalence \mathbb{L}_G of (0.2), there should exist an equivalence

$$(0.3) \quad (\mathbf{D}\text{-mod}(\text{Bun}_G))^{\vee} \simeq \mathbf{D}\text{-mod}(\text{Bun}_G).$$

Now, the pseudo-identity functor $\text{Ps-Id}_{\text{Bun}_G, !}$ mentioned in Sect. 0.1.5 and appearing in Theorem 0.1.6 is exactly supposed to perform this role. More precisely, we can enhance the statement of GLC by specifying how it is supposed to interact with duality:

Conjecture 0.2.3. *The diagram*

$$(0.4) \quad \begin{array}{ccc} \mathbf{D}\text{-mod}(\text{Bun}_G)^{\vee} & \xrightarrow{((\mathbb{L}_G)^{\vee})^{-1}} & (\text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}))^{\vee} \\ & & \downarrow \mathbf{D}_{\text{LocSys}_{\check{G}}}^{\text{Serre}} \\ \text{Ps-Id}_{\text{Bun}_G, !} \downarrow & & \text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}) \\ & & \downarrow \tau \\ \mathbf{D}\text{-mod}(\text{Bun}_G) & \xrightarrow{\mathbb{L}_G} & \text{IndCoh}_{\text{Nilp}_{glob}}(\text{LocSys}_{\check{G}}) \end{array}$$

commutes up to a cohomological shift, where τ denotes the automorphism, induced by the Cartan involution of G .

Remark 0.2.4. Let us comment on the presence of the Cartan involution in Conjecture 0.2.3. In fact, it can be seen already when G is a torus T , in which case τ is the inversion automorphism.

Indeed, we let \mathbb{L}_T be the Fourier-Mukai equivalence, and Conjecture 0.2.3 is known to hold.

0.2.5. *Interaction of GLC with Eisenstein series.* Let us recall (following [AG, Conjecture 12.2.9] or [Gal, Sect. 6.4.5]) how the equivalence \mathbb{L}_G is supposed to be compatible with the functor(s) of Eisenstein series.

For a (standard) parabolic $P \subset G$, let \check{P} be the corresponding parabolic in \check{G} . Consider the diagram

$$(0.5) \quad \begin{array}{ccc} & \text{LocSys}_{\check{P}} & \\ \mathfrak{p}_{\text{spec}} \swarrow & & \searrow \mathfrak{q}_{\text{spec}} \\ \text{LocSys}_{\check{G}} & & \text{LocSys}_{\check{M}}. \end{array}$$

We define the functors of spectral Eisenstein series and constant term

$$\text{Eis}_{\text{spec}} : \text{IndCoh}(\text{LocSys}_{\check{M}}) \rightarrow \text{IndCoh}(\text{LocSys}_{\check{G}}), \quad \text{Eis}_{\text{spec}} := (\mathfrak{p}_{\text{spec}})_* \circ (\mathfrak{q}_{\text{spec}})^*,$$

$$\text{CT}_{\text{spec}} : \text{IndCoh}(\text{LocSys}_{\check{G}}) \rightarrow \text{IndCoh}(\text{LocSys}_{\check{M}}), \quad \text{CT}_{\text{spec}} := (\mathfrak{q}_{\text{spec}})_* \circ (\mathfrak{p}_{\text{spec}})^!$$

see [AG, Sect. 12.2.1] for more details. The functors $(\text{Eis}_{\text{spec}}, \text{CT}_{\text{spec}})$ form an adjoint pair.

Remark 0.2.6. In [AG, Conjecture 12.2.9] a slightly different version of the functor Eis_{spec} is given, where instead of the functor $(\mathfrak{q}_{\text{spec}})^*$ we use $(\mathfrak{q}_{\text{spec}})^!$. The difference between these two functors is given by tensoring by a graded line bundle on $\text{LocSys}_{\check{M}}$; this is due to the fact that the morphism $\mathfrak{q}_{\text{spec}}$ is *Gorenstein*. This difference will be immaterial for the purposes of this paper.

The compatibility of the geometric Langlands equivalence of (0.2) with Eisenstein series reads (see [AG, Conjecture 12.2.9]):

Conjecture 0.2.7. *The diagram*

$$(0.6) \quad \begin{array}{ccc} \text{D-mod}(\text{Bun}_G) & \xrightarrow{\mathbb{L}_G} & \text{IndCoh}_{\text{NilP}_{\text{glob}}}(\text{LocSys}_{\check{G}}) \\ \text{Eis}_! \uparrow & & \uparrow \text{Eis}_{\text{spec}} \\ \text{D-mod}(\text{Bun}_M) & \xrightarrow{\mathbb{L}_M} & \text{IndCoh}_{\text{NilP}_{\text{glob}}}(\text{LocSys}_{\check{M}}) \end{array}$$

commutes up to an automorphism of $\text{IndCoh}_{\text{NilP}_{\text{glob}}}(\text{LocSys}_{\check{M}})$, given by tensoring with a certain canonically defined graded line bundle on $\text{LocSys}_{\check{M}}$.

0.2.8. *Recovering the “strange” functional equation.* Let us now analyze what the combination of Conjectures 0.2.3 and 0.2.7 says about the interaction of the functor $\text{Ps-Id}_{\text{Bun}_G, !}$ with $\text{Eis}_!$. The conclusion that we will draw will amount to Theorem 0.1.8 of the present paper (the reader may safely choose to skip the derivation that follows).

First, passing to the right adjoint and then dual functors in (0.6), we obtain a diagram

$$(0.7) \quad \begin{array}{ccc} \text{D-mod}(\text{Bun}_G)^\vee & \xrightarrow{(\mathbb{L}_G^\vee)^{-1}} & (\text{IndCoh}_{\text{NilP}_{\text{glob}}}(\text{LocSys}_{\check{G}}))^\vee \\ (\text{CT}_*)^\vee \uparrow & & \uparrow (\text{CT}_{\text{spec}})^\vee \\ \text{D-mod}(\text{Bun}_M)^\vee & \xrightarrow{(\mathbb{L}_M^\vee)^{-1}} & (\text{IndCoh}_{\text{NilP}_{\text{glob}}}(\text{LocSys}_{\check{M}}))^\vee \end{array}$$

that commutes up to a tensoring by a graded line bundle on $\text{LocSys}_{\check{M}}$.

Next, we note that the diagram

$$(0.8) \quad \begin{array}{ccc} (\mathrm{IndCoh}_{\mathrm{NilP}_{glob}}(\mathrm{LocSys}_{\check{G}}))^{\vee} & \xrightarrow{\mathbf{D}_{\mathrm{LocSys}_{\check{G}}}^{\mathrm{Serre}}} & \mathrm{IndCoh}_{\mathrm{NilP}_{glob}}(\mathrm{LocSys}_{\check{G}}) \\ (\mathrm{CT}_{\mathrm{spec}})^{\vee} \uparrow & & \mathrm{Eis}_{\mathrm{spec}} \uparrow \\ (\mathrm{IndCoh}_{\mathrm{NilP}_{glob}}(\mathrm{LocSys}_{\check{M}}))^{\vee} & \xrightarrow{\mathbf{D}_{\mathrm{LocSys}_{\check{M}}}^{\mathrm{Serre}}} & \mathrm{IndCoh}_{\mathrm{NilP}_{glob}}(\mathrm{LocSys}_{\check{M}}) \end{array}$$

also commutes up to a tensoring by a graded line bundle on $\mathrm{LocSys}_{\check{M}}$, see Remark 0.2.6.

Now, juxtaposing the diagrams (0.6), (0.7), (0.8) with the diagrams (0.4) for the groups G and M respectively, we obtain a commutative diagram:

$$(0.9) \quad \begin{array}{ccc} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^{\vee} & \xrightarrow{\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !}} & \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \\ (\mathrm{CT}_{\star})^{\vee} \uparrow & & \uparrow \tau_G \circ \mathrm{Eis}_! \circ \tau_M \\ \mathrm{D}\text{-mod}(\mathrm{Bun}_M)^{\vee} & \xrightarrow{\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_M, !}} & \mathrm{D}\text{-mod}(\mathrm{Bun}_M). \end{array}$$

Notice now that $\tau_G \circ \mathrm{Eis}_! \circ \tau_M \simeq \mathrm{Eis}_!^{-}$, so the commutative diagram (0.9) recovers the isomorphism of Theorem 0.1.8.

0.3. The usual functional equation. As was mentioned above, we view the commutativity of the diagram (0.9) as a kind of “strange” functional equation, hence the title of this paper.

Let us now compare it to the usual functional equation of [BG, Theorem 2.1.8].

0.3.1. In *loc.cit.* one considered the case of $P = B$, the Borel subgroup and hence $M = T$, the abstract Cartan. We consider the full subcategory

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_T)^{\mathrm{reg}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_T),$$

defined as in [BG, Sect. 2.1.7]. This is a full subcategory that under the Fourier-Mukai equivalence

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_T) \simeq \mathrm{QCoh}(\mathrm{LocSys}_{\check{T}})$$

corresponds to

$$\mathrm{QCoh}(\mathrm{LocSys}_{\check{T}}^{\mathrm{reg}}) \hookrightarrow \mathrm{QCoh}(\mathrm{LocSys}_{\check{T}}),$$

where $\mathrm{LocSys}_{\check{T}}^{\mathrm{reg}} \subset \mathrm{LocSys}_{\check{T}}$ is the open locus of $\mathrm{LocSys}_{\check{T}}$ consisting of those \check{T} -local systems that for every root α of \check{T} induce a non-trivial local system for \mathbb{G}_m .

Instead of the functor $\mathrm{Eis}_!$, or the functor that we introduce as $\mathrm{Eis}_{\star} := \mathfrak{p}_{\star} \circ \mathfrak{q}^!$ (see Sect. 1.1.6), an intermediate version was considered in [BG, Sect. 2.1], which we will denote here by $\mathrm{Eis}_{!^{\star}}$. The definition of $\mathrm{Eis}_{!^{\star}}$ uses the compactification of the morphism \mathfrak{p} , introduced in [BG, Sect. 1.2]:

$$\begin{array}{ccc} \mathrm{Bun}_B & \xrightarrow{r} & \overline{\mathrm{Bun}}_B \\ & \searrow \bar{\mathfrak{p}} & \swarrow \bar{\mathfrak{q}} \\ & \mathrm{Bun}_G & \mathrm{Bun}_T. \end{array}$$

The assertion of [BG, Theorem 2.1.8] (for the longest element of the Weyl group) is:

Theorem 0.3.2. *The following diagram of functors*

$$\begin{array}{ccc}
 \mathrm{D}\text{-mod}(\mathrm{Bun}_G) & \xrightarrow{\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}} & \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \\
 \mathrm{Eis}_{!,*} \uparrow & & \uparrow \mathrm{Eis}_{!,*}^- \\
 \mathrm{D}\text{-mod}(\mathrm{Bun}_T) & & \mathrm{D}\text{-mod}(\mathrm{Bun}_T) \\
 \uparrow & & \uparrow \\
 \mathrm{D}\text{-mod}(\mathrm{Bun}_T)^{\mathrm{reg}} & \xrightarrow{\rho\text{-shift}} & \mathrm{D}\text{-mod}(\mathrm{Bun}_T)^{\mathrm{reg}},
 \end{array}$$

*commutes up to a cohomological shift, where ρ -shift is the functor of translation by the point $2\rho(\Omega_X)$.*³

Theorem 0.3.2 is a geometric analog of the usual functional equation for Eisenstein series in the theory of automorphic functions.

0.3.3. Let us emphasize the following points of difference between Theorems 0.1.8 and 0.3.2:

- Theorem 0.1.8 compares the functors $\mathrm{Eis}_{!,*}^-$ and $(\mathrm{CT}_*)^\vee$ that take values in different categories, i.e., $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ vs. $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee$, whereas in Theorem 0.3.2 both $\mathrm{Eis}_{!,*}$ and $\mathrm{Eis}_{!,*}^-$ map to $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)$.
- The vertical arrows in Theorem 0.1.8 use geometrically different functors, while in Theorem 0.3.2 these are functors of the same nature, i.e., $\mathrm{Eis}_{!,*}$ and $\mathrm{Eis}_{!,*}^-$.
- The upper horizontal arrow Theorem 0.1.8 is the geometrically non-trivial functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,!}$, while in Theorem 0.3.2 it is the identity functor.
- The lower horizontal arrow in Theorem 0.1.8 for $M = T$ is isomorphic to the identity functor, up to a cohomological shift, while in Theorem 0.3.2 we have the functor of ρ -shift.
- The commutation in 0.1.8 takes place on all of $\mathrm{D}\text{-mod}(\mathrm{Bun}_T)$, whereas in Theorem 0.3.2, it only takes place on $\mathrm{D}\text{-mod}(\mathrm{Bun}_T)^{\mathrm{reg}}$.

0.4. **Interaction with cuspidality.** There is yet one more set of results contained in this paper, which has to do with the notion of cuspidality.

0.4.1. The cuspidal subcategories

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \text{ and } (\mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee)_{\mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee$$

are defined as right-orthogonals of the subcategories generated by the essential images of the functors

$$\mathrm{Eis}_{!,*} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \text{ and } (\mathrm{CT}_*)^\vee : \mathrm{D}\text{-mod}(\mathrm{Bun}_M)^\vee \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee,$$

respectively, for all *proper* parabolics P of G .

0.4.2. Let us return to the setting of Sect. 0.1.4 are recall the “naive” functor

$$\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,\mathrm{naive}} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G),$$

see Sect. 0.1.4.

As was mentioned in *loc.cit.*, the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G,\mathrm{naive}}$ fails to be an equivalence unless G is a torus. However, in Theorem 2.2.7 we show:

³Here $2\rho : \mathbb{G}_m \rightarrow T$ is the coweight equal to the sum of positive coroots, and $\Omega_X \in \mathrm{Pic}(X) = \mathrm{Bun}_{\mathbb{G}_m}$ is the canonical line bundle on X .

Theorem 0.4.3. *The restriction of the functor $\text{Ps-Id}_{\text{Bun}_G, \text{naive}}$ to*

$$(\text{D-mod}(\text{Bun}_G)^\vee)_{\text{cusp}} \subset \text{D-mod}(\text{Bun}_G)^\vee$$

defines an equivalence

$$(\text{D-mod}(\text{Bun}_G)^\vee)_{\text{cusp}} \rightarrow \text{D-mod}(\text{Bun}_G)_{\text{cusp}}.$$

One can view Theorem 0.4.3 as expressing the fact that the objects of $(\text{D-mod}(\text{Bun}_G)^\vee)_{\text{cusp}}$ and $\text{D-mod}(\text{Bun}_G)_{\text{cusp}}$ are “supported” on quasi-compact open substacks (see Propositions 2.3.2 and 2.3.4 for a precise statement).

0.4.4. In addition, in Corollary 3.3.2 we show:

Theorem 0.4.5. *The functors*

$$\text{Ps-Id}_{\text{Bun}_G, \text{naive}} \big|_{(\text{D-mod}(\text{Bun}_G)^\vee)_{\text{cusp}}} \quad \text{and} \quad \text{Ps-Id}_{\text{Bun}_G, !} \big|_{(\text{D-mod}(\text{Bun}_G)^\vee)_{\text{cusp}}}$$

are isomorphic up to a cohomological shift.

Theorem 0.4.5 is responsible for the fact that previous studies in geometric Langlands correspondence that involved only cuspidal objects did not see the appearance of the functor $\text{Ps-Id}_{\text{Bun}_G, !}$ and one could afford to ignore the difference between $\text{D-mod}(\text{Bun}_G)$ and $\text{D-mod}(\text{Bun}_G)^\vee$. In other words, usual manipulations with Verdier duality on cuspidal objects did not produce wrong results.

0.5. Structure of the paper.

0.5.1. In Sect. 1 we recall the setting of [DrGa3], and list the various Eisenstein series and constant term functors for the usual category $\text{D-mod}(\text{Bun}_G)$. In fact there are two adjoint pairs: $(\text{Eis}_!, \text{CT}_*)$ and $(\text{CT}_!^\mu, \text{Eis}_*^\mu)$, where in the latter pair the superscript $\mu \in \pi_1(M) = \pi_0(\text{Bun}_M)$ indicates that we are considering one connected component of Bun_M at a time.

We recall the main result of [DrGa3] that says that the functors CT_* and $\text{CT}_!^-$ are canonically isomorphic.

Next, we consider the category $\text{D-mod}(\text{Bun}_G)_{\text{co}}$, which is nearly tautologically identified with the category that we have so far denoted $\text{D-mod}(\text{Bun}_G)^\vee$, and introduce the corresponding Eisenstein series and constant term functors:

$$(\text{Eis}_{\text{co}, *}, \text{CT}_{\text{co}, ?}) \quad \text{and} \quad (\text{CT}_{\text{co}, *}^\mu, \text{Eis}_{\text{co}, ?}^\mu),$$

where $\text{Eis}_{\text{co}, *} := (\text{CT}_*)^\vee$, $\text{CT}_{\text{co}, *}^\mu := (\text{Eis}_*^\mu)^\vee$.

The functor $\text{CT}_{\text{co}, ?}$ is something that we do not know how to express in terms of the usual functors in the theory of D-modules; it can be regarded as a *non-standard* functor in the terminology of [DrGa2, Sect. 3.3].

A priori, the functor $\text{Eis}_{\text{co}, ?}^\mu$ would also be a non-standard functor. However, the isomorphism $\text{CT}_* \simeq \text{CT}_!^-$ gives rise to an isomorphism

$$\text{Eis}_{\text{co}, ?} \simeq \text{Eis}_{\text{co}, *}^-.$$

0.5.2. In Sect. 2 we recall the definition of the functor

$$\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G),$$

and show that it intertwines the functors $\mathrm{Eis}_{\mathrm{co}, *}$ and Eis_* , and $\mathrm{CT}_{\mathrm{co}, *}$ and CT_* , respectively.

The remainder of this section is devoted to the study of the subcategory

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G),$$

and the proof of Theorem 0.4.3, which says that the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ defines an equivalence from $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$ to $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$.

0.5.3. In Sect. 3 we introduce the functor

$$\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G),$$

and study its behavior vis-à-vis the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$. The relation is expressed by Proposition 3.2.6, whose proof is deferred to [Sch]. Proposition 3.2.6 essentially says that the difference between $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !}$ and $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ can be expressed in terms of the Eisenstein and constant term functors for *proper* parabolics.

We prove Theorem 0.4.5 that says that the functors $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !}$ and $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ are isomorphic (up to a cohomological shift), when evaluated on cuspidal objects.

0.5.4. In Sect. 4 we prove our “strange” functional equation, i.e., Theorem 0.1.8. The proof is basically a formal manipulation from the isomorphism $\mathrm{CT}_* \simeq \mathrm{CT}_1^-$.

Having Theorem 0.1.8, we get control of the behavior of the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !}$ on the Eisenstein part of the category $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$. From here we deduce our main result, Theorem 0.1.6.

0.6. **Conventions.** The conventions in this paper follow those adopted in [DrGa2]. We refer the reader to *loc.cit.* for a review of the theory of DG categories (freely used in this paper), and the theory of D-modules on stacks.

0.7. **Acknowledgements.** The author would like to express his gratitude to V. Drinfeld. The main results of this paper, in particular, Theorem 0.1.8, were obtained in collaboration with him. Even more crucially, the very idea of the pseudo-identity functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, !}$ is the invention of his.

The author is supported by NSF grant DMS-1063470.

1. THE INVENTORY OF CATEGORIES AND FUNCTORS

1.1. Eisenstein series and constant term functors.

1.1.1. Let P be a parabolic in G with Levi quotient M . For

$$\mu \in \pi_1(M) \simeq \pi_0(\mathrm{Bun}_M) \simeq \pi_0(\mathrm{Bun}_P),$$

let Bun_M^μ (resp., Bun_P^μ) denote the corresponding connected component of Bun_M (resp., Bun_P).

1.1.2. Consider the diagram

$$(1.1) \quad \begin{array}{ccc} & \text{Bun}_P^\mu & \\ \text{p} \swarrow & & \searrow \text{q} \\ \text{Bun}_G & & \text{Bun}_M^\mu. \end{array}$$

We consider the functor

$$\text{CT}_*^\mu : \text{D-mod}(\text{Bun}_G) \rightarrow \text{D-mod}(\text{Bun}_M^\mu), \quad \text{CT}_*^\mu = \text{q}_* \circ \text{p}^!$$

1.1.3. According to [DrGa3, Corollary 1.1.3], the functor CT_*^μ admits a left adjoint, denoted by $\text{Eis}_!^\mu$. Explicitly,

$$\text{Eis}_!^\mu = \text{p}_! \circ \text{q}^*.$$

The above expression has to be understood as follows: the functor

$$\text{q}^* : \text{D-mod}(\text{Bun}_M^\mu) \rightarrow \text{D-mod}(\text{Bun}_P^\mu)$$

is defined (because the morphism q is smooth), and the partially defined functor $\text{p}_!$, left adjoint to $\text{p}^!$, is defined on the essential image of q^* by [DrGa3, Proposition 1.1.2].

1.1.4. We define the functor $\text{CT}_* : \text{D-mod}(\text{Bun}_G) \rightarrow \text{D-mod}(\text{Bun}_M)$ as

$$\text{CT}_* \simeq \bigoplus_{\mu} \text{CT}_*^\mu.$$

We define the functor $\text{Eis}_! : \text{D-mod}(\text{Bun}_M) \rightarrow \text{D-mod}(\text{Bun}_G)$ as

$$\text{Eis}_! \simeq \bigoplus_{\mu} \text{Eis}_!^\mu.$$

Lemma 1.1.5. *The functor $\text{Eis}_!$ is the left adjoint of CT_* .*

Proof. Follows from the fact that

$$\bigoplus_{\mu} \text{CT}_*^\mu \simeq \prod_{\mu} \text{CT}_*^\mu.$$

□

1.1.6. We now consider the functor $\text{Eis}_*^\mu : \text{D-mod}(\text{Bun}_M^\mu) \rightarrow \text{D-mod}(\text{Bun}_G)$, defined as

$$\text{Eis}_*^\mu = \text{p}_* \circ \text{q}^!$$

We let $\text{Eis}_*^{\mu,-}$ and $\text{CT}_*^{\mu,-}$ be similarly defined functors when instead of P we use the opposite parabolic P^- (we identify the Levi quotients of P and P^- via the isomorphism $M \simeq P \cap P^-$).

The following is the main result of [DrGa3]:

Theorem 1.1.7. *The functor Eis_*^μ canonically identifies with the right adjoint of $\text{CT}_*^{\mu,-}$.*

1.1.8. We will use the notation

$$\mathrm{CT}_!^\mu : \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M^\mu)$$

for the left adjoint of Eis_*^μ . If $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ is such that the partially defined left adjoint \mathfrak{p}^* of \mathfrak{p}_* is defined on \mathcal{F} , then we have

$$\mathrm{CT}_!^\mu(\mathcal{F}) \simeq \mathfrak{q}_! \circ \mathfrak{p}^*(\mathcal{F}).$$

(The functor $\mathfrak{q}_!$, left adjoint to $\mathfrak{q}^!$, is well-defined by [DrGa3, Sect. 3.1.5].)

Hence, Theorem 1.1.7 can be reformulated as saying that $\mathrm{CT}_!^\mu$ exists and is canonically isomorphic to $\mathrm{CT}_*^{\mu,-}$.

1.1.9. We define the functor $\mathrm{CT}_! : \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M)$ as

$$\mathrm{CT}_! \simeq \bigoplus_{\mu} \mathrm{CT}_!^\mu,$$

so $\mathrm{CT}_! \simeq \mathrm{CT}_*^-$.

We define the functor $\mathrm{Eis}_* : \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ as

$$\mathrm{Eis}_* \simeq \bigoplus_{\mu} \mathrm{Eis}_*^\mu.$$

We note, however, that it is *no longer true* that Eis_* is the right adjoint of $\mathrm{CT}_!$. (Rather, the right adjoint of $\mathrm{CT}_!$ is the functor $\prod_{\mu} \mathrm{Eis}_*^\mu$.)

In fact, one can show that the functor Eis_* *does not admit* a left adjoint, see [DrGa3, Sect. 1.2.1].

1.2. The dual category.

1.2.1. Let $\mathrm{op}\text{-qc}(G)$ denote the poset of open substacks $U \xrightarrow{j} \mathrm{Bun}_G$ such that the intersection of U with every connected component of Bun_G is quasi-compact.

We have

$$(1.2) \quad \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \simeq \lim_{\leftarrow U \in \mathrm{op}\text{-qc}(G)} \mathrm{D}\text{-mod}(U),$$

where for $U_1 \xrightarrow{j_{1,2}} U_2$, the corresponding functor $\mathrm{D}\text{-mod}(U_2) \rightarrow \mathrm{D}\text{-mod}(U_1)$ is $j_{1,2}^*$ (see, e.g., [DrGa2, Lemma 2.3.2] for the proof).

Under the equivalence (1.2), for

$$(U \xrightarrow{j} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G),$$

the tautological evaluation functor $\mathrm{D}\text{-mod}(\mathrm{Bun}_G) \rightarrow \mathrm{D}\text{-mod}(U)$ is j^* .

1.2.2. The following DG category was introduced in [DrGa2, Sect. 4.3.3]:

$$(1.3) \quad \mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}} := \operatorname{colim}_{U \in \text{op-qc}(G)} \mathbf{D}\text{-mod}(U),$$

where for $U_1 \xrightarrow{j_{1,2}} U_2$, the corresponding functor $\mathbf{D}\text{-mod}(U_1) \rightarrow \mathbf{D}\text{-mod}(U_2)$ is $(j_{1,2})_*$, and where the colimit is taken in the category of cocomplete DG categories and continuous functors.

For $(U \xrightarrow{j} \mathbf{Bun}_G) \in \text{op-qc}(G)$ we let $j_{\text{co},*}$ denote the tautological functor

$$(1.4) \quad j_{\text{co},*} : \mathbf{D}\text{-mod}(U) \rightarrow \mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}.$$

1.2.3. Verdier duality functors

$$\mathbf{D}_U : \mathbf{D}\text{-mod}(U)^\vee \simeq \mathbf{D}\text{-mod}(U)$$

for $U \in \text{op-qc}(G)$ and the identifications

$$((j_{1,2})_*)^\vee \simeq (j_{1,2})^*, \quad U_1 \xrightarrow{j_{1,2}} U_2$$

give rise to an identification

$$(1.5) \quad \text{Func}_{\text{cont}}(\mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}, \mathbf{Vect}) \simeq \mathbf{D}\text{-mod}(\mathbf{Bun}_G).$$

Now, the main result of [DrGa2], namely, Theorem 4.1.8, implies:

Theorem 1.2.4. *The category $\mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}$ is compactly generated (and, in particular, dualizable).*

Proof. The truncatability of \mathbf{Bun}_G means that in the presentation of $\mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}$ as a colimit (1.3), we can replace the index poset $\text{op-qc}(G)$ by a *cofinal* poset that consists of quasi-compact open substacks that are *co-truncative*.

Then the resulting colimit

$$\operatorname{colim}_{U} \mathbf{D}\text{-mod}(U)$$

consists of compactly generated categories and functors *that preserve compactness*. In this case, the resulting colimit category is compactly generated, e.g., by [DrGa2, Corollary 1.9.4]. \square

From (1.5), and knowing that $\mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}$ is dualizable, we obtain a canonical identification

$$(1.6) \quad \mathbf{D}_{\mathbf{Bun}_G} : \mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}^\vee \simeq \mathbf{D}\text{-mod}(\mathbf{Bun}_G)_{\text{co}}.$$

Under this identification, for $(U \xrightarrow{j} \mathbf{Bun}_G) \in \text{op-qc}(G)$ we have the following canonical identification of functors

$$(j_{\text{co},*})^\vee \simeq j^*.$$

1.2.5. Similar constructions and notation apply when instead of all of \mathbf{Bun}_G we consider one of its connected components \mathbf{Bun}_G^λ , $\lambda \in \pi_1(G)$.

1.3. Dual, adjoint and conjugate functors.

1.3.1. Let \mathbf{C}_1 and \mathbf{C}_2 be two DG categories, and let

$$\mathbf{F} : \mathbf{C}_1 \rightleftarrows \mathbf{C}_2 : \mathbf{G}$$

be a pair of *continuous* mutually adjoint functors.

1.3.2. By passing to dual functors, the adjunction data

$$\mathrm{Id}_{\mathbf{C}_1} \rightarrow \mathbf{G} \circ \mathbf{F} \text{ and } \mathbf{F} \circ \mathbf{G} \rightarrow \mathrm{Id}_{\mathbf{C}_2}$$

gives rise to

$$\mathrm{Id}_{\mathbf{C}_1^\vee} \rightarrow \mathbf{F}^\vee \circ \mathbf{G}^\vee \text{ and } \mathbf{G}^\vee \circ \mathbf{F}^\vee \rightarrow \mathrm{Id}_{\mathbf{C}_2^\vee},$$

making

$$\mathbf{G}^\vee : \mathbf{C}_1^\vee \rightleftarrows \mathbf{C}_2^\vee : \mathbf{F}^\vee$$

into a pair of adjoint functors.

1.3.3. Assume now that \mathbf{C}_1 is compactly generated. In this case, the fact that the right adjoint \mathbf{G} of \mathbf{F} is continuous is equivalent to the fact that \mathbf{F} preserves compactness. I.e., it defines a functor between non-cocomplete DG categories

$$\mathbf{C}_1^c \rightarrow \mathbf{C}_2^c,$$

and hence, by passing to the opposite categories, a functor

$$(1.7) \quad (\mathbf{C}_1^c)^{\mathrm{op}} \rightarrow (\mathbf{C}_2^c)^{\mathrm{op}},$$

Following [Ga2, Sect. 1.5], we let

$$\mathbf{F}^{\mathrm{op}} : \mathbf{C}_1^\vee \rightarrow \mathbf{C}_2^\vee$$

denote the functor obtained as the composition of:

- (i) The identification $\mathbf{C}_1^\vee \simeq \mathrm{Ind}((\mathbf{C}_1^c)^{\mathrm{op}})$;
- (ii) The ind-extension $\mathrm{Ind}((\mathbf{C}_1^c)^{\mathrm{op}}) \rightarrow \mathrm{Ind}((\mathbf{C}_2^c)^{\mathrm{op}})$ of (1.7);
- (iii) The fully faithful embedding $(\mathbf{C}_2^c)^{\mathrm{op}} \hookrightarrow \mathbf{C}_2^\vee$.

We call \mathbf{F}^{op} the functor *conjugate* to \mathbf{F} .

1.3.4. The following is [Ga2, Lemma 1.5.3]:

Lemma 1.3.5. *We have a canonical isomorphism of functors $\mathbf{F}^{\mathrm{op}} \simeq \mathbf{G}^\vee$.*

1.4. Dual Eisenstein series and constant term functors.

1.4.1. We define the functor

$$\mathrm{Eis}_{\mathrm{co},*}^\mu : \mathrm{D}\text{-mod}(\mathrm{Bun}_M^\mu)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

as

$$\mathrm{Eis}_{\mathrm{co},*}^\mu \simeq (\mathrm{CT}_*^\mu)^\vee$$

under the identifications (1.6) and

$$\mathbf{D}_{\mathrm{Bun}_M^\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M^\mu)^\vee \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_M^\mu)_{\mathrm{co}}.$$

We define

$$\mathrm{Eis}_{\mathrm{co},*} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

as

$$\mathrm{Eis}_{\mathrm{co},*} := \bigoplus_{\mu} \mathrm{Eis}_{\mathrm{co},*}^\mu \simeq (\mathrm{CT}_*)^\vee.$$

Note that by Lemma 1.3.5, we have:

Corollary 1.4.2. *There are canonical isomorphisms*

$$\mathrm{Eis}_{\mathrm{co},*} \simeq (\mathrm{Eis}_!)^{\mathrm{op}} \text{ and } \mathrm{Eis}_{\mathrm{co},*}^\mu \simeq (\mathrm{Eis}_!^\mu)^{\mathrm{op}}.$$

1.4.3. We define the functor

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M^{\mu})_{\mathrm{co}}$$

as

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} \simeq (\mathrm{Eis}_*^{\mu})^{\vee}.$$

We define

$$\mathrm{CT}_{\mathrm{co},*} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}}$$

as

$$\mathrm{CT}_{\mathrm{co},*} := \bigoplus_{\mu} \mathrm{CT}_{\mathrm{co},*}^{\mu} \simeq (\mathrm{Eis}_*)^{\vee}.$$

From Lemma 1.3.5, we obtain:

Corollary 1.4.4. *There is a canonical isomorphism*

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} \simeq (\mathrm{CT}_!^{\mu})^{\mathrm{op}}.$$

1.4.5. Define also

$$\mathrm{CT}_{\mathrm{co},?}^{\mu} := (\mathrm{Eis}_!^{\mu})^{\vee} \text{ and } \mathrm{CT}_{\mathrm{co},?} := (\mathrm{Eis}_!)^{\vee} \simeq \bigoplus_{\mu} \mathrm{CT}_{\mathrm{co},?}^{\mu},$$

$$\mathrm{Eis}_{\mathrm{co},?}^{\mu} := (\mathrm{CT}_!^{\mu})^{\vee} \text{ and } \mathrm{Eis}_{\mathrm{co},?} := (\mathrm{CT}_!)^{\vee} \simeq \bigoplus_{\mu} \mathrm{Eis}_{\mathrm{co},?}^{\mu}.$$

By Sect. 1.3.2, we obtain the following pairs of adjoint functors

$$\mathrm{Eis}_{\mathrm{co},*}^{\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M^{\mu})_{\mathrm{co}} \rightleftarrows \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} : \mathrm{CT}_{\mathrm{co},?}^{\mu},$$

$$\mathrm{Eis}_{\mathrm{co},*} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}} \rightleftarrows \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} : \mathrm{CT}_{\mathrm{co},?},$$

and

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightleftarrows \mathrm{D}\text{-mod}(\mathrm{Bun}_M^{\mu})_{\mathrm{co}} : \mathrm{Eis}_{\mathrm{co},?}^{\mu}.$$

1.4.6. Finally, from Theorem 1.1.7, we obtain:

Corollary 1.4.7. *There are canonical isomorphisms of functors*

$$\mathrm{Eis}_{\mathrm{co},?}^{\mu} \simeq \mathrm{Eis}_{\mathrm{co},*}^{\mu,-}, \quad \mathrm{Eis}_{\mathrm{co},?} \simeq \mathrm{Eis}_{\mathrm{co},*}^-$$

and

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} \simeq (\mathrm{CT}_*^{\mu,-})^{\mathrm{op}}.$$

To summarize, we also obtain an adjunction

$$\mathrm{CT}_{\mathrm{co},*}^{\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightleftarrows \mathrm{D}\text{-mod}(\mathrm{Bun}_M^{\mu})_{\mathrm{co}} : \mathrm{Eis}_{\mathrm{co},*}^{\mu,-}.$$

1.4.8. We can ask the following question: does the functor $\mathrm{CT}_{\mathrm{co},*}^{\mu}$ admit a *left* adjoint? The answer is “no”:

Proof. If $\mathrm{CT}_{\mathrm{co},*}^{\mu}$ had admitted a left adjoint, by Sect. 1.3.2, the functor Eis_*^{μ} would have admitted a *continuous* right adjoint. However, this is not the case, since the functor

$$\mathrm{Eis}_*^{\mu} : \mathrm{D}\text{-mod}(\mathrm{Bun}_M^{\mu}) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

does not preserve compactness. □

1.5. **Explicit description of the dual functors.**

1.5.1. For $(U \xrightarrow{j} \text{Bun}_G) \in \text{op-qc}(G)$ we consider the functor $j^* : \text{D-mod}(\text{Bun}_G) \rightarrow \text{D-mod}(U)$, and its right adjoint j_* .

Define

$$j_{\text{co}}^* : \text{D-mod}(\text{Bun}_G)_{\text{co}} \rightarrow \text{D-mod}(U)$$

as

$$j_{\text{co}}^* := (j_*)^\vee.$$

By Sect. 1.3.2, the functors

$$j_{\text{co}}^* : \text{D-mod}(\text{Bun}_G)_{\text{co}} \rightleftarrows \text{D-mod}(U) : j_{\text{co},*}$$

form an adjoint pair, where $j_{\text{co},*}$ is as in (1.4).

Lemma 1.5.2. *The functor $j_{\text{co},*}$ is fully faithful.*

Proof. We need to show that the co-unit of the adjunction

$$j_{\text{co}}^* \circ j_{\text{co},*} \rightarrow \text{Id}_{\text{D-mod}(U)}$$

is an isomorphism. But this follows from the fact that the corresponding map between the dual functors, i.e.,

$$j^* \circ j_* \rightarrow \text{Id}_{\text{D-mod}(U)},$$

is an isomorphism (the latter because $j_* : \text{D-mod}(U) \rightarrow \text{D-mod}(\text{Bun}_G)$ is fully faithful). \square

1.5.3. By the definition of $\text{D-mod}(\text{Bun}_G)_{\text{co}}$, the functor j_{co}^* amounts to a compatible family of functors

$$j_{\text{co}}^* \circ (j_1)_{\text{co},*} : \text{D-mod}(U_1) \rightarrow \text{D-mod}(U)$$

for $(U_1 \xrightarrow{j_1} \text{Bun}_G) \in \text{op-qc}(G)$.

It is easy to see from the definitions that

$$j_{\text{co}}^* \circ (j_1)_{\text{co},*} \simeq (j_1)_* \circ (j')^*,$$

where

$$\begin{array}{ccc} U \cap U_1 & \xrightarrow{j'_1} & U \\ j' \downarrow & & \downarrow j \\ U_1 & \xrightarrow{j_1} & \text{Bun}_G. \end{array}$$

1.5.4. Again, by the definition of the category $\text{D-mod}(\text{Bun}_M)_{\text{co}}$, the functor $\text{Eis}_{\text{co},*}$ amounts to a compatible family of functors

$$\text{Eis}_{\text{co},*} \circ (j_M)_{\text{co},*} : \text{D-mod}(U_M) \rightarrow \text{D-mod}(\text{Bun}_G)_{\text{co}}$$

for $(U_M \xrightarrow{j_M} \text{Bun}_M) \in \text{op-qc}(M)$.

We now claim:

Proposition 1.5.5. *For a given $(U_M \xrightarrow{j_M} \text{Bun}_M) \in \text{op-qc}(M)$, let $(U_G \xrightarrow{j_G} \text{Bun}_G) \in \text{op-qc}(G)$ be such that*

$$\mathfrak{p}(\mathfrak{q}^{-1}(U_M)) \subset U_G.$$

Then there is a canonical isomorphism

$$\text{Eis}_{\text{co},*} \circ (j_M)_{\text{co},*} \simeq (j_G)_{\text{co},*} \circ (j_G)^* \circ \text{Eis}_* \circ (j_M)_* : \text{D-mod}(U_M) \rightarrow \text{D-mod}(\text{Bun}_G)_{\text{co}}.$$

Proof. First, we claim that there is a canonical isomorphism

$$(1.8) \quad \mathrm{Eis}_{\mathrm{co},*} \circ (j_M)_{\mathrm{co},*} \simeq (j_G)_{\mathrm{co},*} \circ (j_G)_{\mathrm{co}}^* \circ \mathrm{Eis}_{\mathrm{co},*} \circ (j_M)_{\mathrm{co},*}.$$

Indeed, (1.8) follows by passing to dual functors in the isomorphism

$$(j_M)^* \circ \mathrm{CT}_* \simeq (j_M)^* \circ \mathrm{CT}_* \circ (j_G)_* \circ (j_G)^*,$$

where the latter follows by base change from the definition.

Hence, it remains to establish a canonical isomorphism of functors

$$(j_G)_{\mathrm{co}}^* \circ \mathrm{Eis}_{\mathrm{co},*} \circ (j_M)_{\mathrm{co},*} \simeq (j_G)^* \circ \mathrm{Eis}_* \circ (j_M)_*, \quad \mathrm{D}\text{-mod}(U_M) \rightarrow \mathrm{D}\text{-mod}(U_G),$$

i.e., an isomorphism

$$((j_M)^* \circ \mathrm{CT}_* \circ (j_G)_*)^\vee \simeq (j_G)^* \circ \mathrm{Eis}_* \circ (j_M)_*.$$

However, the latter amounts to pull-push along the diagram

$$\begin{array}{ccccc} & & U_P & & \\ & & \swarrow \mathrm{p}|_{U_P} & & \searrow \mathrm{q}|_{U_P} \\ \mathrm{Bun}_G & \xleftarrow{j_P} & U_G & & U_M \xrightarrow{j_M} \mathrm{Bun}_M, \end{array}$$

where $U_P := \mathrm{q}^{-1}(U_M)$.

□

1.5.6. The functor $\mathrm{CT}_{\mathrm{co},*}$ amounts to a compatible family of functors

$$\mathrm{CT}_{\mathrm{co},*} \circ (j_G)_{\mathrm{co},*} : \mathrm{D}\text{-mod}(U_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}}$$

for $(U_G \xrightarrow{j_G} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G)$.

In a similar way to Proposition 1.5.5, we have:

Proposition 1.5.7. *For a given $(U_G \xrightarrow{j_G} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G)$, let $(U_M \xrightarrow{j_M} \mathrm{Bun}_M) \in \mathrm{op}\text{-qc}(M)$ be such that*

$$\mathrm{q}(\mathrm{p}^{-1}(U_G)) \subset U_M.$$

Then there is a canonical isomorphism

$$\mathrm{CT}_{\mathrm{co},*} \circ (j_G)_{\mathrm{co},*} \simeq (j_M)_{\mathrm{co},*} \circ (j_M)^* \circ \mathrm{CT}_* \circ (j_G)_* : \mathrm{D}\text{-mod}(U_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}}.$$

2. INTERACTION WITH THE NAIVE PSEUDO-IDENTITY AND CUSPIDALITY

2.1. The naive pseudo-identity functor.

2.1.1. The following functor

$$\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

was introduced in [DrGa2, Sect. 4.4.2]:

For $(U_G \xrightarrow{j_G} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G)$, the composition

$$\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}} \circ j_{\mathrm{co},*} : \mathrm{D}\text{-mod}(U_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

is by definition the functor j_* .

Remark 2.1.2. The functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ is very far from being an equivalence, unless G is a torus. For example, in [Ga2, Theorem 7.7.2], a particular object of $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$ was constructed, which belongs to $\ker(\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}})$, as soon as the semi-simple part of G is non-trivial.

2.1.3. Recall the equivalence:

$$\begin{aligned} \text{Funct}_{\text{cont}}(\text{D-mod}(\text{Bun}_G)_{\text{co}}, \text{D-mod}(\text{Bun}_G)) &\simeq (\text{D-mod}(\text{Bun}_G)_{\text{co}})^\vee \otimes \text{D-mod}(\text{Bun}_G) \simeq \\ &\simeq \text{D-mod}(\text{Bun}_G) \otimes \text{D-mod}(\text{Bun}_G) \simeq \text{D-mod}(\text{Bun}_G \times \text{Bun}_G). \end{aligned}$$

According to [DrGa2, Sect. 4.4.3], the functor $\text{Ps-Id}_{\text{Bun}_G, \text{naive}}$ corresponds to the object

$$(\Delta_{\text{Bun}_G})_*(\omega_{\text{Bun}_G}) \in \text{D-mod}(\text{Bun}_G \times \text{Bun}_G),$$

where Δ_{Bun_G} denotes the diagonal morphism on Bun_G , and $\omega_{\mathcal{Y}}$ is the dualizing object on a stack \mathcal{Y} (we take $\mathcal{Y} = \text{Bun}_G$).

From here we obtain:

Lemma 2.1.4. *There exists a canonical isomorphism $\text{Ps-Id}_{\text{Bun}_G, \text{naive}}^\vee \simeq \text{Ps-Id}_{\text{Bun}_G, \text{naive}}$.*

Proof. This expresses the fact that $(\Delta_{\text{Bun}_G})_*(\omega_{\text{Bun}_G})$ is equivariant with respect to the flip automorphism of $\text{D-mod}(\text{Bun}_G \times \text{Bun}_G)$. \square

Corollary 2.1.5. *For $(U_G \xrightarrow{j_G} \text{Bun}_G) \in \text{op-qc}(G)$, we have a canonical isomorphism:*

$$j^* \circ \text{Ps-Id}_{\text{Bun}_G, \text{naive}} \simeq j_{\text{co}}^*.$$

Proof. Obtained by passing to the dual functors is

$$\text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ j_{\text{co},*} \simeq j_*.$$

\square

2.1.6. We now claim:

Proposition 2.1.7. *There are canonical isomorphisms*

$$\text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ \text{Eis}_{\text{co},*} \simeq \text{Eis}_* \circ \text{Ps-Id}_{\text{Bun}_M, \text{naive}}$$

and

$$\text{Ps-Id}_{\text{Bun}_M, \text{naive}} \circ \text{CT}_{\text{co},*} \simeq \text{CT}_* \circ \text{Ps-Id}_{\text{Bun}_G, \text{naive}}.$$

Proof. We will prove the first isomorphism, while the second one is similar.

By definition, we need to construct a compatible family of isomorphisms of functors

$$\text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ \text{Eis}_{\text{co},*} \circ (j_M)_{\text{co},*} \simeq \text{Eis}_* \circ \text{Ps-Id}_{\text{Bun}_M, \text{naive}} \circ (j_M)_{\text{co},*}$$

for $(U_M \xrightarrow{j_M} \text{Bun}_M) \in \text{op-qc}(M)$.

For a given U_M , let U_G be as in Proposition 1.5.5. We rewrite

$$\begin{aligned} \text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ \text{Eis}_{\text{co},*} \circ (j_M)_{\text{co},*} &\simeq \text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ (j_G)_{\text{co},*} \circ (j_G)^* \circ \text{Eis}_* \circ (j_M)_* \simeq \\ &\simeq (j_G)_* \circ (j_G)^* \circ \text{Eis}_* \circ (j_M)_*. \end{aligned}$$

However, it is easy to see that for the above choice of U_G , the natural map

$$\text{Eis}_* \circ (j_M)_* \rightarrow (j_G)_* \circ (j_G)^* \circ \text{Eis}_* \circ (j_M)_*$$

is an isomorphism.

Now, by definition,

$$\text{Eis}_* \circ \text{Ps-Id}_{\text{Bun}_M, \text{naive}} \circ (j_M)_{\text{co},*} \simeq \text{Eis}_* \circ (j_M)_*,$$

and the assertion follows. (It is clear that these isomorphisms are independent of the choice of U_G , and hence are compatible under $(U_1)_M \hookrightarrow (U_2)_M$.) \square

2.2. Cuspidality.

2.2.1. Recall that in [DrGa3, Sect. 1.4] the full subcategory

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

was defined as the intersection of the kernels of the functors CT_* for *all proper parabolic subgroups* $P \subset G$.

Equivalently, let

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{Eis}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

be the full subcategory, generated by the essential images of the functors $\mathrm{Eis}_!$ for all proper parabolics. From the $(\mathrm{Eis}_!, \mathrm{CT}_*)$ -adjunction, we obtain

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} = (\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{Eis}})^\perp.$$

2.2.2. We let

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{Eis}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

be the full subcategory generated by the essential images of the functors

$$\mathrm{Eis}_{\mathrm{co}, *}: \mathrm{D}\text{-mod}(\mathrm{Bun}_M)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}.$$

We define

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} := (\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{Eis}})^\perp.$$

Equivalently, $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$ is the intersection of the kernels of the functors $\mathrm{CT}_{\mathrm{co}, ?}$ for all proper parabolics.

2.2.3. From Corollary 1.4.2 we obtain:

Corollary 2.2.4.

(1) *An object of $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$ is cuspidal if and only if its pairing with every object of $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{Eis}}$ is zero under the canonical map*

$$\langle -, - \rangle_{\mathrm{Bun}_G}: \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \times \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{Vect}$$

corresponding to $\mathbf{D}_{\mathrm{Bun}_G}$.

(2) *The identification $\mathbf{D}_{\mathrm{Bun}_G}: \mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$ induces identifications*

$$(\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{Eis}})^\vee \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{Eis}} \text{ and } (\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}})^\vee \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}.$$

Remark 2.2.5. We will see shortly that $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$ belongs to the intersection of the kernels of the functors $\mathrm{CT}_{\mathrm{co}, *}$ for all proper parabolics. But this inclusion is strict. For example for $G = SL_2$, the object from [Ga2, Theorem 7.7.2] belongs to $\mathrm{CT}_{\mathrm{co}, *}$ (there is only one parabolic to consider), but it does not belong to $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$.

2.2.6. Our goal for the rest of this section is to prove:

Theorem 2.2.7. *The restriction of the functor $\mathrm{Ps}\text{-Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ to*

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

takes values in $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \subset \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$, and defines an equivalence

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}.$$

2.3. Support of cuspidal objects.

2.3.1. The following crucial property of $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$ was established in [DrGa3, Proposition 1.4.6]:

Proposition 2.3.2. *There exists an element $(\mathcal{U}_G \xrightarrow{J_G} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G)$, such that for any $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$, the maps*

$$(J_G)! \circ (J_G)^*(\mathcal{F}) \rightarrow \mathcal{F} \rightarrow (J_G)_* \circ (J_G)^*(\mathcal{F})$$

are isomorphisms.

2.3.3. We now claim that a parallel phenomenon takes place for $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co},\mathrm{cusp}}$:

Proposition 2.3.4. *For any $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co},\mathrm{cusp}}$, the map*

$$\mathcal{F} \rightarrow (J_G)_{\mathrm{co},*} \circ (J_G)_{\mathrm{co}}^*(\mathcal{F})$$

is an isomorphism.

Proof. We need to show that the map from the tautological embedding

$$(2.1) \quad \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co},\mathrm{cusp}} \xrightarrow{\mathbf{e}_{\mathrm{co}}} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

to the composition

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co},\mathrm{cusp}} \xrightarrow{\mathbf{e}_{\mathrm{co}}} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \xrightarrow{(J_G)_{\mathrm{co}}^*} \mathrm{D}\text{-mod}(\mathcal{U}_G) \xrightarrow{(J_G)_{\mathrm{co},*}} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

is an isomorphism.

Note that in terms of the identification of Corollary 2.2.4(b), the dual of the embedding \mathbf{e}_{co} of (2.1) is the functor

$$(2.2) \quad \mathbf{f} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}},$$

left adjoint to the tautological embedding $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \xrightarrow{\mathbf{e}} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$.

Hence, by duality, we need to show that the functor (2.2) maps isomorphically to the composition

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G) \xrightarrow{(J_G)^*} \mathrm{D}\text{-mod}(\mathcal{U}_G) \xrightarrow{(J_G)_*} \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \xrightarrow{\mathbf{f}} \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}.$$

The latter is equivalent to the fact that any $\mathcal{F}' \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ for which $J_G^*(\mathcal{F}') = 0$, is left-orthogonal to $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$. However, this follows from the isomorphism

$$\mathcal{F} \rightarrow (J_G)_* \circ (J_G)^*(\mathcal{F}), \quad \mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$$

of Proposition 2.3.2. □

2.4. Description of the cuspidal category.

2.4.1. We claim:

Proposition 2.4.2. *Let $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$ be such that there exists $(U \xrightarrow{j} \mathrm{Bun}_G) \in \mathrm{op}\text{-qc}(G)$ such that the map*

$$\mathcal{F} \rightarrow j_{\mathrm{co},*} \circ j_{\mathrm{co}}^*(\mathcal{F})$$

is an isomorphism. Then $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co},\mathrm{cusp}}$ if and only if $\mathrm{CT}_{\mathrm{co},}(\mathcal{F}) = 0$ for all proper parabolics.*

Proof. Recall that

$$\langle -, - \rangle_{\text{Bun}_G} : \text{D-mod}(\text{Bun}_G)_{\text{co}} \times \text{D-mod}(\text{Bun}_G) \rightarrow \text{Vect}$$

denotes the pairing corresponding to the identification

$$\mathbf{D}_{\text{Bun}_G} : \text{D-mod}(\text{Bun}_G)^\vee \simeq \text{D-mod}(\text{Bun}_G)_{\text{co}}.$$

On the one hand, by Corollary 1.4.2, for $\mathcal{F}_G \in \text{D-mod}(\text{Bun}_G)_{\text{co}}$, the condition that \mathcal{F}_G be right-orthogonal to the essential image of $\text{Eis}_{\text{co},*}$ for a given parabolic P is equivalent to

$$\langle \text{Eis}_!(\mathcal{F}_M), \mathcal{F}_G \rangle_{\text{Bun}_G} = 0, \quad \mathcal{F}_M \in \text{D-mod}(\text{Bun}_M).$$

If $\mathcal{F}_G = j_{\text{co},*}(\mathcal{F}_U)$, then the above is equivalent to

$$\langle j^* \circ \text{Eis}_!(\mathcal{F}_M), \mathcal{F}_U \rangle_U = 0,$$

where

$$\langle -, - \rangle_U : \text{D-mod}(U)_{\text{co}} \times \text{D-mod}(U) \rightarrow \text{Vect}$$

is the pairing corresponding to $\mathbf{D}_U : \text{D-mod}(U)^\vee \simeq \text{D-mod}(U)$.

On the other hand, the condition that $\text{CT}_{\text{co},*}(\mathcal{F}_G) = 0$ for the same parabolic is equivalent to

$$\langle \text{Eis}_*(\mathcal{F}_M), \mathcal{F}_G \rangle_{\text{Bun}_G},$$

i.e.,

$$\langle j^* \circ \text{Eis}_*(\mathcal{F}_M), \mathcal{F}_U \rangle_U = 0.$$

Hence, the assertion of Proposition 2.4.2 follows from the next one, proved in Sect. 2.5:

Proposition 2.4.3.

(a) For $\mathcal{F}_M \in \text{D-mod}(\text{Bun}_M)$, the object $\text{Eis}_*(\mathcal{F}_M)$ admits an increasing filtration (indexed by a poset) with subquotients of the form $\text{Eis}_!(\mathcal{F}_M^\alpha)$, $\mathcal{F}_M^\alpha \in \text{D-mod}(\text{Bun}_M)$.

(b) Assume that \mathcal{F}_M is supported on finitely many connected components of Bun_M , and let $(U \xrightarrow{j} \text{Bun}_G) \in \text{op-qc}(G)$. Then:

(i) The objects $j^* \circ \text{Eis}_!(\mathcal{F}_M^\alpha)$ from point (a) are zero for all but finitely many α 's.

(ii) The object $j^* \circ \text{Eis}_!(\mathcal{F}_M)$ is a finite successive extension of objects of the form $j^* \circ \text{Eis}_*(\mathcal{F}_M^\alpha)$, $\mathcal{F}_M^\alpha \in \text{D-mod}(\text{Bun}_M)$.

□

2.4.4. We now observe:

Proposition 2.4.5. Let $\mathcal{F} \in \text{D-mod}(\text{Bun}_G)_{\text{co}}$ be such that there exists $(U_G \xrightarrow{j_G} \text{Bun}_G) \in \text{op-qc}(G)$ such that the map

$$\mathcal{F} \rightarrow (j_G)_{\text{co},*} \circ (j_G)_{\text{co}}^*(\mathcal{F})$$

is an isomorphism. Then $\text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}) \in \text{D-mod}(\text{Bun}_G)_{\text{cusp}}$ if and only if $\text{CT}_{\text{co},*}(\mathcal{F}) = 0$ for all proper parabolics.

Proof. We claim that for \mathcal{F} satisfying the condition of the proposition, for a given parabolic P ,

$$\text{CT}_* \circ \text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}) = 0 \Leftrightarrow \text{CT}_{\text{co},*}(\mathcal{F}) = 0.$$

Indeed, the implication \Leftarrow holds for any \mathcal{F} by Proposition 2.1.7.

Conversely, let $(U_M \xrightarrow{j_M} \text{Bun}_M) \in \text{op-qc}(M)$ be as in Proposition 1.5.7. For

$$\mathcal{F} \simeq (j_G)_{\text{co},*}(\mathcal{F}_{U_G}),$$

by Proposition 1.5.7, we have

$$\begin{aligned} \mathrm{CT}_{\mathrm{co},*}(\mathcal{F}) &\simeq (j_M)_{\mathrm{co},*} \circ (j_M)^* \circ \mathrm{CT}_* \circ (j_G)_*(\mathcal{F}_{U_G}) \simeq \\ &\simeq (j_M)_{\mathrm{co},*} \circ (j_M)^* \circ \mathrm{CT}_* \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}} \circ (j_G)_{\mathrm{co},*}(\mathcal{F}_{U_G}) \simeq \\ &\simeq (j_M)_{\mathrm{co},*} \circ (j_M)^* \circ \mathrm{CT}_* \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}}(\mathcal{F}). \end{aligned}$$

□

2.4.6. Combining Propositions 2.3.4, 2.4.2 and 2.4.5 we obtain:

Corollary 2.4.7. *For $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$ the following conditions are equivalent:*

- (i) $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$;
- (ii) *There exists $(U \xrightarrow{j} \mathrm{Bun}_G) \in \mathrm{op}\text{-}\mathrm{qc}(G)$ such that the map $\mathcal{F} \rightarrow j_{\mathrm{co},*} \circ j_{\mathrm{co}}^*(\mathcal{F})$ is an isomorphism and $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}}(\mathcal{F}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$.*
- (ii') *There exists $(U \xrightarrow{j} \mathrm{Bun}_G) \in \mathrm{op}\text{-}\mathrm{qc}(G)$ such that the map $\mathcal{F} \rightarrow j_{\mathrm{co},*} \circ j_{\mathrm{co}}^*(\mathcal{F})$ is an isomorphism and $\mathrm{CT}_{\mathrm{co},*}(\mathcal{F}) = 0$ for all proper parabolics.*
- (iii) *For $(\mathcal{U}_G \xrightarrow{j^G} \mathrm{Bun}_G) \in \mathrm{op}\text{-}\mathrm{qc}(G)$ as in Proposition 2.3.2, the map $\mathcal{F} \rightarrow (j_G)_{\mathrm{co},*} \circ (j_G)_{\mathrm{co}}^*(\mathcal{F})$ is an isomorphism and $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}}(\mathcal{F}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$.*
- (iii') *For $(\mathcal{U}_G \xrightarrow{j^G} \mathrm{Bun}_G) \in \mathrm{op}\text{-}\mathrm{qc}(G)$ as in Proposition 2.3.2, the map $\mathcal{F} \rightarrow (j_G)_{\mathrm{co},*} \circ (j_G)_{\mathrm{co}}^*(\mathcal{F})$ is an isomorphism and $\mathrm{CT}_{\mathrm{co},*}(\mathcal{F}) = 0$ for all proper parabolics.*

2.4.8. *Proof of Theorem 2.2.7.* From Corollary 2.4.7 we obtain that the functor $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}}$ sends

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}.$$

We construct the inverse functor as follows. Let $(\mathcal{U}_G \xrightarrow{j^G} \mathrm{Bun}_G) \in \mathrm{op}\text{-}\mathrm{qc}(G)$ be as in Proposition 2.3.2. The sought-for functor

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$$

is

$$\mathcal{F} \mapsto (j_G)_{\mathrm{co},*} \circ (j_G)^*(\mathcal{F}).$$

We claim that the image of this functor lands in $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$. Indeed, by Proposition 2.4.5, it suffices to check that

$$\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}} \circ (j_G)_{\mathrm{co},*} \circ (j_G)^*(\mathcal{F}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}.$$

However,

$$\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}} \circ (j_G)_{\mathrm{co},*} \circ (j_G)^*(\mathcal{F}) \simeq (j_G)_* \circ (j_G)^*(\mathcal{F}),$$

and the latter is isomorphic to \mathcal{F} by Proposition 2.4.5.

Let us now check that the two functors are inverses of each other. However, we have just shown that the composition

$$\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{cusp}}$$

is isomorphic to the identity functor.

For the composition in the other direction, for $\mathcal{F} \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}, \mathrm{cusp}}$ we consider

$$(j_G)_{\mathrm{co},*} \circ (j_G)^* \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, \mathrm{naive}}(\mathcal{F}),$$

which by Corollary 2.1.5 is isomorphic to

$$(j_G)_{\mathrm{co},*} \circ (j_G)_{\mathrm{co}}^*(\mathcal{F}),$$

and the latter is isomorphic to \mathcal{F} by Proposition 2.3.4. □

2.5. Proof of Proposition 2.4.3.

2.5.1. The proof of the proposition uses the relative compactification $\text{Bun}_P \xrightarrow{r} \widetilde{\text{Bun}}_P$ of the map \mathfrak{p} , introduced in [BG, Sect. 1.3.6]:

$$(2.3) \quad \begin{array}{ccc} \text{Bun}_P & \xrightarrow{r} & \widetilde{\text{Bun}}_P \\ & \searrow \tilde{\mathfrak{p}} & \swarrow \tilde{\mathfrak{q}} \\ & \text{Bun}_G & \text{Bun}_M \end{array}$$

Note that for $\mathcal{F}_M \in \text{D-mod}(\text{Bun}_M)$, we have

$$\text{Eis}_*(\mathcal{F}_M) \simeq \tilde{\mathfrak{p}}_* \left(\tilde{\mathfrak{q}}^!(\mathcal{F}_M) \overset{!}{\otimes} r_*(\omega_{\text{Bun}_P}) \right) \simeq \tilde{\mathfrak{p}}! \left(\tilde{\mathfrak{q}}^!(\mathcal{F}_M) \overset{!}{\otimes} r_*(\omega_{\text{Bun}_P}) \right),$$

the latter isomorphism due to the fact that $\tilde{\mathfrak{p}}$ is proper. Here the notation $\overset{!}{\otimes}$ (and, in the sequel, $\overset{*}{\otimes}$) follows [DrGa3, Sect. 1.1.5].

Recall now that according to [BG, Theorem 5.1.5], the object

$$r_*(\omega_{\text{Bun}_P}) \in \text{D-mod}(\widetilde{\text{Bun}}_P)$$

is *universally locally acyclic* (a.k.a. ULA)⁴ with respect to the map $\tilde{\mathfrak{q}}$. This implies that

$$\tilde{\mathfrak{q}}^!(\mathcal{F}_M) \overset{!}{\otimes} r_*(\omega_{\text{Bun}_P}) \simeq \tilde{\mathfrak{q}}^*(\mathcal{F}_M) \overset{*}{\otimes} r_*(\omega_{\text{Bun}_P})[-2 \dim(\text{Bun}_M)].$$

Thus, we obtain that, up to a cohomological shift, $\text{Eis}_*(\mathcal{F}_M)$ is isomorphic to

$$(2.4) \quad \tilde{\mathfrak{p}}! \left(\tilde{\mathfrak{q}}^*(\mathcal{F}_M) \overset{*}{\otimes} r_*(\omega_{\text{Bun}_P}) \right).$$

2.5.2. Let $\Lambda_{G,P}^{\text{pos}}$ be the monoid of linear combinations

$$\theta = \sum_i n_i \cdot \alpha_i,$$

where $n_i \in \mathbb{Z}^{\geq 0}$ and α_i is a simple coroot of G , which is not in M .

For each θ , we let $\text{Mod}_{\text{Bun}_M}^{\theta,+}$ be a version of the Hecke stack, introduced in [BFGM, Sect. 3.1]:

$$\begin{array}{ccc} & \text{Mod}_{\text{Bun}_M}^{\theta,+} & \\ \swarrow \tilde{h} & & \searrow \tilde{h} \\ \text{Bun}_M & & \text{Bun}_M \end{array}$$

Set

$$\text{Mod}_{\text{Bun}_P}^{\theta,+} := \text{Bun}_P \times_{\text{Bun}_M} \text{Mod}_{\text{Bun}_M}^{\theta,+},$$

⁴See [DrGa3, Sect. 1.1.5] for what the ULA condition means.

where the fiber product is formed using the map $\overleftarrow{h} : \text{Mod}_{\text{Bun}_M}^{\theta,+} \rightarrow \text{Bun}_M$.

According to [BG, Proposition 6.2.5], there is a canonically defined locally closed embedding

$$r^\theta : \text{Mod}_{\text{Bun}_P}^{\theta,+} \rightarrow \widetilde{\text{Bun}}_P,$$

making the following diagram commute

$$\begin{array}{ccccc}
 & & \text{Mod}_{\text{Bun}_P}^{\theta,+} & & \\
 & \swarrow \overleftarrow{h} & \downarrow \text{'q} & \searrow r^\theta & \\
 \text{Bun}_P & & & & \widetilde{\text{Bun}}_P \\
 \downarrow \text{q} & & \downarrow \text{'q} & & \downarrow \tilde{\text{q}} \\
 & & \text{Mod}_{\text{Bun}_M}^{\theta,+} & & \\
 & \swarrow \overleftarrow{h} & & \searrow \overrightarrow{h} & \\
 \text{Bun}_M & & & & \text{Bun}_M .
 \end{array}$$

(The right diamond is intentionally lopsided to emphasize that it is *not* Cartesian.)

Furthermore,

$$(2.5) \quad \widetilde{\text{Bun}}_P = \bigsqcup_{\theta \in \Lambda_{G,P}^{\text{pos}}} r^\theta(\text{Mod}_{\text{Bun}_P}^{\theta,+}).$$

For $\theta = 0$, the map \overleftarrow{h} is an isomorphism, and the resulting map

$$\text{Bun}_P \simeq \text{Mod}_{\text{Bun}_P}^{0,+} \xrightarrow{r^0} \widetilde{\text{Bun}}_P$$

is the map r in (2.3).

The following is easy to see from the construction:

Lemma 2.5.3. *For $(U \xrightarrow{j} \text{Bun}_G) \in \text{op-qc}(G)$ and $\mu \in \pi_1(M)$, the preimage of $U \times \text{Bun}_M^\mu$ under the map*

$$\text{Mod}_{\text{Bun}_P}^{\theta,+} \xrightarrow{r^\theta} \widetilde{\text{Bun}}_P \xrightarrow{\tilde{\text{p}} \times \tilde{\text{q}}} \text{Bun}_G \times \text{Bun}_M$$

is empty for all but finitely many elements θ .

2.5.4. The decomposition (2.5) endows the object

$$r_*(\omega_{\text{Bun}_P}) \in \text{D-mod}(\widetilde{\text{Bun}}_P)$$

with an increasing filtration, indexed by the poset $\Lambda_{G,P}^{\text{pos}}$, with the θ subquotient equal to

$$(r^\theta)_! \circ (r^\theta)^* \circ r_*(\omega_{\text{Bun}_P}).$$

Hence, by the projection formula, the object in (2.4) admits a filtration, indexed by $\Lambda_{G,P}^{\text{pos}}$, with the θ subquotient equal to

$$(2.6) \quad \tilde{\text{p}}_! \circ r_!^\theta \left((r^\theta)^* \circ \tilde{\text{q}}^*(\mathcal{F}_M) \otimes^* (r^\theta)^* \circ r_*(\omega_{\text{Bun}_P}) \right).$$

Moreover, if \mathcal{F}_M is supported on finitely many components of Bun_M , the restriction of the subquotient (2.6) to $U \in \text{op-qc}(G)$ is zero for all but finitely many θ by Lemma 2.5.3.

2.5.5. We have the following assertion, proved by the same argument as [BG, Theorem 6.2.10]:

Lemma 2.5.6. *The object $(r^\theta)^* \circ r_*(\omega_{\text{Bun}_P}) \in \text{D-mod}(\text{Mod}_{\text{Bun}_P}^{\theta,+})$ is lisse when $!$ -restricted to the fiber of the map*

$$'q : \text{Mod}_{\text{Bun}_P}^{\theta,+} \rightarrow \text{Mod}_{\text{Bun}_M}^{\theta,+}$$

over any k -point of $\text{Mod}_{\text{Bun}_M}^{\theta,+}$.

Corollary 2.5.7. $(r^\theta)^* \circ r_*(\omega_{\text{Bun}_P}) \simeq 'q^*(\mathcal{K}^\theta)$ for some $\mathcal{K}^\theta \in \text{D-mod}(\text{Mod}_{\text{Bun}_M}^{\theta,+})$.

Proof. Follows from Lemma 2.5.6 plus the combination of the following three facts: (1) the map $'q$ is smooth; (2) $(r^\theta)^* \circ r_*(\omega_{\text{Bun}_P})$ is holonomic with regular singularities; (3) the fibers of the map $'q$ are contractible (and hence any RS local system on such a fiber is canonically trivial). □

2.5.8. By Corollary 2.5.7, we can rewrite the subquotient (2.6) as

$$\tilde{p}_! \circ r_!^\theta \left((r^\theta)^* \circ \tilde{q}^*(\mathcal{F}_M) \otimes^* ('q)^*(\mathcal{K}^\theta) \right),$$

and further, using the fact that

$$\tilde{q} \circ r^\theta = \vec{h} \circ 'q \text{ and } \tilde{p} \circ r^\theta = p \circ 'h$$

as

$$p_! \circ 'h_! \circ 'q^* \left(\vec{h}^*(\mathcal{F}_M) \otimes^* \mathcal{K}^\theta \right) \simeq p_! \circ q^* \left(\overleftarrow{h}_!(\vec{h}^*(\mathcal{F}_M) \otimes^* \mathcal{K}^\theta) \right).$$

To summarize, we identify the subquotient (2.6) with

$$\text{Eis}_! \left(\overleftarrow{h}_!(\vec{h}^*(\mathcal{F}_M) \otimes^* \mathcal{K}^\theta) \right),$$

as required in Proposition 2.4.3(a). The finiteness assertion in Proposition 2.4.3(b)(i) follows from the finiteness at the end of Sect. 2.5.4.

2.5.9. The proof of Proposition 2.4.3(b)(ii) is similar, but with the following modification:

Let $k_{\text{Bun}_P} \in \text{D-mod}(\text{Bun}_P)$ be the “constant sheaf” D-module, i.e., the Verdier dual of ω_{Bun_P} .

Then the object

$$r_!(k_{\text{Bun}_P}) \in \text{D-mod}(\widetilde{\text{Bun}}_P)$$

admits a *decreasing* filtration, indexed by the poset $\Lambda_{G,P}^{\text{pos}}$, with the θ subquotient being

$$(r^\theta)_* \circ (r^\theta)^! \circ r_!(k_{\text{Bun}_P}).$$

However, this filtration is finite on the preimage of $U \times \text{Bun}_M^\mu$ for any $U \in \text{op-qc}(G)$ and $\mu \in \pi_1(M)$ under the map $\tilde{p} \times \tilde{q}$, again by Lemma 2.5.3.

3. INTERACTION WITH THE GENUINE PSEUDO-IDENTITY FUNCTOR

3.1. The pseudo-identity functor.

3.1.1. We now recall that in [DrGa2, Sect. 4.4.8] another functor, denoted

$$\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!} : \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}} \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$$

was introduced.

Namely, in terms of the equivalences

$$(3.1) \quad \begin{aligned} \mathrm{Funct}_{\mathrm{cont}}(\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}, \mathrm{D}\text{-mod}(\mathrm{Bun}_G)) &\simeq (\mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}})^\vee \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \simeq \\ &\simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G \times \mathrm{Bun}_G), \end{aligned}$$

the functor $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!}$ corresponds to the object

$$(\Delta_{\mathrm{Bun}_G})!(k_{\mathrm{Bun}_G}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G \times \mathrm{Bun}_G).$$

3.1.2. Note the following feature of the functor $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!}$, parallel to one for $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,\mathrm{naive}}$, given by Lemma 2.1.4.

Lemma 3.1.3. *Under the identification $\mathrm{D}\text{-mod}(\mathrm{Bun}_G)^\vee \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G)_{\mathrm{co}}$, we have*

$$(\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!})^\vee \simeq \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!}.$$

Proof. This is just the fact that the object $(\Delta_{\mathrm{Bun}_G})!(k_{\mathrm{Bun}_G}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G \times \mathrm{Bun}_G)$ is equivariant with respect to the flip. \square

3.1.4. The goal of this section and the next is to prove:

Theorem 3.1.5. *The functor $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!}$ is an equivalence.*

The proof will rely on a certain geometric result, namely, Proposition 3.2.6 which will be proved in [Sch].

3.2. Relation between the two functors.

3.2.1. Consider again the map

$$\Delta_{\mathrm{Bun}_G} : \mathrm{Bun}_G \rightarrow \mathrm{Bun}_G \times \mathrm{Bun}_G.$$

It naturally factors as

$$\mathrm{Bun}_G \xrightarrow{\mathrm{id}_{\mathrm{Bun}_G}^Z} \mathrm{Bun}_G \times B(Z_G) \xrightarrow{\Delta_{\mathrm{Bun}_G}^Z} \mathrm{Bun}_G \times \mathrm{Bun}_G,$$

where:

- Z_G denotes the center of G , and $B(Z_G)$ is its classifying stack;
- The map $\mathrm{id}_{\mathrm{Bun}_G}^Z$ is given by the identity map $\mathrm{Bun}_G \rightarrow \mathrm{Bun}_G$, and

$$\mathrm{Bun}_G \rightarrow \mathrm{pt} \xrightarrow{\mathrm{triv}} B(Z_G),$$

where $\mathrm{triv} : \mathrm{pt} \rightarrow B(Z_G)$ corresponds to the trivial Z_G -bundle;

- The composition $\mathrm{pr}_1 \circ \Delta_{\mathrm{Bun}_G}^Z$ is projection on the first factor $\mathrm{Bun}_G \times B(Z_G) \rightarrow \mathrm{Bun}_G$;
- The composition $\mathrm{pr}_2 \circ \Delta_{\mathrm{Bun}_G}^Z$ is given by the natural action of $B(Z_G)$ on Bun_G .

Remark 3.2.2. Note that if G is a torus, the map $\Delta_{\mathrm{Bun}_G}^Z$ is an isomorphism.

3.2.3. We write

$$(\Delta_{\text{Bun}_G})_*(\omega_{\text{Bun}_G}) \simeq (\Delta_{\text{Bun}_G}^Z)_* \circ (\text{id}_{\text{Bun}_G}^Z)_*(\omega_{\text{Bun}_G}).$$

In addition,

$$(\Delta_{\text{Bun}_G})!(k_{\text{Bun}_G}) \simeq (\Delta_{\text{Bun}_G}^Z)!(\text{id}_{\text{Bun}_G}^Z)!(k_{\text{Bun}_G}) \simeq (\Delta_{\text{Bun}_G}^Z)!(\text{id}_{\text{Bun}_G}^Z)!(\omega_{\text{Bun}_G})[-2 \dim(\text{Bun}_G)],$$

the latter isomorphism is due to the fact that Bun_G is smooth.

It is easy to see that

$$\text{triv}_!(k) \simeq \text{triv}_*(k)[- \dim(Z_G)].$$

Hence,

$$(\Delta_{\text{Bun}_G})!(k_{\text{Bun}_G}) \simeq (\Delta_{\text{Bun}_G}^Z)!(\text{id}_{\text{Bun}_G}^Z)_*(\omega_{\text{Bun}_G})[-2 \dim(\text{Bun}_G) - \dim(Z_G)].$$

Now, the morphism

$$\Delta_{\text{Bun}_G}^Z : \text{Bun}_G \times B(Z_G) \rightarrow \text{Bun}_G \times \text{Bun}_G$$

is schematic and separated. Hence, we obtain a natural transformation

$$(3.2) \quad (\Delta_{\text{Bun}_G}^Z)!(k_{\text{Bun}_G}) \rightarrow (\Delta_{\text{Bun}_G}^Z)_*(\omega_{\text{Bun}_G}).$$

Summarizing, we obtain a map

$$(3.3) \quad (\Delta_{\text{Bun}_G})!(k_{\text{Bun}_G}) \rightarrow (\Delta_{\text{Bun}_G})_*(\omega_{\text{Bun}_G})[-2 \dim(\text{Bun}_G) - \dim(Z_G)].$$

3.2.4. From (3.3) and Sect. 2.1.3, we obtain a natural transformation:

$$(3.4) \quad \text{Ps-Id}_{\text{Bun}_G,!} \rightarrow \text{Ps-Id}_{\text{Bun}_G,\text{naive}}[-2 \dim(\text{Bun}_G) - \dim(Z_G)]$$

as functors $\text{D-mod}(\text{Bun}_G)_{\text{co}} \rightarrow \text{D-mod}(\text{Bun}_G)$.

Let

$$\text{Ps-Id}_{\text{Bun}_G,\text{diff}} : \text{D-mod}(\text{Bun}_G)_{\text{co}} \rightarrow \text{D-mod}(\text{Bun}_G)$$

denote the cone of the natural transformation (3.4).

3.2.5. We claim:

Proposition 3.2.6. *The functor $\text{Ps-Id}_{\text{Bun}_G,\text{diff}}$ admits a decreasing filtration, indexed by a poset, with subquotients being functors of the form*

$$\begin{aligned} \text{D-mod}(\text{Bun}_G)_{\text{co}} &\xrightarrow{\text{CT}_{\text{co},*}^\mu} \text{D-mod}(\text{Bun}_M^\mu)_{\text{co}} \xrightarrow{\text{Ps-Id}_{\text{Bun}_M^\mu,\text{naive}}} \text{D-mod}(\text{Bun}_M^\mu) \xrightarrow{\mathbf{F}^{\mu,\mu'}} \\ &\rightarrow \text{D-mod}(\text{Bun}_M^{\mu'}) \xrightarrow{\text{Eis}_*^{\mu',-}} \text{D-mod}(\text{Bun}_G), \end{aligned}$$

for a proper parabolic P with Levi quotient M , where $\mu, \mu' \in \pi_1(M)$ and $\mathbf{F}^{\mu,\mu'}$ is some functor $\text{D-mod}(\text{Bun}_M^\mu) \rightarrow \text{D-mod}(\text{Bun}_M^{\mu'})$. Furthermore, for a pair

$$(U_1 \xrightarrow{j_1} \text{Bun}_G), (U_2 \xrightarrow{j_2} \text{Bun}_G) \in \text{op-qc}(G),$$

the induced filtration on

$$j_1^* \circ \text{Ps-Id}_{\text{Bun}_G,\text{diff}} \circ (j_2)_{\text{co},*}$$

is finite.

The proof of Proposition 3.2.6 is analogous to that of Proposition 2.4.3 and will be given in [Sch].

As its geometric ingredient, instead of the stack $\widetilde{\text{Bun}}_P$ appearing in the proof of Proposition 2.4.3, one uses a compactification of the morphism $\Delta_{\text{Bun}_G}^{Z_G}$ which can be constructed using Vinberg's canonical semi-group of [Vi] attached to G .

3.3. Pseudo-identity and cuspidality.

3.3.1. As a consequence of Proposition 3.2.6, we obtain:

Corollary 3.3.2. *The morphism (3.4) induces an isomorphism*

$$\text{Ps-Id}_{\text{Bun}_G,!} |_{\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}} \simeq \text{Ps-Id}_{\text{Bun}_G,\text{naive}} |_{\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}} [-2 \dim(\text{Bun}_G) - \dim(Z_G)].$$

Proof. By the definition of $\text{D-mod}(\text{Bun}_G)$, it is sufficient to show that for any $(U_1 \xrightarrow{j_1} \text{Bun}_G) \in \text{op-qc}(G)$, the map (3.4) induces an isomorphism

$$\begin{aligned} j_1^* \circ \text{Ps-Id}_{\text{Bun}_G,!} |_{\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}} &\rightarrow \\ &\rightarrow j_1^* \circ \text{Ps-Id}_{\text{Bun}_G,\text{naive}} |_{\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}} [-2 \dim(\text{Bun}_G) - \dim(Z_G)]. \end{aligned}$$

Let us take $U_2 := \mathcal{U}_G$ as in Proposition 2.3.2. By Proposition 2.3.4, it suffices to show that for $\mathcal{F} \in \text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}$

$$j_1^* \circ \text{Ps-Id}_{\text{Bun}_G,\text{diff}} \circ (j_2)_{\text{co},*} \circ (j_2)_{\text{co}}^*(\mathcal{F}) = 0.$$

However, this follows from Proposition 3.2.6:

Indeed, the object in question has a finite filtration, with subquotients isomorphic to

$$j_1^* \circ \text{Eis}_*^{\mu', -} \circ \mathcal{F}^{\mu, \mu'} \circ \text{Ps-Id}_{\text{Bun}_M^{\mu}, \text{naive}} \circ \text{CT}_{\text{co},*}^{\mu} \circ (j_2)_{\text{co},*} \circ (j_2)_{\text{co}}^*(\mathcal{F}),$$

which, by Proposition 2.3.4, is isomorphic to

$$j_1^* \circ \text{Eis}_*^{\mu', -} \circ \mathcal{F}^{\mu, \mu'} \circ \text{Ps-Id}_{\text{Bun}_M^{\mu}, \text{naive}}(\text{CT}_{\text{co},*}^{\mu}(\mathcal{F})),$$

while $\text{CT}_{\text{co},*}^{\mu}(\mathcal{F}) = 0$ by Corollary 2.4.7. □

Corollary 3.3.3. *The functor $\text{Ps-Id}_{\text{Bun}_G,!}$ induces an equivalence*

$$\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}} \rightarrow \text{D-mod}(\text{Bun}_G)_{\text{cusp}}.$$

Proof. Follows from Theorem 2.2.7 and Corollary 3.3.2. □

3.3.4. The next assertion is a crucial step in the proof of Theorem 3.1.5:

Proposition 3.3.5. *The functor $\text{Ps-Id}_{\text{Bun}_G,!}$ induces an isomorphism*

$$\text{Hom}_{\text{D-mod}(\text{Bun}_G)_{\text{co}}}(\mathcal{F}', \mathcal{F}) \rightarrow \text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})),$$

provided that $\mathcal{F}' \in \text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}$.

3.4. Proof of Proposition 3.3.5.

3.4.1. Let us first assume that \mathcal{F} has the form $j_{\text{co},*}(\mathcal{F}_U)$ for some $(U \xrightarrow{j} \text{Bun}_G) \in \text{op-qc}(G)$.

Consider the commutative diagram

$$(3.5) \quad \begin{array}{ccc} \text{Hom}(\mathcal{F}', \mathcal{F}) & \longrightarrow & \text{Hom}(\text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F})) \\ \downarrow & & \downarrow \\ \text{Hom}(\text{Ps-Id}_{\text{Bun}_G, !}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G, !}(\mathcal{F})) & \longrightarrow & \text{Hom}(\text{Ps-Id}_{\text{Bun}_G, !}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F})[d]), \end{array}$$

where $d = -2 \dim(\text{Bun}_G) - \dim(Z_G)$.

We need to show that the left vertical arrow is an isomorphism. We will do so by showing that all the other arrows are isomorphisms.

3.4.2. First, we claim that upper horizontal arrow in (3.5) is an isomorphism for any $\mathcal{F}' \in \text{D-mod}(\text{Bun}_G)_{\text{co}}$ and $\mathcal{F} = j_*(\mathcal{F}_U)$. Indeed, the map in question fits into a commutative diagram

$$\begin{array}{ccc} \text{Hom}_{\text{D-mod}(\text{Bun}_G)_{\text{co}}}(\mathcal{F}', j_{\text{co},*}(\mathcal{F}_U)) & \longrightarrow & \text{Hom}(\text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G, \text{naive}} \circ j_{\text{co},*}(\mathcal{F}_U)) \\ \sim \downarrow & & \downarrow \sim \\ \text{Hom}_{\text{D-mod}(U)}(j_{\text{co}}^*(\mathcal{F}'), \mathcal{F}_U) & & \text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}'), j_*(\mathcal{F}_U)) \\ \text{id} \downarrow & & \downarrow \sim \\ \text{Hom}_{\text{D-mod}(U)}(j_{\text{co}}^*(\mathcal{F}'), \mathcal{F}_U) & \xrightarrow{\sim} & \text{Hom}_{\text{D-mod}(U)}(j^* \circ \text{Ps-Id}_{\text{Bun}_G, \text{naive}}(\mathcal{F}'), \mathcal{F}_U). \end{array}$$

3.4.3. The right vertical arrow in (3.5) is an isomorphism by Corollary 3.3.2.

To show that the lower horizontal arrow is an isomorphism, using Corollary 3.3.3, it suffices to show that for any $\mathcal{F}'' \in \text{D-mod}(\text{Bun}_G)_{\text{cusp}}$, we have

$$\text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\mathcal{F}'', \text{Ps-Id}_{\text{Bun}_G, \text{diff}} \circ j_{\text{co},*}(\mathcal{F}_U)) = 0.$$

By Proposition 2.3.2,

$$\begin{aligned} \text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\mathcal{F}'', \text{Ps-Id}_{\text{Bun}_G, \text{diff}} \circ j_{\text{co},*}(\mathcal{F}_U)) &\simeq \\ &\simeq \text{Hom}_{\text{D-mod}(\mathcal{U}_G)}(j_G^*(\mathcal{F}''), j_G^* \circ \text{Ps-Id}_{\text{Bun}_G, \text{diff}} \circ j_{\text{co},*}(\mathcal{F}_U)). \end{aligned}$$

Applying Proposition 3.2.6, we obtain that it suffices to show that for $\mathcal{F}'' \in \text{D-mod}(\text{Bun}_G)_{\text{cusp}}$

$$\text{Hom}_{\text{D-mod}(\mathcal{U}_G)}\left(j_G^*(\mathcal{F}''), j_G^* \circ \text{Eis}_*^{\mu', -} \circ F^{\mu, \mu'} \circ \text{Ps-Id}_{\text{Bun}_M^\mu, \text{naive}} \circ \text{CT}_{\text{co},*}^\mu \circ j_{\text{co},*}(\mathcal{F}_U)\right) = 0,$$

which by Proposition 2.3.2 is equivalent to

$$\text{Hom}_{\text{D-mod}(\text{Bun}_G)}\left(\mathcal{F}'', \text{Eis}_*^{\mu', -} \circ F^{\mu, \mu'} \circ \text{Ps-Id}_{\text{Bun}_M^\mu, \text{naive}} \circ \text{CT}_{\text{co},*}^\mu \circ j_{\text{co},*}(\mathcal{F}_U)\right) = 0.$$

Now, $\text{D-mod}(\text{Bun}_G)_{\text{cusp}}$ is *left-orthogonal* to the essential image of $\text{Eis}_*^{\mu', -}$ by Theorem 1.1.7, implying the desired vanishing.

3.4.4. We will now reduce the assertion of Proposition 3.3.5 to the situation of Sect. 3.4.1.

Let us recall that according to [DrGa2, Theorem 4.1.8], any element $(U \xrightarrow{j} \text{Bun}_G) \in \text{op-qc}(G)$ is contained in one which is *co-truncative*. See [DrGa2, Sect. 3.8] for what it means for an open substack to be co-truncative. In particular, the open substack \mathcal{U}_G of Proposition 2.3.2 can be enlarged so that it is co-truncative.

Recall also that for a co-truncative open substack $U \xrightarrow{j} \text{Bun}_G$, the functor $j_{\text{co},*}$ has a (continuous) right adjoint, denoted $j^?$, see [DrGa2, Sect. 4.3].

Any $\mathcal{F} \in \text{D-mod}(\text{Bun}_G)_{\text{co}}$ fits into an exact triangle

$$\mathcal{F}_1 \rightarrow \mathcal{F} \rightarrow j_{\text{co},*} \circ j^?(\mathcal{F}),$$

where $j^?(\mathcal{F}_1) = 0$ by Lemma 1.5.2.

We take U to contain the substack \mathcal{U}_G as in Proposition 2.3.2, and assume that it is co-truncative. In view of Proposition 2.3.4 and Corollary 3.3.3, it remains to show that if $j^?(\mathcal{F}) = 0$, then

$$\text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\mathcal{F}'', \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})) = 0, \quad \mathcal{F}'' \in \text{D-mod}(\text{Bun}_G)_{\text{cusp}}.$$

By Proposition 2.3.2, it suffices to show that

$$j^?(\mathcal{F}) = 0 \Rightarrow j^* \circ \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F}) = 0.$$

However, this follows from (the nearly tautological) [Ga2, Corollary 6.6.3]. □

4. THE STRANGE FUNCTIONAL EQUATION AND PROOF OF THE EQUIVALENCE

In this section we will carry out the two main tasks of this paper: we will prove the strange functional equation (Theorem 4.1.2 below) and finish the proof of Theorem 3.1.5 (that says that the functor $\text{Ps-Id}_{\text{Bun}_G,!}$ is an equivalence).

4.1. The strange functional equation. In this subsection we will study the behavior of the functor $\text{Ps-Id}_{\text{Bun}_G,!}$ on the subcategory

$$\text{D-mod}(\text{Bun}_G)_{\text{co,Eis}} \subset \text{D-mod}(\text{Bun}_G)_{\text{co}}.$$

4.1.1. First, we have the following “strange” result:

Theorem 4.1.2. *For a parabolic P and its opposite P^- we have a canonical isomorphism of functors*

$$\text{Eis}_! \circ \text{Ps-Id}_{\text{Bun}_M,!} \simeq \text{Ps-Id}_{\text{Bun}_G,!} \circ \text{Eis}_{\text{co},*}^-.$$

Proof. Both sides are continuous functors

$$\text{D-mod}(\text{Bun}_M)_{\text{co},*} \rightarrow \text{D-mod}(\text{Bun}_G),$$

that correspond to objects of

$$\text{D-mod}(\text{Bun}_M \times \text{Bun}_G)$$

under the identification

$$\begin{aligned} \text{Funct}_{\text{cont}}(\text{D-mod}(\text{Bun}_M)_{\text{co}}, \text{D-mod}(\text{Bun}_G)) &\simeq (\text{D-mod}(\text{Bun}_M)_{\text{co}})^\vee \otimes \text{D-mod}(\text{Bun}_G) \simeq \\ &\simeq \text{D-mod}(\text{Bun}_M) \otimes \text{D-mod}(\text{Bun}_G) \simeq \text{D-mod}(\text{Bun}_M \times \text{Bun}_G), \end{aligned}$$

We claim that both objects identify canonically with

$$((\mathfrak{q} \times \mathfrak{p}) \circ \Delta_{\text{Bun}_P})!(k_{\text{Bun}_P}),$$

where the map in the formula is the same as

$$\mathrm{Bun}_P \xrightarrow{\mathfrak{q} \times \mathfrak{p}} \mathrm{Bun}_M \times \mathrm{Bun}_G.$$

The functor $\mathrm{Eis}_! \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_M, !}$ corresponds to the object, obtained by applying the functor $(\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)} \otimes \mathrm{Eis}_!) : \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G)$ to

$$(\Delta_{\mathrm{Bun}_M})!(k_{\mathrm{Bun}_M}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_M \times \mathrm{Bun}_M) \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_M).$$

The functor $\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)} \otimes \mathrm{Eis}_!$ is left adjoint to the functor

$$\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)} \otimes \mathrm{CT}_* \simeq (\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{q})_* \circ (\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{p})^!,$$

and hence is the !-Eisenstein series functor for the group $M \times G$ with respect to the parabolic $M \times P$. I.e., it is given by

$$(\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{p})! \times (\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{q})^*,$$

when applied to holonomic objects.

Base change along the diagram

$$\begin{array}{ccccc} \mathrm{Bun}_P & \xrightarrow{\Gamma_{\mathfrak{q}}} & \mathrm{Bun}_M \times \mathrm{Bun}_P & \xrightarrow{\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{p}} & \mathrm{Bun}_M \times \mathrm{Bun}_G \\ \mathfrak{q} \downarrow & & \downarrow \mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{q} & & \\ \mathrm{Bun}_M & \xrightarrow{\Delta_{\mathrm{Bun}_M}} & \mathrm{Bun}_M \times \mathrm{Bun}_M & & \end{array}$$

shows that

$$(\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{p})! \times (\mathrm{id}_{\mathrm{Bun}_M} \times \mathfrak{q})^* \circ (\Delta_{\mathrm{Bun}_M})!(k_{\mathrm{Bun}_M}) \simeq ((\mathfrak{q} \times \mathfrak{p}) \circ \Delta_{\mathrm{Bun}_P})!(k_{\mathrm{Bun}_P}),$$

as required.

The functor $\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G, !} \circ \mathrm{Eis}_{\mathrm{co}, *}^-$ corresponds to the object, obtained by applying the functor

$$\begin{aligned} & ((\mathrm{Eis}_{\mathrm{co}, *}^-)^\vee \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}) : \\ & \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \rightarrow \mathrm{D}\text{-mod}(\mathrm{Bun}_M) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \end{aligned}$$

to the object

$$(\Delta_{\mathrm{Bun}_G})!(k_{\mathrm{Bun}_G}) \in \mathrm{D}\text{-mod}(\mathrm{Bun}_G \times \mathrm{Bun}_G) \simeq \mathrm{D}\text{-mod}(\mathrm{Bun}_G) \otimes \mathrm{D}\text{-mod}(\mathrm{Bun}_G).$$

We have:

$$(\mathrm{Eis}_{\mathrm{co}, *}^-)^\vee \simeq \mathrm{CT}_*^-,$$

and we recall that by Theorem 1.1.7

$$\mathrm{CT}_*^- \simeq \mathrm{CT}_! := \bigoplus_{\mu} \mathrm{CT}_!^{\mu},$$

where $\mathrm{CT}_!^{\mu}$ is the left adjoint of Eis_*^{μ} .

Since $\mathrm{CT}_!^{\mu}$ is the left adjoint of Eis_*^{μ} , we obtain that $\mathrm{CT}_!^{\mu} \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}$ is the left adjoint of $\mathrm{Eis}_*^{\mu} \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}$, i.e., is the !-constant term functor for the group $G \times G$ with respect to the parabolic $P \times G$. Hence,

$$\mathrm{CT}_!^{\mu} \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)} \simeq (\mathfrak{q}^{\mu} \times \mathrm{id}_{\mathrm{Bun}_G})! \circ (\mathfrak{p}^{\mu} \times \mathrm{id}_{\mathrm{Bun}_G})^*,$$

when applied to holonomic objects (the superscript μ indicates that we are taking only the μ -connected component of Bun_P).

Taking the direct sum over μ , we thus obtain

$$(\mathrm{Eis}_{\mathrm{co},*}^-)^\vee \otimes \mathrm{Id}_{\mathrm{D-mod}(\mathrm{Bun}_G)} \simeq (\mathfrak{q} \times \mathrm{id}_{\mathrm{Bun}_G})! \circ (\mathfrak{p} \times \mathrm{id}_{\mathrm{Bun}_G})^*,$$

when applied to holonomic objects.

Now, base change along the diagram

$$\begin{array}{ccccc} \mathrm{Bun}_P & \xrightarrow{\Gamma_p} & \mathrm{Bun}_P \times \mathrm{Bun}_G & \xrightarrow{\mathfrak{q} \times \mathrm{id}_{\mathrm{Bun}_G}} & \mathrm{Bun}_M \times \mathrm{Bun}_G \\ \mathfrak{p} \downarrow & & \downarrow \mathfrak{p} \times \mathrm{id}_{\mathrm{Bun}_G} & & \\ \mathrm{Bun}_G & \xrightarrow{\Delta_{\mathrm{Bun}_G}} & \mathrm{Bun}_G \times \mathrm{Bun}_G & & \end{array}$$

shows that

$$(\mathfrak{q} \times \mathrm{id}_{\mathrm{Bun}_G})! \circ (\mathfrak{p} \times \mathrm{id}_{\mathrm{Bun}_G})^* \circ (\Delta_{\mathrm{Bun}_G})!(k_{\mathrm{Bun}_G}) \simeq ((\mathfrak{q} \times \mathfrak{p}) \circ \Delta_{\mathrm{Bun}_P})!(k_{\mathrm{Bun}_P}),$$

as required. \square

4.1.3. By passing to dual functors in the isomorphism

$$(4.1) \quad \mathrm{Eis}_! \circ \mathrm{Ps-Id}_{\mathrm{Bun}_M,!} \simeq \mathrm{Ps-Id}_{\mathrm{Bun}_G,!} \circ \mathrm{Eis}_{\mathrm{co},*}^-$$

of Theorem 4.1.2, we obtain:

Corollary 4.1.4. *There is a canonical isomorphism*

$$(4.2) \quad \mathrm{Ps-Id}_{\mathrm{Bun}_M,!} \circ \mathrm{CT}_{\mathrm{co},?} \simeq \mathrm{CT}_*^- \circ \mathrm{Ps-Id}_{\mathrm{Bun}_G,!}.$$

4.1.5. Consider now the commutative diagram:

$$(4.3) \quad \begin{array}{ccc} \mathrm{D-mod}(\mathrm{Bun}_G)_{\mathrm{co}} & \xrightarrow{\mathrm{Ps-Id}_{\mathrm{Bun}_G,!}} & \mathrm{D-mod}(\mathrm{Bun}_G) \\ \mathrm{Eis}_{\mathrm{co},*} \uparrow & & \uparrow \mathrm{Eis}_!^- \\ \mathrm{D-mod}(\mathrm{Bun}_M)_{\mathrm{co}} & \xrightarrow{\mathrm{Ps-Id}_{\mathrm{Bun}_M,!}} & \mathrm{D-mod}(\mathrm{Bun}_M). \end{array}$$

By passing to the right adjoint functors along the vertical arrows, we obtain a natural transformation

$$(4.4) \quad \mathrm{Ps-Id}_{\mathrm{Bun}_M,!} \circ \mathrm{CT}_{\mathrm{co},?} \rightarrow \mathrm{CT}_*^- \circ \mathrm{Ps-Id}_{\mathrm{Bun}_G,!}.$$

We now claim:

Proposition 4.1.6. *The map (4.4) equals the map (4.2), and, in particular, is an isomorphism.*

4.2. **Proof of Proposition 4.1.6.** The proof of the proposition is *not* a formal manipulation, as its statement involves the isomorphism of Theorem 4.1.2 for the two different parabolics, namely, P and P^- . The corresponding geometric input is provided by Lemma 4.2.3 below.

4.2.1. Let us identify

$$\mathrm{CT}_*^- \simeq \mathrm{CT}_! \quad \text{and} \quad \mathrm{Eis}_{\mathrm{co},*} \simeq (\mathrm{CT}_!^-)^\vee$$

via Theorem 1.1.7.

Then the map

$$\mathrm{Ps-Id}_{\mathrm{Bun}_M,!} \circ \mathrm{CT}_{\mathrm{co},?} \rightarrow \mathrm{CT}_! \circ \mathrm{Ps-Id}_{\mathrm{Bun}_G,!},$$

corresponding to (4.4), equals by definition the composition

$$\begin{aligned}
\mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_M,!} \circ \mathrm{CT}_{\mathrm{co},?} &\rightarrow \mathrm{CT}_! \circ \mathrm{Eis}_!^- \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_M,!} \circ \mathrm{CT}_{\mathrm{co},?} \stackrel{(4.3)}{\simeq} \\
&\simeq \mathrm{CT}_! \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!} \circ \mathrm{Eis}_{\mathrm{co},*} \circ \mathrm{CT}_{\mathrm{co},?} \simeq \mathrm{CT}_! \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!} \circ (\mathrm{CT}_!^-)^\vee \circ (\mathrm{Eis}_!)^\vee = \\
&= \mathrm{CT}_! \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!} \circ (\mathrm{Eis}_! \circ \mathrm{CT}_!^-)^\vee \rightarrow \mathrm{CT}_! \circ \mathrm{Ps}\text{-}\mathrm{Id}_{\mathrm{Bun}_G,!},
\end{aligned}$$

where the first arrows comes from the unit of the $(\mathrm{Eis}_!^-, \mathrm{CT}_!)$ -adjunction, and the last arrow comes from the co-unit of the $(\mathrm{Eis}_!, \mathrm{CT}_!^-)$ -adjunction.

This corresponds to the following map of objects in $\mathrm{D}\text{-mod}(\mathrm{Bun}_G \times \mathrm{Bun}_M)$:

$$\begin{aligned}
(4.5) \quad &(\mathrm{Eis}_! \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)}) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)})!(k_{\mathrm{Bun}_M}) \rightarrow \\
&\rightarrow (\mathrm{Eis}_! \otimes (\mathrm{CT}_! \circ \mathrm{Eis}_!^-)) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)})!(k_{\mathrm{Bun}_M}) = \\
&= (\mathrm{Eis}_! \otimes \mathrm{CT}_!) \circ (\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)} \otimes \mathrm{Eis}_!^-) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)})!(k_{\mathrm{Bun}_M}) \simeq \\
&\simeq (\mathrm{Eis}_! \otimes \mathrm{CT}_!) \circ (\mathrm{CT}_!^- \circ \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)})!(k_{\mathrm{Bun}_G}) = \\
&= ((\mathrm{Eis}_! \circ \mathrm{CT}_!^-) \otimes \mathrm{CT}_!) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)})!(k_{\mathrm{Bun}_G}) \rightarrow \\
&\rightarrow (\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)} \otimes \mathrm{CT}_!) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)})!(k_{\mathrm{Bun}_G}),
\end{aligned}$$

where the isomorphism between the 3rd and the 4th lines is

$$\begin{aligned}
&(\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)} \otimes \mathrm{Eis}_!^-) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)})!(k_{\mathrm{Bun}_M}) \simeq \\
&\simeq ((\mathfrak{q}^- \times \mathfrak{p}^-) \circ \Delta_{\mathrm{Bun}_{P^-}})!(k_{\mathrm{Bun}_{P^-}}) \simeq \\
&\simeq (\mathrm{CT}_!^- \circ \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)}) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)})!(k_{\mathrm{Bun}_G}),
\end{aligned}$$

used in the proof of Theorem 4.1.2.

The assertion of the proposition amounts to showing that the composed map in (4.5) equals

$$\begin{aligned}
&(\mathrm{Eis}_! \otimes \mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)}) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)})!(k_{\mathrm{Bun}_M}) \simeq \\
&\simeq ((\mathfrak{p} \times \mathfrak{q}) \circ \Delta_{\mathrm{Bun}_P})!(k_{\mathrm{Bun}_P}) \simeq \\
&\simeq (\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)} \otimes \mathrm{CT}_!) \circ (\Delta_{\mathrm{D}\text{-mod}(\mathrm{Bun}_G)})!(k_{\mathrm{Bun}_G}).
\end{aligned}$$

4.2.2. The geometric input is provided by the following assertion, proved at the end of this subsection:

Lemma 4.2.3. *The following diagram commutes:*

$$\begin{array}{ccc}
(\Delta_M)!(k_M) & \longrightarrow & (\mathrm{Id}_M \otimes (\mathrm{CT}_! \circ \mathrm{Eis}_!^-)) \circ (\Delta_M)!(k_M) \\
\downarrow & & \downarrow \sim \\
((\mathrm{CT}_!^- \circ \mathrm{Eis}_!) \otimes \mathrm{Id}_M) \circ (\Delta_M)!(k_M) & & (\mathrm{Id}_M \otimes \mathrm{CT}_!) \circ (\mathrm{Id}_M \otimes \mathrm{Eis}_!^-) \circ (\Delta_M)!(k_M) \\
\sim \downarrow & & \downarrow \sim \\
(\mathrm{CT}_!^- \otimes \mathrm{Id}_M) \circ (\mathrm{Eis}_! \otimes \mathrm{Id}_M) \circ (\Delta_M)!(k_M) & & (\mathrm{Id}_M \otimes \mathrm{CT}_!) \circ (\mathrm{CT}_!^- \otimes \mathrm{Id}_G) \circ (\Delta_G)!(k_G) \\
\sim \downarrow & & \downarrow \sim \\
(\mathrm{CT}_!^- \otimes \mathrm{Id}_M) \circ (\mathrm{Id}_G \otimes \mathrm{CT}_!) \circ (\Delta_G)!(k_G) & \xrightarrow{\sim} & (\mathrm{CT}_!^- \otimes \mathrm{CT}_!) \circ (\Delta_G)!(k_G)
\end{array}$$

where we use short-hand $\mathrm{Id}_M, \Delta_M, k_M$ for $\mathrm{Id}_{\mathrm{D}\text{-mod}(\mathrm{Bun}_M)}, \Delta_{\mathrm{Bun}_M}$ and k_{Bun_M} , respectively, and similarly for G .

Using the lemma, we rewrite the map in (4.5) as follows:

$$\begin{aligned}
& (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \rightarrow \\
& \quad \rightarrow ((\text{Eis}_! \circ \text{CT}_!^- \circ \text{Eis}_!) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) = \\
& = ((\text{Eis}_! \circ \text{CT}_!^-) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \simeq \\
& \simeq ((\text{Eis}_! \circ \text{CT}_!^-) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\text{Id}_{\text{D-mod}(\text{Bun}_G)} \otimes \text{CT}_!) \circ (\Delta_{\text{D-mod}(\text{Bun}_G)})_!(k_{\text{Bun}_G}) \rightarrow \\
& \quad \rightarrow (\text{Id}_{\text{D-mod}(\text{Bun}_G)} \otimes \text{CT}_!) \circ (\Delta_{\text{D-mod}(\text{Bun}_G)})_!(k_{\text{Bun}_G}),
\end{aligned}$$

and further as

$$\begin{aligned}
& (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \rightarrow \\
& \quad \rightarrow ((\text{Eis}_! \circ \text{CT}_!^- \circ \text{Eis}_!) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) = \\
& = ((\text{Eis}_! \circ \text{CT}_!^-) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \rightarrow \\
& \quad \rightarrow (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \simeq \\
& \quad \simeq (\text{Id}_{\text{D-mod}(\text{Bun}_G)} \otimes \text{CT}_!) \circ (\Delta_{\text{D-mod}(\text{Bun}_G)})_!(k_{\text{Bun}_G}).
\end{aligned}$$

However, the composition

$$\begin{aligned}
& (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \rightarrow \\
& \quad \rightarrow ((\text{Eis}_! \circ \text{CT}_!^- \circ \text{Eis}_!) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) = \\
& = ((\text{Eis}_! \circ \text{CT}_!^-) \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M}) \rightarrow \\
& \quad \rightarrow (\text{Eis}_! \otimes \text{Id}_{\text{D-mod}(\text{Bun}_M)}) \circ (\Delta_{\text{D-mod}(\text{Bun}_M)})_!(k_{\text{Bun}_M})
\end{aligned}$$

is the identity map, as it is induced by the map

$$\text{Eis}_! \rightarrow \text{Eis}_! \circ \text{CT}_!^- \circ \text{Eis}_! \rightarrow \text{Eis}_!,$$

comprised by the unit and co-unit of the $(\text{Eis}_!, \text{CT}_!^-)$ -adjunction, and the assertion follows.

4.2.4. *Proof of Lemma 4.2.3.* Let us recall from [DrGa3, Sect. 1.3.2] that the unit for the $(\text{Eis}_!, \text{CT}_!^-)$ can be described as follows. The functor

$$\text{CT}_! \circ \text{Eis}_!^- : \text{D-mod}(\text{Bun}_M) \rightarrow \text{D-mod}(\text{Bun}_M)$$

is given by

$$(\mathfrak{q})_! \circ (\mathfrak{p})^* \circ (\mathfrak{p}^-)_! \circ (\mathfrak{q}^-)^*,$$

which by base change along the diagram

$$\begin{array}{ccccc}
 & & \text{Bun}_M & & \\
 & & \downarrow \mathbf{j} & & \\
 & & \text{Bun}_{P^-} \times_{\text{Bun}_G} \text{Bun}_P & & \\
 & \text{id} \swarrow & & \searrow \text{id} & \\
 & \text{Bun}_{P^-} & & \text{Bun}_P & \\
 & \swarrow \mathbf{q}^- & & \swarrow \mathbf{p} & \searrow \mathbf{q} \\
 \text{Bun}_M & & \text{Bun}_G & & \text{Bun}_M
 \end{array}$$

can be rewritten as

$$(\mathbf{q})_! \circ (\mathbf{p}^-)_! \circ (\mathbf{p})^* \circ (\mathbf{q}^-)^*.$$

The natural transformation

$$\text{Id}_{\text{D-mod}(\text{Bun}_M)} \rightarrow \text{CT}_! \circ \text{Eis}_!^-$$

is given by

$$(\text{id}_{\text{Bun}_M})_! \circ (\text{id}_{\text{Bun}_M})^* = (\mathbf{q})_! \circ (\mathbf{p}^-)_! \circ \mathbf{j}_! \circ \mathbf{j}^* \circ (\mathbf{p})^* \circ (\mathbf{q}^-)^* \rightarrow (\mathbf{q})_! \circ (\mathbf{p}^-)_! \circ (\mathbf{p})^* \circ (\mathbf{q}^-)^*,$$

where the second arrow comes from the $(\mathbf{j}_!, \mathbf{j}^*)$ -adjunction.

The natural transformation

$$\text{Id}_{\text{D-mod}(\text{Bun}_M)} \rightarrow \text{CT}_!^- \circ \text{Eis}_!$$

is described similarly, with the roles of P and P^- swapped.

Base change along

$$\begin{array}{ccc}
 \text{Bun}_{P^-} \times_{\text{Bun}_G} \text{Bun}_P & \longrightarrow & \text{Bun}_{P^-} \times \text{Bun}_P \xrightarrow{\mathbf{q}^- \times \mathbf{q}} \text{Bun}_M \times \text{Bun}_M \\
 \downarrow & & \downarrow \mathbf{p}^- \times \mathbf{p} \\
 \text{Bun}_G & \xrightarrow{\Delta_{\text{Bun}_G}} & \text{Bun}_G \times \text{Bun}_G
 \end{array}$$

implies that the object

$$(\text{CT}_!^- \otimes \text{CT}_!) \circ (\Delta_{\text{Bun}_G})_!(k_{\text{Bun}_G}) \in \text{D-mod}(\text{Bun}_M \times \text{Bun}_M)$$

identifies with

$$(\mathbf{q}^- \times_{\text{Bun}_G} \mathbf{q})_!(k_{\text{Bun}_{P^-} \times_{\text{Bun}_G} \text{Bun}_P}),$$

where $\mathbf{q}^- \times_{\text{Bun}_G} \mathbf{q}$ denotes the map

$$\text{Bun}_{P^-} \times_{\text{Bun}_G} \text{Bun}_P \rightarrow \text{Bun}_{P^-} \times \text{Bun}_P \xrightarrow{\mathbf{q}^- \times \mathbf{q}} \text{Bun}_M \times \text{Bun}_M.$$

Now, the above description of the unit of the adjunctions implies that both circuits in the diagram in Lemma 4.2.3 are equal to the map

$$(\Delta_{\text{Bun}_M})!(k_{\text{Bun}_M}) \rightarrow (\mathfrak{q}^- \times_{\text{Bun}_G} \mathfrak{q})!(k_{\text{Bun}_{P^-}} \times_{\text{Bun}_G} \text{Bun}_P),$$

that corresponds to the open embedding

$$\text{Bun}_M \xrightarrow{\mathbf{j}} \text{Bun}_{P^-} \times_{\text{Bun}_G} \text{Bun}_P.$$

□

4.3. Proof of Theorem 3.1.5. We are finally ready to prove Theorem 3.1.5.

We proceed by induction on the semi-simple rank of G . The case of a torus follows immediately from Corollary 3.3.3. Hence, we will assume that the assertion holds for all proper Levi subgroups of G .

4.3.1. Theorem 4.1.2, together with the induction hypothesis, imply that the essential image of $\text{D-mod}(\text{Bun}_G)_{\text{co,Eis}}$ under $\text{Ps-Id}_{\text{Bun}_G,!}$ generates $\text{D-mod}(\text{Bun}_G)_{\text{Eis}}$.

Corollary 3.3.3 implies that the essential image of $\text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}$ under $\text{Ps-Id}_{\text{Bun}_G,!}$ generates (in fact, equals) $\text{D-mod}(\text{Bun}_G)_{\text{cusp}}$.

Hence, it remains to show that $\text{Ps-Id}_{\text{Bun}_G,!}$ is fully faithful.

4.3.2. The fact that $\text{Ps-Id}_{\text{Bun}_G,!}$ induces an isomorphism

$$(4.6) \quad \text{Hom}_{\text{D-mod}(\text{Bun}_G)_{\text{co}}}(\mathcal{F}', \mathcal{F}) \rightarrow \text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F}'), \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F}))$$

for $\mathcal{F}' \in \text{D-mod}(\text{Bun}_G)_{\text{co,cusp}}$ follows from Proposition 3.3.5.

Hence, it remains to show that (4.6) is an isomorphism for $\mathcal{F}' \in \text{D-mod}(\text{Bun}_G)_{\text{co,Eis}}$. The latter amounts to showing that the functor $\text{Ps-Id}_{\text{Bun}_G,!}$ induces an isomorphism

$$\begin{aligned} \text{Hom}_{\text{D-mod}(\text{Bun}_G)_{\text{co}}}(\text{Eis}_{\text{co},*}(\mathcal{F}_M), \mathcal{F}) &\rightarrow \\ &\rightarrow \text{Hom}_{\text{D-mod}(\text{Bun}_G)}(\text{Ps-Id}_{\text{Bun}_G,!} \circ \text{Eis}_{\text{co},*}(\mathcal{F}_M), \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})) \end{aligned}$$

for $\mathcal{F}_M \in \text{D-mod}(\text{Bun}_M)_{\text{co}}$ for a *proper* parabolic P with Levi quotient M .

4.3.3. Note that for $\mathcal{F}_M \in \text{D-mod}(\text{Bun}_M)_{\text{co}}$ and $\mathcal{F} \in \text{D-mod}(\text{Bun}_G)_{\text{co}}$ we have a commutative diagram:

$$\begin{array}{ccc} \text{Hom}(\text{Eis}_{\text{co},*}(\mathcal{F}_M), \mathcal{F}) & \longrightarrow & \text{Hom}(\text{Ps-Id}_{\text{Bun}_G,!} \circ \text{Eis}_{\text{co},*}(\mathcal{F}_M), \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})) \\ & & \begin{array}{c} (4.3) \downarrow \sim \\ \text{Hom}(\text{Eis}_!^- \circ \text{Ps-Id}_{\text{Bun}_M,!}(\mathcal{F}_M), \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})) \\ \sim \downarrow \\ \text{Hom}(\text{Ps-Id}_{\text{Bun}_M,!}(\mathcal{F}_M), \text{CT}_*^- \circ \text{Ps-Id}_{\text{Bun}_G,!}(\mathcal{F})) \\ (4.4) \uparrow \end{array} \\ \sim \downarrow & & \text{Hom}(\mathcal{F}_M, \text{CT}_{\text{co},?}(\mathcal{F})) \longrightarrow \text{Hom}(\text{Ps-Id}_{\text{Bun}_M,!}(\mathcal{F}_M), \text{Ps-Id}_{\text{Bun}_M,!} \circ \text{CT}_{\text{co},?}(\mathcal{F})). \end{array}$$

The bottom horizontal arrow in the above diagram is an isomorphism by the induction hypothesis. Now, Proposition 4.1.6 implies that the lower right vertical arrow is also an isomorphism.

Hence, the upper horizontal arrow is also an isomorphism, as required.

REFERENCES

- [AG] D. Arinkin and D. Gaitsgory, *Singular support of coherent sheaves, and the geometric Langlands conjecture*, arXiv:1201.6343.
- [BG] A. Braverman, D. Gaitsgory, *Geometric Eisenstein Series*, Invent. Math **150** (2002), 287–384. Also: arXiv:math/9912097.
- [BFGM] A. Braverman, M. Finkelberg, D. Gaitsgory and I. Mirkovic, *Intersection cohomology of Drinfeld compactifications*, Selecta Math (N.S.) **8** (2002), 381–418. Also: arXiv:math/0012129.
- [DrGa1] V. Drinfeld and D. Gaitsgory, *On some finiteness questions for algebraic stacks*, Geometric and Functional Analysis **23** (2013), 149–294. Also: arXiv:1108.5351.
- [DrGa2] V. Drinfeld and D. Gaitsgory, *Compact generation of the category of D -modules on the stack of G -bundles on a curve*, arXiv:1112.2402.
- [DrGa3] V. Drinfeld and D. Gaitsgory, *Geometric constant term functor(s)*, arXiv:1301.2071.
- [Ga1] D. Gaitsgory, *Outline of the proof of the geometric Langlands conjecture for GL_2* , arXiv:1302.2506
- [Ga2] D. Gaitsgory, *Functors given by kernels, adjunctions, and duality*, arXiv:1303.2763.
- [Sch] S. Schieder, in preparation.
- [Vi] E. B. Vinberg, *On reductive algebraic semigroups*, E. B. Dynkins Seminar, 145–182, Amer. Math. Soc. Transl. Ser. 2 **169**, Amer. Math. Soc., Providence, RI, 1995.