

DIFFERENTIAL GEOMETRY

MATH 136

Lecture 18: Riemannian Manifolds

18.1. A **topological manifold** M of dimension m is a subset of some \mathbb{R}^n such that every $x \in M$ has a neighborhood U , that is homeomorphic to an open subset $R = \phi(U)$ of \mathbb{R}^m . The pair (U, ϕ) is called a **chart**. It produces a **coordinate system** on U : there is a parametrization $r(\phi(x)) = x$, which is a regular map from $R \subset \mathbb{R}^m$ to \mathbb{R}^n , meaning that dr has rank m everywhere. A C^k **atlas** on M is a collection $\mathcal{F} = \{U_i, \phi_i\}_{i \in I}$ of charts such that $\bigcup_{i \in I} U_i = M$, and that all coordinate change maps $\phi_{ij} = \phi_i \circ \phi_j^{-1}$ are in $C^k(\phi_i(U_j \cap U_i), \mathbb{R}^m)$. An atlas is called **maximal**, if (U, ϕ) is a chart such that $\phi \circ \phi_i^{-1}$ and $\phi_i \circ \phi^{-1}$ are all C^k for all $i \in I$, then $(U, \phi) \in \mathcal{F}$. Two atlases \mathcal{F}, \mathcal{G} are called **equivalent** if their union $\mathcal{F} \cup \mathcal{G}$ is an atlas. Given an atlas \mathcal{F} , the union of all atlases equivalent to \mathcal{F} is called a **differentiable structure of \mathcal{F}** .¹




FIGURE 1. A m -manifold $M \subset \mathbb{R}^n$ is shown with part of an atlas \mathcal{F} . Each patch U_i is a regularly parametrized by $r : R_i \rightarrow U_i$ with $R_i = \phi_i(U_i) \subset \mathbb{R}^m$. The map r has maximal rank m everywhere on R_i .

18.2. A m -dimensional C^k -**differentiable manifold** is a pair (M, \mathcal{F}) , where M is a m -dimensional topological manifold and \mathcal{F} is a differentiable C^k structure on M . This means is that every point $p \in M$ is contained in a region $U = \phi(R)$. The inverse r of ϕ defines a parametrization from $R \subset \mathbb{R}^m$ to $M \subset \mathbb{R}^n$. We are now in the frame

¹The concept can be difficult: \mathcal{F} is not unique in general. On S^7 , there are 28 different smooth structures. The smooth Poincare conjecture claims that S^4 has a unique differentiable structure.

work used before. We can for example define r_u, r_v and the fundamental forms. We say **smooth** if we just want to be able to differentiate as often as necessary. ²

18.3. If $E = \mathbb{R}^m$ is the space of **column vectors** of dimension m , its **dual** E^* is defined as the space of all linear maps $f : E \rightarrow \mathbb{R}$. It is the space of **row vectors**. If $\{e_1, \dots, e_m\}$ is a basis of E , then $\{e^1, \dots, e^m\}$ denotes a basis of E^* . Every element in E can be written as $v = \sum_i v^i e_i$, every element in E^* can be written as $v = \sum_i v_i e^i$. For $p, q \geq 0$, the linear space T_q^p of all multi-linear maps $(E^*)^p \times E^q \rightarrow \mathbb{R}$ is called the space of **tensors of type** (p, q) . Column vectors are $(1, 0)$ -tensors in $T_0^1 = E$, while row vectors are $(0, 1)$ -tensors in $T_1^0 = E^*$, bilinear maps are $(0, 2)$ tensors in T_2^0 . A **tensor field of type** (p, q) on a m -manifold M is a smooth assignment of a (p, q) tensor to every point. Such a map is also called a **section** of the tensor bundle, generalizing that a **vector field**. A vector field is a section of the **tangent bundle** TM . For a $(0, 2)$ tensor field g for example, the attribute "smooth" means that for any vector fields X, Y , the function $x \rightarrow g(x)(X(x), Y(x))$ is smooth from M to \mathbb{R} . If $f : M \rightarrow \mathbb{R}^k$ is a smooth map, then df is a $(0, 1)$ tensor field. This is also called a **1-form**. **Vector field** is an abbreviation for a $(1, 0)$ tensor field. The first fundamental form g is by definition a $(2, 0)$ tensor field; think of a symmetric bilinear form attached to every point. A **Riemannian manifold** (M, g) is a smooth manifold M with a positive definite symmetric $(2, 0)$ tensor field g .

18.4. Let M be a m -manifold and $f : M \rightarrow \mathbb{R}^k$ be smooth. A point $x \in M$ is called a **critical point** and $f(x)$ a **critical value**, if the rank of $df(x)$ is not m . Non-critical points are called **regular points**.

Theorem 1. *If M is a m -manifold and $f : M \rightarrow \mathbb{R}^k$ is smooth and y is a regular value of f , then $M_f = f^{-1}(y)$ is a manifold of dimension $m - k$.*

Proof. At $x \in f^{-1}(y)$, the Jacobean map $df(x)$ has rank k and the kernel $H = \ker(df)$ of $df(x)$ is $(m - k)$ -dimensional and H^\perp is k dimensional. Take a chart (U, ϕ) in M which contains x . It defines a parametrization $r = \phi^{-1} : R \rightarrow U$. Define $g = f \circ r : R \subset \mathbb{R}^m \rightarrow \mathbb{R}^k$. Define eh projection $L : \mathbb{R}^m = H \oplus H^\perp \rightarrow \mathbb{R}^{m-k}, (h, h') \mapsto h'$ onto the orthogonal complement of the kernel. The map $F : R \subset \mathbb{R}^m \rightarrow \mathbb{R}^m = \mathbb{R}^k \times \mathbb{R}^{m-k}$ defined by $F(z) = (g(z), Lz)$ has the non-singular derivative $dF(u) = (dg(u), Lu)$. By the **inverse function theorem**, a neighborhood of $u = \phi(x)$ is mapped by the diffeomorphism F onto a neighborhood $F(y)$ of $F(u)$. We get so a chart $U_x = r(R_x)$, where $R_x = F^{-1}(H)$. It is a chart of M_f . Doing the same construction at any point $x \in M$ produces an atlas for $f^{-1}(y)$ and verifies that $f^{-1}(y)$ is a manifold. \square

18.5. Examples: a) The **d - sphere** is the set $M = S^d = \{x \in \mathbb{R}^{d+1} \mid x_1^2 + x_2^2 + \dots + x_{d+1}^2 = 1\} \subset \mathbb{R}^{d+1}$. The **standard differentiable structure** \mathcal{F} on S^d is generated by $\mathcal{F} = \{(S^d \setminus \{A\}, \phi_A), (S^d \setminus \{B\}, \phi_B)\}$, where $\phi_A \phi_B$ are **stereographic projections** from antipodal points A, B .

b) The set $SL(n, \mathbb{R})$ of $n \times n$ matrices of determinant 1 is a manifold.

OLIVER KNILL, KNILL@MATH.HARVARD.EDU, MATH 136, FALL, 2025

²A theorem of Whitney assures that a smooth compact m -manifold M given abstractly as paracompact Hausdorff space can be realized within \mathbb{R}^n if $n = 2m + 1$.