

# DIFFERENTIAL GEOMETRY

MATH 136

## Lecture 19: Discrete Manifolds

**19.1.** A **discrete m-manifold** is a finite graph  $G = (V, E)$  for which every unit sphere  $S(v)$  is a discrete  $(m-1)$ -sphere. A **discrete m-sphere** is a discrete m-manifold which has the property that removing a point renders it contractible. Inductively, a graph is called **contractible**, if both  $S(v)$  and  $S \setminus v$  are contractible for some  $v \in V$ . The 1-point space 1 is contractible. The empty graph is the  $(-1)$ -sphere. Let  $F_k$  denote the set of  $K_{k+1}$  subgraphs ( $k$ -simplices) and  $f_k = |F_k|$ . We have  $F_0 = V, F_1 = E$ . The **Euler characteristic** of  $M$  is defined as  $\chi(M) = \sum_{k=0}^m (-1)^k f_k = f_0 - f_1 + f_2 - f_3 + \dots + (-1)^m f_m$ . This definition of Ludwig Schläfli generalizes  $\chi(M) = f_0 - f_1 + f_2 = V - E + F$  for 2-manifolds.




FIGURE 1. This 2-manifold  $M$  of genus  $g = 2$  has  $\chi(M) = 2 - 2g = -2$ .

**19.2.** A graph without edges is a 0-manifold. A 0-manifold with 2 points is a 0-sphere. The reason is that removing a vertex produces the contractible  $K_1$ . Every connected 1-manifold is a 1-sphere, a circular graph  $C_n$  with  $n \geq 4$ . Every finite 2-manifold is either a 2-sphere  $S^2$  or a connected sum of tori or projective planes:  $M = S^2, M = \mathbb{T}^2 \# \dots \# \mathbb{T}^2$  or  $M = \mathbb{P}^2 \# \dots \# \mathbb{P}^2$ . A 2-sphere can be characterized as 2-manifold of Euler characteristic 2. The 16 cell and the 600 cells are examples of 3-spheres. The join of two 1-spheres is a 3-sphere. The join of a  $k$ -sphere with a  $m$ -sphere is a  $(k+m+1)$ -sphere. The join of  $G$  with the 0-sphere is called **suspension**.

**19.3.** Euler's formula  $\chi(M) = V - E + F = 2$  for 2-spheres generalizes to higher dimension. The 0-sphere has  $\chi(M) = V = 2$ , every 1-sphere has  $\chi(M) = V - E = 0$ . Every 2-sphere has  $\chi(M) = V - E + F = 2$ . This pattern continues:

**Theorem 1** (Euler's Gem). *If  $M$  is a  $m$ -sphere, then  $\chi(M) = 1 + (-1)^m$ .*

*Proof.* Use induction with respect to dimension  $m$ . For  $m = 0$ , we have  $\chi(M) = 2$ . The induction assumption is that all  $(m - 1)$ -spheres  $S$  satisfy  $\chi(S) = 1 + (-1)^{m-1}$ . Pick a vertex  $v$ . As the unit sphere  $S(v)$  is a  $(m - 1)$ -sphere and  $S(v) = B(v) \cap G \setminus v$ , where both the unit ball  $B(v)$  and  $G \setminus v$  are contractible with Euler characteristic 1, we have, using the induction assumption,  $\chi(M) = \chi(G \setminus v) + \chi(B(v)) - \chi(G \setminus v \cap B(v)) = 2 - (1 - (-1)^{m-1}) = 1 + (-1)^m$ .  $\square$

**19.4.** In the continuum, manifolds can be constructed as level surfaces of functions like  $x^2 + y^2 + z^2 = 1$ . We can do that also in the discrete. Take an arbitrary function  $f : V \rightarrow Z_k = \{0, \dots, k\}$ . It defines a new graph  $M_f$ , where the vertices are the set of complete subgraphs on which  $f$  attains all  $k$  values. Connect two of these points by an edge, if one is contained in the other. The new graph  $M_f$  is a sub-graph of the **Barycentric refinement** of  $M$ . There is the analog of what we have seen classically for functions on manifolds.

**Theorem 2** (Level Sets). *If  $M$  is a  $m$ -manifold and  $f : M \rightarrow Z_k$  is an arbitrary function, then either  $M_f$  is empty or then  $M_f$  is a  $(m - k)$ -manifold.*

*Proof.* Let  $x$  be a  $n$ -simplex on which  $f$  takes all values. This means  $f(x) = Z_k$ . The graph  $S^-(x) = \{y \subset x, y \neq x\}$  is a  $(n - 1)$ -sphere in the Barycentric refinement of  $M$ . The simplices in  $S^-(x)$  on which  $f$  still reaches  $Z_k$  is by induction a  $(n - 1 - k)$ -manifold and since we are in a simplex, it has to be a  $(n - 1 - k)$ -sphere. Every unit sphere  $S(x)$  in the Barycentric refinement is a  $(m - 1)$ -sphere as it is the join of  $S^-(x)$  with  $S^+(x) = \{y, x \subset y, x \neq y\}$ . (The join of two spheres is always a sphere.) The sphere  $S_f^+(x)$  in  $M_f$  is the same than  $S^+(x)$  in  $M$  because every simplex  $z$  in  $M$  containing  $x$  automatically has the property that  $f(z) = Z_k$ . So, the unit sphere  $S(x)$  in  $M_f$  is the join of a  $(n - k - 1)$ -sphere and the  $(m - n - 1)$ -sphere and so a  $(m - k - 1)$ -sphere. Having shown that every unit sphere in  $M_f$  is a  $(m - k - 1)$ -sphere, we see that  $M_f$  is a  $(m - k)$ -manifold.  $\square$

**19.5.** Differential geometry works too: define **curvature** as

$$K(v) = \sum_{k=0}^m \frac{(-1)^k f_{k-1}(S(v))}{k+1} = 1 - \frac{f_0(S(v))}{2} - \frac{f_1(S(v))}{3} + \dots$$

In the case of a 2-manifold, this boils down to  $1 - f_0(S(v))/2 + f_1(S(v))/3 = 1 - d(v)/6$ , where  $d(v)$  is the vertex degree. For odd-dimensional manifolds, the curvature is constant zero.

**Theorem 3** (General Gauss-Bonnet).  $\sum_{v \in V} K(v) = \chi(M)$

*Proof.* The proof is the same as in the 2-dimensional case. Again look at the energies  $\omega(x) = (-1)^{\dim(x)}$  attached to each simplex  $x$  in the graph (complete subgraph with  $\dim(x) + 1$  vertices). Then  $\chi(M) = \sum_x \omega(x)$ . Now distribute all these energies of a  $k$ -simplex  $x$  equally to the  $k + 1$  vertices contained in  $x$ . As there are  $f_{k-1}$  simplices in  $S(v)$  which correspond do simplices containing  $v$ , this adds  $\frac{(-1)^k f_{k-1}(S(v))}{k+1}$  to each vertex  $v$ . Now just collect all up at a vertex  $v$  to get the curvature  $K(v)$ . The transactions of energies preserved the total energy = Euler characteristic.  $\square$