

PROBABILITY THEORY

MATH 154

Unit 10: Problem Seminar

10.1. Finite probability taps into **combinatorics**. Here are some examples:

$n!$	There are $5! = 120$ possible ways to redistribute 5 coats to 5 people.
$\frac{n!}{n_1! \cdots n_k!}$	With $\{A, A, B, B, B, B, E, U, U, U\}$, form $10!/(2!4!1!3!)$ ten letter words.
$\frac{n!}{(n-k)!}$	10 people can sit $10!/6! = 10 * 9 * 8 * 7$ possible ways on 4 chairs.
n^k	There are 6^{10} possible ways to throw 10 dices.
$\frac{n!}{(n-k)!k!}$	There are $52!/(5!47!)$ hands of 5 cards a deck of 52.

- Problem 1:**
- You pick 7 cards at random from a deck of 81 cards of the game "set". What is the probability that all of them are red (27 are red)?
 - You throw a dice 7 times. What is the probability that all 7 numbers are even?
 - Your gym lock consist of 3 different numbers from 1 to 40. Having forgotten the number, you try a random combinations. What is the probability to open it?
 - How many bijective functions $X \rightarrow X$ are there on $X = \{1, 2, 3, 4, 5, 6, 7\}$?
 - How many functions $X \rightarrow Y$ are there from $X = \{1, 2, 3, 4, 5, 6, 7\}$ to $\{1, 2\}$?

10.2. The **laboratory** Ω is a set of experiments. The σ -**algebra** \mathcal{A} consists of events. A σ -algebra is a Boolean algebra which allows to perform **countably** many operations.

$A \cdot B = A \cap B = \{\omega \in \Omega \mid \omega \in A \text{ and } \omega \in B\}$	"Both events A and B happen"
$A \cup B = \{\omega \in \Omega \mid \omega \in A \text{ or } \omega \in B\}$	"Either A or B happens"
$A + B = A \Delta B = \{\omega \in \Omega \mid \omega \in A \text{ xor } \omega \in B\}$	"One of the events A or B happens"
$A \setminus B = \{\omega \in \Omega \mid \omega \in A \text{ but not } \omega \in B\}$	" A but not B happens"
$A^c = \{\omega \in \Omega \mid \omega \notin A\}$	" A does not happen"
$\bigcap_n A_n = \{\omega \in \Omega \mid \omega \in A_n, \text{ for all } n\}$	"All events A_n happen"
$\bigcup_n A_n = \{\omega \in \Omega \mid \omega \in A_n, \text{ for at least one } n\}$	"At least one event A_n happens"

Problem 2: We use the notation $A \cdot B = A \cap B$ and $A + B = A \Delta B$ and $1 = \Omega$ and $0 = \emptyset$ in the Boolean algebra $\mathcal{P} = 2^\Omega$ of all subsets of Ω .

- Draw the Venn diagram picture proving that $A(B - C) = AB - AC$.
- Simplify $(5A + 2)(3A^2 + A - 1)$.
- Write $A^n = A \cdot A \cdot A \cdots A$ for the n 'th power. Simplify $(A - 1)(A + A^2 + A^3)$.
- Why is $(1 + A)^3 = 1 + A$?
- Show that $A \cup B = A + B + AB$.

10.3. If B has positive probability, then $P[A|B] = P[A \cap B]/P[B]$ is called the **conditional probability** of A under the condition that event B takes place.

Problem 3: a) If the probability that a student is sick at a given day is 1 percent and the probability that a student has an exam at a given day is 5 percent. Suppose that 6 percent of the students with exams are ill. What is the probability that an ill student has an exam on a given day?
 b) Suppose that A, B are subsets of a sample space with a probability function P . We know that $P[A] = 4/5$ and $P[B] = 3/5$. Explain why $P[B|A]$ is at least $1/2$.

10.4. The linear space \mathcal{L}^2 has an **inner product** $\mathcal{X} \cdot \mathcal{Y} = E[XY]$ and so a **length** $|\mathcal{X}| = \sqrt{\mathcal{X} \cdot \mathcal{X}}$. The **standard deviation** of X is the length of **centered random variable** $X - E[X]$. The **correlation** $-1 \leq \text{Cov}[X, Y]/(\sigma[X]\sigma[Y]) \leq 1$ is $\cos(\alpha)$ and defines an **angle** α between $X - E[X]$ and $Y - E[Y]$. If X takes finitely many values (which means $X \in \mathcal{S}$), then $E[X^n] = \sum_{x \in X(\Omega)} x^n P[X = x]$. For $X \in \mathcal{L}^n$ with a PDF f , then $E[X^n] = \int_{-\infty}^{\infty} x^n f(x) dx$. In the box, c, λ are constants.

$E[X + Y] = E[X] + E[Y]$	$E[\lambda X] = \lambda E[X]$
$X \leq Y \Rightarrow E[X] \leq E[Y]$	$E[X^2] = 0 \Leftrightarrow X = 0$
$E[X] = c$ if $X(\omega) = c$	$E[X - E[X]] = 0$
$\text{Var}[X] \geq 0$	$\text{Var}[X] = E[X^2] - E[X]^2$
$\text{Var}[\lambda X] = \lambda^2 \text{Var}[X]$	$\text{Cov}[X, X] = \text{Var}[X]$

Problem 4: Let $([-\pi, \pi], \mathcal{B}, P)$ be the Lebesgue probability space, where $P[[a, b]] = (b - a)/(2\pi)$ on the π system of all half open intervals on $\Omega = [-\pi, \pi]$.
 a) Which theorem assures that the measure P exists?
 b) Let $X(x) = \sin(3x), Y(x) = \cos(3x)$. Compute the $E[X], E[Y], \sigma[X], \sigma[Y]$.
 c) What is the correlation $\text{Cor}[X, Y] = \text{Cov}[X, Y]/\sigma(X)\sigma(Y)$? Are X, Y independent?

10.5. Assume X has a probability density $F' = f$ then $E[X^n] = \int x^n f(x) dx$ and $E[e^{tX}] = \int e^{tx} f(x) dx, E[e^{itX}] = \int e^{itx} f(x) dx$. Now form $\text{Var}[X] = E[X^2] - E[X]^2$ etc.

Problem 5: The PDF $f(x) = \frac{2}{\pi\sqrt{1-x^2}}$ is supported on $[-1, 1]$.
 a) Compute the cumulative distribution function $F(x) = \int_{-\infty}^x f(t) dt = \int_{-1}^x f(t) dt$.
 b) Write down the integral for the moment generating function $M_X(t)$.
 c) Express the variance in terms of $M'_X(0)$ and $M''_X(0)$.
 d) Relate $M_X(t)$ and $M_Y(t)$ and $M_{X+Y}(t)$ for independent random variables! State the same law for the characteristic function $\phi_X(t), \phi_Y(t)$.

10.6. Assume X is a random variable taking a finite or countable number of values $P[X = x_k] = p_k$. Then $E[X^n] = \sum_k x_k^n p_k, E[e^{tX}] = \sum_k e^{tx_k} p_k$ and $E[e^{itX}] = \sum_k e^{itx_k} p_k$.

Problem 6: Assume X is a random variable that takes the value 3 with probability $1/3$ and the value 6 with probability $2/3$.
 a) Find the expectation $m = E[X]$ and $n = E[X^2]$.
 b) Are X, X^2 independent? Explain.
 c) Write down the characteristic function.