

5/7/2021: Final Practice B

”By signing, I affirm my awareness of the standards of the Harvard College Honor Code.”

Your Name:

- Solutions are submitted to knill@math.harvard.edu as PDF handwritten in a file carrying your name. Capitalize the first letters like in OliverKnill.pdf. The paper has to **feature your personal handwriting** and contain no typed part. If you like, you can start writing on a new paper. For 1), you could write 1: False, 2: False ... but you then need to copy the above Honor Code statement and sign.
- No books, calculators, computers, or other electronic aids are allowed. You can use a double sided page of your own handwritten notes when writing the paper. It is your responsibility to submit the paper on time and get within that time also a confirmation. The exam is due at 9 AM on May 8th. Do not communicate with anybody related to the class during the exam period and with nobody at all about the exam.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
Total:		140


Problem 1) TF questions (20 points). No justifications are needed.

- 1) T F $\cos(17\pi/4) = \sqrt{2}/2$.
- 2) T F The tangent function is monotonically increasing on the open interval $(-\pi/2, \pi/2)$.
- 3) T F The arccot function is monotonically increasing from $\pi/4$ to $3\pi/4$.
- 4) T F If f is a probability density function, then $\int_{-\infty}^{\infty} f(x) dx = 0$
- 5) T F $\frac{d}{dx} e^{\log(x)} = 1$.
- 6) T F If $f''(0) = -1$ then f has a local maximum at $x = 0$.
- 7) T F The improper integral $\int_{-1}^1 1/|x| dx$ is finite.
- 8) T F The function $-\cos(x) - x$ has a root in the interval $(-100, 100)$.
- 9) T F If a function f has a local maximum in $(0, 1)$ then it also has a local minimum in $(0, 1)$.
- 10) T F The anti derivative of $1/(1 - x^2)$ is equal to $\arctan(x)$.
- 11) T F The function $f(x) = (e^x - e^{2x})/(x - x^2)$ has the limit 1 as x goes to zero.
- 12) T F If you listen to the sound $e^{-x} \sin(10000x)$, then it gets louder and louder as time goes on.
- 13) T F The function $f(x) = e^{x^2}$ has a local minimum at $x = 0$
- 14) T F The function $f(x) = (x^{55} - 1)/(x - 1)$ has the limit 1 for $x \rightarrow 1$.
- 15) T F If the total cost $F(x)$ of an entity is extremal at x , then we have a break even point $f(x) = g(x)$.
- 16) T F The value $\int_{-\infty}^{\infty} xf(x) dx$ is called the expectation of the PDF f .
- 17) T F The trapezoid rule is an integration method in which the left and right Riemann sum are averaged.
- 18) T F $\tan(\pi/3) = \sqrt{3}$.
- 19) T F A Newton step for the function f is $T(x) = x + \frac{f(x)}{f'(x)}$.
- 20) T F $\sin(\arctan(1)) = \sqrt{3}$.


Problem 2) Matching problem (10 points) No justifications needed

(5 points) Match the functions names with their graphs (1-4) their derivatives (A-D) (middle row) and second derivatives (a-d) (last row).


Function	fill in 1)-4)	fill in A)-D)	fill in a)-d)
$\sin(x)/x$			
$\tan(x)$			
$\arcsin(x)$			
$1/(1+x^2)$			




1)




2)




3)




4)




A)




B)




C)




D)




a)



b)



c)




d)

(5 points) Which of the following limits exists in the limit $x \rightarrow 0$.

Function	exists	does not exist
$\sin^4(x)/x^4$		
$1/\log x $		
$\arctan(x)/x$		
$\log x /(x-1)$		
$\cos(x)/(x-1)$		
$(x^{10}-1)/(x-1)$		

Problem 3) Matching or short answer problem (10 points). No justifications are needed.

a) (4 points) On the Boston Esplanade is a sculpture of **Arthur Fiedler** (1894-1979) a long-time conductor of the Boston Pops Orchestra. His head is sliced into thin slices. Assume that the thickness of each level is $h = 1.5$ inch and the area of each of the 100 slices k is $A(k)$. Which formula gives the volume of the head? (One applies.)



Formula	Check if true
$1.5[A(1) + \dots + A(100)]$	<input type="checkbox"/>
$\frac{1}{1.5}[A(1) + \dots + A(100)]$	<input type="checkbox"/>

Formula	Check if true
$1.5[\frac{1}{A(1)} + \dots + \frac{1}{A(100)}]$	<input type="checkbox"/>
$\frac{1.5}{100}[A(1) + \dots + A(100)]$	<input type="checkbox"/>

b) (4 points) The summer has arrived on May 12 2014 for a day before it cooled down again. Harvard students enjoy the **Lampon pool** that day in front of the **Lampon castle**. Assume the water volume at height z is $V(z) = 1 + 5z - \cos(z)$. Assume water evaporates at a rate of $V'(z) = -1$ gallon per day. How fast does the water level drop at $z = \pi/2$ meters? Check the right answer: (one applies)



Rate	Check if true
-6	<input type="checkbox"/>
-1/6	<input type="checkbox"/>

Rate	Check if true
-4	<input type="checkbox"/>
-1/4	<input type="checkbox"/>

c) (2 points) Speaking of weather: the temperature on April 28, 2021 in Cambridge was 53 degrees Fahrenheit. Four days before, on April 24, the temperature had been 70 degrees and have us dream about **summer time**. Which of the following theorems assures that there was a moment during the night of April 24 to May 28 that the temperature was exactly 61.5 degrees? (One applies.)



Theorem	check if true
Mean value theorem	<input type="checkbox"/>
Fermat's theorem	<input type="checkbox"/>

Theorem	check if true
Intermediate value theorem	<input type="checkbox"/>
Bolzano theorem	<input type="checkbox"/>


Problem 4) Area computation (10 points)

Find the area enclosed by the graphs of the functions

$$f(x) = \log|x|$$

and

$$g(x) = \sqrt{1-x^2}.$$



Problem 5) Volume computation (10 points)

The lamps near the front entrance of the **Harvard Malkin Athletic Center (MAC)** have octagonal cross sections, where at height z , the area is

$$A(z) = 2(1 + \sqrt{2})(1 + z)^2$$

with $0 \leq z \leq 3$. What is the volume of the lamp?



Problem 6) Improper integrals (10 points)

Which of the following limits $R \rightarrow \infty$ exist? If the limit exist, compute it.

a) (2 points) $\int_1^R \sin(2\pi x) dx$

b) (2 points) $\int_1^R \frac{1}{x^2} dx$

c) (2 points) $\int_1^R \frac{1}{\sqrt{x}} dx$

d) (2 points) $\int_1^R \frac{1}{1+x^2} dx$


e) (2 points) $\int_1^R x dx$

Problem 7) Extrema (10 points)

In Newton's masterpiece "The Method of Fluxions" on the bottom of page 45, Newton asks: "In a given triangle or in a segment of any given curve, to inscribe the greatest rectangle." Lets be more specific and find rectangle with largest area

$$A = xy$$

in the triangle given by the x-axes, y-axes and line $y = 2 - 2x$. Use the second derivative test to make sure you have found the maximum.



Problem 8) Integration by parts (10 points)

a) (5 points) Find

$$\int (1 + x + x^2 + x^3 + x^4)(\sin(x) + e^x) dx .$$

b) (5 points) Find

$$\int \log(x) \frac{1}{x^2} dx .$$

Problem 9) Substitution (10 points)

a) (5 points) **“One,Two,Three,Four Five, once I caught a fish alive!”**

$$\int \frac{(1 + 2x + 3x^2 + 4x^3 + 5x^4)}{(1 + x + x^2 + x^3 + x^4 + x^5)} dx .$$


b) (5 points) A **“Trig Trick-or-Treat”** problem:

$$\int (1 - x^2)^{-3/2} + (1 - x^2)^{-1/2} + (1 - x^2)^{1/2} dx .$$

Problem 10) Partial fractions (10 points)

Integrate

$$\int_{-1}^1 \frac{1}{(x + 3)(x + 2)(x - 2)(x - 3)} dx .$$



The graph of the function is shown to the right.

Lets call it the **friendship graph**.

Problem 11) Chain rule. (10 points)

a) Find the derivative of

$$f(x) = (\sin(7x + x \cos(x)) - 3x) .$$

in general.

b) Now evaluate at $x = 0$.

Problem 12) Various integration problems (10 points)

a) (2 points) $\int_0^{2\pi} 2 \cos^2(x) - \sin(x) dx$

b) (2 points) $\int x^2 e^{3x} dx$

c) (2 points) $\int_1^{\infty} \frac{1}{(x+2)^2} dx$

d) (2 points) $\int \sqrt{x} \log(x) dx$

e) (2 points) $\int_1^e \log(x)^2 dx$

Problem 13) Applications (10 points)

a) (2 points) [**Agnesi density**]

The CDF of the PDF $f(x) = \pi^{-1}/(1+x^2)$ is

b) (2 points) [**Piano man**]

The upper hull of $f(x) = x^2 \sin(1000x)$ is the function

c) (2 points) [**Rower's wisdom**]

If f is power, F is work and $g = F/x$ then $f = g$ if and only if $g'(x) =$

d) (2 points) [**Catastrophes**]

For $f(x) = c(x-1)^2$ there is a catastrophe at $c =$

e) (2 points) [**Randomness**]

We can use chance to compute integrals. It is called the method.