

3/5/2021: First hourly Practice E

”By signing, I affirm my awareness of the standards of the Harvard College Honor Code.”

Your Name:

- Solutions are submitted to knill@math.harvard.edu as PDF handwritten in a file carrying your name. Capitalize the first letters like in OliverKnill.pdf. The paper has to **feature your personal handwriting** and contain no typed part. If you like, you can start writing on a new paper. For 1), you could write 1: False, 2: False ... but you then need to copy the above Honor Code statement and sign.
- No books, calculators, computers, or other electronic aids are allowed. You can use a double sided page of your own handwritten notes when writing the paper. It is your responsibility to submit the paper on time and get within that time also a confirmation. The exam is due at 6 AM on March 6th.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
Total:		100

Problem 1) TF questions (20 points) No justifications are needed.







- 1) T F The function $\arcsin(x)$ is defined as $1/\sin(x)$.
- 2) T F The function $f(x) = \sin(1/x^2)$ can be defined at 0 so that it becomes a continuous everywhere on the real line.
- 3) T F The function $x/\sin(x)$ can be defined at $x = 0$ so that it becomes a continuous function on the real line.
- 4) T F The function $f(x) = \sin^2(x)/x^2$ has the limit 1 at $x = 0$.
- 5) T F The function $f(x) = 1/\log|x|$ has the limit 1 at $x = 0$.
- 6) T F The function $f(x) = (1+h)^{x/h}$ has the property that $Df(x) = [f(x+h) - f(x)]/h = f(x)$.
- 7) T F $\cos(3\pi/2) = 1$.
- 8) T F If a function f is continuous on the interval $[3, 10]$, then it has a global maximum on this interval.
- 9) T F The reciprocal rule assures that $d/dx(1/g(x)) = 1/g(x)^2$.
- 10) T F If $f(0) = g(0) = f'(0) = g'(0) = 0$ and $g''(0) = f''(0) = 1$, then $\lim_{x \rightarrow 0}(f(x)/g(x)) = 1$
- 11) T F An inflection point is a point where the function $f''(x)$ changes sign.
- 12) T F If $f''(x) > 0$ then f is concave up at x .
- 13) T F The chain rule assures that $d/dx f(g(x)) = f'(x)g'(x)$.
- 14) T F The function $f(x) = 1/x + \log(x)$ is continuous on the interval $[1, 2]$.
- 15) T F The function $(e^x - 1)/\cos(x)$ defines an indefinite form at ∞ .
- 16) T F The graph of the function $f(x) = x/(1+x^2)$ has slope 1 at 0.
- 17) T F There is a differentiable function for which $f'(0) = 0$ but for which 0 is not a local extremum.
- 18) T F The second derivative test assures that $x = p$ is a local minimum if $f'(p) = 0$ and $f''(p) < 0$.
- 19) T F The identity $(x^7 - 1)/(x - 1) = x^6 + x^5 + x^4 + x^3 + x^2 + x + 1$ holds for all $x \neq 1$.
- 20) T F The slope of the tangent at a point $(x, f(x))$ of the graph of a differentiable function f is equal to $1/f'(x)$.


Problem 2) Matching problem (10 points) No justifications are needed.

Match the functions with the graphs. Naturally, only 10 of the 12 graphs will appear.

Function	Enter 1-12
$\cot(x)$	
$\cos(2x)$	
$2x$	
$\tan(x)$	
$\log(1/ x)$	

Function	Enter 1-12
x^2	
$\exp(x)$	
$-\sin(x)$	
x^3	
$\text{sinc}(x)$	






Problem 3) Matching problem (10 points) No justifications are needed.

Match the graph of the functions f in $a) - h)$ with the second derivatives f'' in 1)-8).

Function	Second derivative (Enter 1- 8 here)
a)	
b)	
c)	
d)	
e)	
f)	
g)	
h)	



Problem 4) Continuity (10 points)

Some of the following functions might a priori not be defined yet at the point a . In each case, decide whether f can be made a continuous function by assigning a value $f(a)$ at the point a . If no such value exist, state that the function is not continuous.

a) (2 points) $f(x) = \frac{(x^3-1)}{(x-1)}$, at $x = 1$

b) (2 points) $f(x) = \sin(\frac{1}{x}) + \cos(x)$, at $x = 0$

c) (2 points) $f(x) = \sin(\frac{1}{\log(|x|)})$, at $x = 0$

d) (2 points) $f(x) = \log(|\sin(x)|)$, at $x = 0$

e) (2 points) $f(x) = \frac{(x-1)}{x}$, at $x = 0$

Problem 5) Chain rule (10 points)

a) (2 points) Write $1 + \cot^2(x)$ as an expression which only involves the function $\sin(x)$.

b) (3 points) Find the derivative of the function $\operatorname{arccot}(x)$ by using the chain rule for

$$\cot(\operatorname{arccot}(x)) = x .$$

c) (2 points) Write $1 + \tan^2(x)$ as an expression which only involves the function $\cos(x)$.

d) (3 points) Find the derivative of the function $\operatorname{arctan}(x)$ by using the chain rule for

$$\tan(\operatorname{arctan}(x)) = x .$$

Remark: even if you should know the derivatives of arccot or arctan , we want to see the derivations in b) and d).

Problem 6) Derivatives (10 points)

Find the derivatives of the following functions:

a) (2 points) $f(x) = \frac{\cos(3x)}{\cos(x)}$

b) (2 points) $f(x) = \sin^2(x) \log(1 + x^2)$

c) (2 points) $f(x) = 5x^4 - \frac{1}{x^2+1}$

d) (2 points) $f(x) = \tan(x) + \exp(-\sin(x^2))$

e) (2 points) $f(x) = \frac{x^3}{(1+x^2)}$

Problem 7) Limits (10 points)

Find the limits $\lim_{x \rightarrow 0} f(x)$ for the following functions f at $x = 0$ or state (providing reasoning as usual) that the limit does not exist.

a) (2 points) $f(x) = \frac{\sin(3x)}{\sin(x)}$

b) (2 points) $f(x) = \frac{\sin^2(x)}{x^2}$

c) (2 points) $f(x) = \sin(\log(|x|))$

d) (2 points) $f(x) = \tan(x) \log(x)$

e) (2 points) $f(x) = \frac{(5x^4 - 1)}{(x^2 + 1)}$

Problem 8) Extrema (10 points)


A rectangular shoe-box of width x , length x and height y is of volume 2 so that $x^2y = 2$. The surface area adds up three rectangular parts of size $(x \times y)$ and 2 square parts of size $(x \times x)$ and leads to

$$f = 2x^2 + 3xy .$$

a) (2 points) Write down the function $f(x)$ of the single variable x you want to minimize.

b) (6 points) Find the value of x for which the surface area is minimal.

c) (2 points) Check with the second derivative test, whether the point you found is a local minimum.



Problem 9) Global extrema (10 points)

In this problem we study the function $f(x) = 3x^5 - 5x^3$ on the interval $[-2, 2]$.

a) (2 points) Find all roots of f .

b) (2 points) Find all local extrema of the function.

c) (2 points) Use the second derivative test to analyze the critical points, where applicable.

d) (2 points) Find the **global** maximum and minimum of f on the interval $[-2, 2]$.

e) (2 points) Bring the function $x \log(x)$ into indefinite form at 0.