

# INTRODUCTION TO CALCULUS

MATH 1A

## Unit 12: Maxima and Minima

**12.1.** Pierre Fermat made a simple but profound observation: if  $f'(x)$  is not zero, then  $x$  can not be a maximum nor be a minimum. His reasoning was: if you make a step  $h$  then you end up at  $f(x+h) \sim L(x+h) = f(x) + hf'(x)$ . Indeed, we all know that if there is a slope and do a step we end up a bit higher.

**12.2.** Lets call a point  $x$  a **local maximum** of  $f$  if  $f(y) \leq f(x)$  for all  $y$  near enough to  $x$ . The function  $f(x) = x^3 - 2x$  for example has a local minimum at  $x = 1$  and a local maximum at  $x = -1$ . The observation of Fermat is equivalent to:

**Fermat's principle:** If a differentiable function  $f$  has a local maximum or minimum at  $x$ , then  $f'(x) = 0$ .

**12.3.** The function  $f(x) = x^2$  for example has the derivative  $f'(x) = 2x$ . This is zero at  $x = 0$ , the minimum of  $f$ . Note that the converse of Fermat's statement is not necessarily true: if  $f'(x) = 0$ , then  $x$  does not need to be a maximum or minimum. The standard example is  $f(x) = x^3$ . We have  $f'(x) = 3x^2$  which is zero at  $x = 0$ . But  $x = 0$  is neither a maximum nor minimum of  $f$ .

**12.4.** A point  $x$  is called a **critical point** of  $f$ , if  $f'(x) = 0$ . Critical points are important because they are **candidates for maxima and minima**.

**12.5.** The next test allows to see whether we have a maximum or minimum. The derivative should exists near  $a$  but not necessarily at  $a$ . Like for  $f(x) = |x|$  and  $a = 0$ .

**First derivative test:** If  $a$  is a critical point of  $f$  and the slope  $f'(x)$  changes from negative to positive at  $a$  then  $a$  is a local minimum. If  $f'(x)$  changes from positive to negative at  $a$ , then  $a$  is a local maximum. If  $f'(x)$  does not change sign, the  $a$  is neither a local maximum nor local minimum.

**12.6.** Second derivatives help. It assumes that the second derivative exists at  $a$ .

**Second derivative test:** If  $a$  is a critical point of  $f$  and  $f''(a) > 0$ , then  $f$  is a local minimum. If  $f''(a) < 0$ , then  $f$  is a local maximum. If  $f''(a) = 0$ , the test is inconclusive.

# Homework

This PSet is due Wednesday February 21, 2024.

**Problem 12.1:** Find the critical points of the following two functions

a)  $f(x) = x^4 - 4x^3 + 4x^2$ .

b)  $f(x) = x^4(x - 3)^2$ .

**Problem 12.2:** Use the second derivative test to determine the nature of the critical points in the same two functions:

a)  $f(x) = x^4 - 4x^3 + 4x^2$ .

b)  $f(x) = x^4(x - 3)^2$ .

**Problem 12.3:** What does the first derivative test tell you about the behavior at the critical points in the two cases

a)  $f(x) = x^4 - 4x^3 + 4x^2$ .

b)  $f(x) = x^4(x - 3)^2$ .

In each case, is there a point, where the first derivative test gives more information than the second derivative test?

**Problem 12.4:** Find all the critical points and determine whether it is a local maximum, a local minimum or neither. You can use either test. You will see that some of the cases are a bit unusual.

a)  $f(x) = e^2x - e^x$

b)  $f(x) = e^x + x$

c)  $f(t) = t^4 + t^3$

d)  $f(t) = |2t - 8|$

e)  $f(x) = 5$ .

**Problem 12.5:** a) Both the first and second derivative test do not work for the **tamed devil function**  $f(x) = x \sin(1/x)$  at  $x = 0$ . Why not? (Since we have no chain rule yet, you can certainly look up the first and second derivative using a tool like Wolfram alpha.)

b) Function  $f(x) = \arcsin(\sin(x))$  has appeared in our ground hog movie. Where are the maxima and minima? To do so, plot the function  $f(x)$  and its derivative  $f'(x)$  and use one of the derivative tests at the maxima and minima.