

Name:

MWF 9 Oliver Knill
MWF 9 Chao Li
MWF 10 Gijs Heuts
MWF 10 Adrian Zahariuc
MWF 10 Yihang Zhu
MWF 11 Peter Garfield
MWF 11 Matthew Woolf
MWF 12 Charmaine Sia
MWF 12 Steve Wang
MWF 14 Mike Hopkins
TTH 10 Oliver Knill
TTH 10 Francesco Cavazzani
TTH 11:30 Kate Penner
TTH 11:30 Francesco Cavazzani

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see **details** of your computation.
- All functions can be differentiated arbitrarily often unless otherwise specified.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) True/False questions (20 points), no justifications needed

- | | | |
|-----|---|--|
| 1) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | Every function $f(x, y)$ of two variables has either a global minimum or a global maximum. |
| 2) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The linearization of the function $f(x, y) = e^{x+3y}$ at $(0, 0)$ is $L(x, y) = 1 + x + 3y$. |
| 3) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The function $f(x, y, z) = x^2 \cos(z) + x^3 y^2 z + (y - 2)^3 y^5$ satisfies the partial differential equation $f_{xyxzxy} = 12$. |
| 4) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | If $xe^z = y^2 z$, then $\partial z / \partial x = e^z / (y^2 - xe^z)$. |
| 5) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The function $\cos(x^2) \cos(y^2)$ has a local maximum at $(0, 0)$. |
| 6) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The value of the double integral $\int_0^{\pi/4} \int_0^2 x^3 \cos(y) \, dx dy$ is the same as $(\int_0^2 x^3 \, dx)(\int_0^{\pi/4} \cos(y) \, dy)$. |
| 7) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The gradient of $f(x, y)$ is always tangent to the level curves of f . |
| 8) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | If $f(x, y, z) = x - 2y + z$, then the largest possible directional derivative $D_{\vec{u}} f$ at any point in space is $\sqrt{6}$. |
| 9) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | $\int_0^1 \int_0^1 (x^2 + y^2) \, dx dy = \int_0^1 \int_0^1 r^3 \, dr d\theta$. |
| 10) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | It is possible that the directional derivative $D_{\vec{v}} f$ is positive for all unit vectors $\vec{v}$. |
| 11) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | Using linearization of $f(x, y) = xy$ we can estimate $f(0.999, 1.01) \sim 1 - 0.001 + 0.01 = 1.009$. |
| 12) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | Given a curve $\vec{r}(t)$ on a surface $g(x, y, z) = -1$, then $\frac{d}{dt} g(\vec{r}(t)) < 0$. |
| 13) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | If $f(x, y)$ has a local minimum at $(0, 0)$ then it is possible that $f_{xy}(0, 0) > 0$. |
| 14) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The function $f(x, y) = -x^8 - 2x^6 - y^8$ has a local minimum at $(0, 0)$. |
| 15) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | If $\vec{r}(t)$ is a curve in space and f is a function of three variables, then $\frac{d}{dt} f(\vec{r}(t)) = 0$ for $t = 0$ implies that $\vec{r}(0)$ is a critical point of $f(x, y, z)$. |
| 16) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | Let a, b, c be the number of saddle points, maxima and minima of a function $f(x, y)$. Then $a \leq b + c$. |
| 17) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | If $f(x, y)$ is a nonzero function of two variables and R is a region, then $\int \int_R f(x, y) \, dx dy$ is the volume under the graph of f and therefore a positive value. |
| 18) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | We extremize $f(x, y)$ under the constraint $g(x, y) = c$ and obtain a solution (x_0, y_0) . If the Lagrange multiplier λ is positive, then the solution is a minimum. |
| 19) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | The tangent plane to a surface $f(x, y, z) = 1$ intersects the surface in exactly one point. |
| 20) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">F</div> | Let $\vec{v}$ be a vector of length 1 in space. Given a function $f(x, y, z)$ of three variables. If (x_0, y_0, z_0) is a critical point of f , then it is a critical point of $g(x, y, z) = D_{\vec{v}} f(x, y, z)$. |

Problem 2) (10 points)

a) (6 points) Match the regions with the corresponding double integrals



Enter a,b,c,d,e or f	Integral of $f(x,y)$	Enter a,b,c,d,e or f	Integral of $f(x,y)$
	$\int_0^1 \int_{x^2}^{\sqrt{x}} f(x,y) \, dydx$		$\int_0^1 \int_0^{\sqrt{1-x^2}} f(x,y) \, dydx$
	$\int_0^1 \int_0^{\sqrt{y}} f(x,y) \, dxdy$		$\int_0^1 \int_{(1-x)^2}^1 f(x,y) \, dydx$
	$\int_0^1 \int_{y^2}^1 f(x,y) \, dxdy$		$\int_0^1 \int_{(1-x)^2}^{\sqrt{1-x^2}} f(x,y) \, dydx$

b) (4 points) Match the PDE's with the names. No justifications are needed.

Enter A,B,C,D here	PDE
	$f_{xx} = -f_{yy}$
	$f_x = f_y$

Enter A,B,C,D here	PDE
	$f_{xx} = f_{yy}$
	$f_x = f_{yy}$


A) Wave equation	B) Heat equation	C) Transport equation	D) Laplace equation
------------------	------------------	-----------------------	---------------------

Problem 3) (10 points)

- a) (3 points) Find and classify all the critical points of $f(x, y) = xy - x$ on the plane.
- b) (2 points) Decide whether an absolute maximum or an absolute minimum of f exists on the plane $\mathbb{R}^2$.
- c) (3 points) Use the method of Lagrange multipliers to find the maximum and minimum of f on the boundary $x^2 + 4y^2 = 12$ of the elliptical region $G : x^2 + 4y^2 \leq 12$.
- d) (2 points) Find the absolute maximum and absolute minimum of f on the region G given in c).

Problem 4) (10 points)

Find the cylindrical basket which is open on the top has the largest volume for fixed area π . If x is the radius and y is the height, we have to extremize $f(x, y) = \pi x^2 y$ under the constraint $g(x, y) = 2\pi xy + \pi x^2 = \pi$. Use the method of Lagrange multipliers.




Problem 5) (10 points)

The Pac-Man region R is bounded by the lines $y = x, y = -x$ and the unit circle. The number

$$a = \frac{\int \int_R x \, dx dy}{\int \int_R 1 \, dx dy}$$

defines the point $C = (a, 0)$ called center of mass of the region. Find it.




Problem 6) (10 points)

- a) (5 points) Find the tangent plane to the surface $\sqrt{xyz} = 60$ at $(x, y, z) = (100, 36, 1)$.
- b) (5 points) Estimate $\sqrt{100.1 * 36.1 * 0.999}$ using linear approximation. Here, for clarity reasons, we use $*$ for the usual multiplication for numbers.

Problem 7) (10 points)

Oliver got a diammagnetic kit, where strong magnets produce a force field in which pyrolytic graphitic floats. The gravitational field produces a well of the form $f(x, y) = x^4 + y^3 - 2x^2 - 3y$. Find all critical points of this function and classify them. Is there a global minimum?



Right picture credit: Wikipedia.

Problem 8) (10 points)

Let $f(x, y) = xy$.

- a) (2 points) Find the direction of maximal increase at the point $(1, 1)$.
- b) (3 points) Find the directional derivative at $(1, 1)$ in the direction $\langle 3/5, 4/5 \rangle$.
- c) (2 points) The curve $\vec{r}(t) = \langle \sqrt{2} \sin(t), \sqrt{2} \cos(t) \rangle$ passes through the point $(1, 1)$ at some time t_0 . Find $\frac{d}{dt}f(\vec{r}(t))$ at time t_0 directly.
- d) (3 points) Find $\frac{d}{dt}f(\vec{r}(t))$ at time t_0 using the multivariable chain rule.

Problem 9) (10 points)

Integrate the function

$$f(x, y) = \frac{y^5 - 1}{y^{1/3} - y^{1/4}}$$

on the finite region bounded by the curves $y = x^3$ and $y = x^4$.

Problem 10) (10 points)

The main building of a mill has a cone shaped roof and cylindrical walls. If the cylinder has radius r , the height of the side wall is h and the height of the roof is h , then the volume is

$$V(h, r) = \pi r^2 h + h \pi r^2 / 3 = (4\pi/3) h r^2$$

and assume the cost of the building is

$$A(h, r) = \pi r^2 + 2\pi r h + \pi 2r^2 = \pi(3r^2 + 2rh)$$

which is the area of the ground plus the area of the wall plus $2\pi r h$, the cost for the roof. For fixed volume $V(h, r) = 4\pi/3$, minimize the cost $A(h, r)$ using the Lagrange multiplier method.

