

Name:

--

MWF 9 Oliver Knill
MWF 9 Chao Li
MWF 10 Gijs Heuts
MWF 10 Yu-Wen Hsu
MWF 10 Yong-Suk Moon
MWF 11 Rosalie Belanger-Rioux
MWF 11 Gijs Heuts
MWF 11 Siu-Cheong Lau
MWF 12 Erick Knight
MWF 12 Kate Penner
TTH 10 Peter Smillie
TTH 10 Jeff Kuan
TTH 10 Yi Xie
TTH 11:30 Jeff Kuan
TTH 11:30 Jameel Al-Aidroos

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see details of your computation.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 180 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
14		10
Total:		150

Problem 1) True/False questions (20 points)

- 1) T F The projection vector $\text{proj}_{\vec{v}}(\vec{w})$ is parallel to \vec{w} .

Solution:

It is parallel to \vec{v} .

- 2) T F Any parametrized surface S is a graph of a function $f(x, y)$.

Solution:

A counter example is the sphere.

- 3) T F If the directional derivatives $D_{\vec{v}}(f)(1, 1)$ and $D_{\vec{w}}(f)(1, 1)$ are both 0 for $\vec{v} = \langle 1, 1 \rangle / \sqrt{2}$ and $\vec{w} = \langle 1, -1 \rangle / \sqrt{2}$, then $(1, 1)$ is a critical point.

Solution:

Indeed $\nabla f(1, 1)$ must be perpendicular to \vec{v} and \vec{w} and so be the zero vector.

- 4) T F The linearization $L(x, y)$ of $f(x, y) = x + y + 4$ at $(0, 0)$ satisfies $L(x, y) = f(x, y)$.

Solution:

The linearization of any linear function at $(0, 0)$ is the function itself.

- 5) T F For any function $f(x, y)$ of two variables, the line integral of the vector field $\vec{F} = \nabla f$ on a level curve $\{f = c\}$ is always zero.

Solution:

The gradient is perpendicular to the velocity vector.

- 6) T F If \vec{F} is a vector field of unit vectors defined in $1/2 \leq x^2 + y^2 \leq 2$ and \vec{F} is tangent to the unit circle C , then $\int_C \vec{F} \cdot d\vec{r}$ is either equal to 2π or -2π .

Solution:

\vec{r}' is parallel to \vec{F} so that $\vec{F} \cdot \vec{r}'$ is equal to 1 or -1 .

- 7) T F If a curve C intersects a surface S at a right angle, then at the point of intersection, the tangent vector to the curve is parallel to the normal vector of the surface.

Solution:

This is clear once you know what the question means.

- 8) T F The curvature of the curve $\vec{r}(t) = \langle \cos(3t), \sin(6t) \rangle$ at the point $\vec{r}(0)$ is smaller than the curvature of the curve $\vec{r}(t) = \langle \cos(30t), \sin(60t) \rangle$ at the point $\vec{r}(0)$.

Solution:

The curvature is independent of the parametrization of the curve.

- 9) T F At every point (x, y, z) on the hyperboloid $x^2 + 2y^2 - z^2 = 1$, the vector $\langle x, 2y, -z \rangle$ is normal to the hyperboloid.

Solution:

Look at the gradient. It is parallel to the vector.

- 10) T F The set $\{\phi = \pi/2, \theta = \pi\}$ in spherical coordinates is the negative x axis.

Solution:

$\phi = \pi/2$ forces us to be on the xy -plane. $\theta = \pi$ is the negative x axis

- 11) T F The integral $\int_0^1 \int_0^{2\pi} \int_0^\pi \rho^2 \sin^2(\phi) \, d\phi \, d\theta \, d\rho$ is equal to the volume of the unit ball.

Solution:

If $\sin^2(\phi)$ would be $\sin(\phi)$, then it would be the integral in spherical coordinates.

- 12) T F Four points A, B, C, D are located in a single common plane if $(B - A) \cdot ((C - A) \times (D - A)) = 0$.

Solution:

This is the volume of the parallelepiped with corners A, B, C, D . If the volume is zero, then the parallelepiped is flat and the points in a plane.

- 13) T F If a function $f(x, y)$ has a local maximum at $(0, 0)$, then the discriminant D is negative.

Solution:

False, we also can have $D = 0$ like for $f(x, y) = 1 - x^4 - y^4$.

- 14) T F The integral $\int_0^x \int_y^1 f(x, y) dx dy$ represents a double integral over a bounded region in the plane.

Solution:

The integral is not properly defined. There can be no variable in the most outer integral.

- 15) T F The following identity is true: $\int_0^3 \int_0^{2x} x^2 dy dx = \int_0^6 \int_{y/2}^3 x^2 dx dy$

Solution:

Make a picture and draw the triangle.

- 16) T F The integral $\iint_S \text{curl}(\vec{F}) \cdot d\vec{S}$ over the surface S of a cube is zero for all vector fields \vec{F} .

Solution:

By the divergence theorem, this flux integral is equal to the triple integral of $\text{div}(\vec{F})$ over the cube. But since $\text{div}(\text{curl}(\vec{F})) = 0$, this integral is zero.

- 17) T F A vector field \vec{F} defined on three space which is incompressible ($\text{div}(\vec{F}) = 0$) and irrotational ($\text{curl}(\vec{F}) = 0$) can be written as $\vec{F} = \nabla f$ with $\Delta f = \nabla^2 f = 0$.

Solution:

Every gradient field $F = \nabla f$, for which $\Delta(f) = 0$, is also incompressible.

- 18) T F If a vector field \vec{F} is defined at all points of three-space except the origin, and $\text{curl}(\vec{F}) = \vec{0}$ everywhere, then the line integral of \vec{F} around the circle $x^2 + y^2 = 1$ in the xy -plane is equal to zero.

Solution:

The circle is the boundary of a hemisphere which is contained in the region, where \vec{F} is defined.

- 19) T F The identity $\text{curl}(\text{grad}(\text{div}(\vec{F}))) = \vec{0}$ is true for all vector fields $\vec{F}(x, y, z)$.

Solution:

Already $\text{curl}(\text{grad}(f)) = \vec{0}$.


- 20) T F If $\vec{F} = \text{curl}(\vec{G})$, where $\vec{G} = \langle e^{e^x}, 5^x z^5, \sin y \rangle$, then $\text{div}(\vec{F}(x, y, z)) > 0$ for all (x, y, z) .

Solution:


$\text{div}(\text{curl}(\vec{F})) = 0$.

Problem 2) (10 points)


Match the equations with the space curves. No justifications are needed.




I



II



III



IV


Enter I,II,III,IV here	Equation
	$\vec{r}(t) = \langle t^2, t^3 - t, t \rangle$
	$\vec{r}(t) = \langle 1 - t , t - t - 1 , t \rangle$
	$\vec{r}(t) = \langle 2 \sin(5t), \cos(11t), t \rangle$
	$\vec{r}(t) = \langle t \sin(1/t), t \cos(1/t) , t \rangle$

Solution:


Enter I,II,III,IV here	Equation
II	$\vec{r}(t) = \langle t^2, t^3 - t, t \rangle$
III	$\vec{r}(t) = \langle 1 - t , t - t - 1 , t \rangle$
I	$\vec{r}(t) = \langle 2 \sin(5t), \cos(11t), t \rangle$
IV	$\vec{r}(t) = \langle t \sin(1/t), t \cos(1/t) , t \rangle$

Problem 3) (10 points)


Match the equations with the objects. No justifications are needed.




I




II




III



IV




V



VI



VII



VIII

Enter I,II,III,IV,V,VI,VII,VIII here	Equation
	$\vec{r}(s, t) = \langle (2 + \cos(s)) \cos(t), (2 + \cos(s)) \sin(t), \sin(s) \rangle$
	$\vec{r}(t) = \langle \cos(3t), \sin(5t) \rangle$
	$x^2 + y^2 - z^2 = 1$
	$\vec{F}(x, y, z) = \langle -y, x, 1 \rangle$
	$x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq z^2, z \geq 0$
	$z = f(x, y) = x^2 - y$
	$g(x, y) = x^2 - y^2 = 1$
	$\vec{F}(x, y) = \langle -y, x \rangle$

Solution:

Enter I,II,III,IV,V,VI,VII,VIII here	Equation
II	$\vec{r}(s, t) = \langle (2 + \cos(s)) \cos(t), (2 + \cos(s)) \sin(t), \sin(s) \rangle$
VIII	$\vec{r}(t) = \langle \cos(3t), \sin(5t) \rangle$
V	$x^2 + y^2 - z^2 = 1$
I	$\vec{F}(x, y, z) = \langle -y, x, 1 \rangle$
VI	$x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq z^2, z \geq 0$
IV	$z = f(x, y) = x^2 - y$
VII	$g(x, y) = x^2 - y^2 = 1$
III	$\vec{F}(x, y) = \langle -y, x \rangle$

Problem 4) (10 points)

a) Find an equation for the plane Σ passing through the points $\vec{r}(0), \vec{r}(1), \vec{r}(2)$, where $\vec{r}(t) = \langle t^2, t^4, t \rangle$.

b) Find the distance between the point $\vec{r}(-1)$ and the plane Σ found in a).

Solution:

a) $r(0) = (0, 0, 0), r(1) = (1, 1, 1), r(2) = (4, 16, 2)$. The normal vector is $n = \langle a, b, c \rangle = \langle 1, 1, 1 \rangle \times \langle 4, 16, 2 \rangle = \langle -14, 2, 12 \rangle$. The plane has the equation $ax + by + cz = d$, where d is obtained from plugging in one of the points. The answer is $\boxed{-14x + 2y + 12z = 0}$.

b) $Q = (0, 0, 0)$ is a point on the plane. The point $P = \vec{r}(-1) = (1, 1, -1)$ has distance $|\vec{PQ} \cdot \vec{n}|/|\vec{n}| = |\langle 1, 1, -1 \rangle \cdot \langle -14, 2, 12 \rangle|/\sqrt{196 + 4 + 144} = \boxed{24/\sqrt{344}}$ from the plane.

Problem 5) (10 points)

A vector field $\vec{F}(x, y)$ in the plane is given by $\vec{F}(x, y) = \langle x^2 + 5, y^2 - 1 \rangle$. Find all the critical points of $|\vec{F}(x, y)|$ and classify them. At which point or points is $|\vec{F}(x, y)|$ minimal?

Solution:

Instead of extremizing the length, we can instead extremize the square of the length. This has the advantage of not involving square roots.

Extremize $f(x, y) = x^4 + 10x^2 + 25 + y^4 - 2y^2 + 1$. The gradient is $\nabla f(x, y) = \langle 4x^3 + 20x, 4y^3 - 4y \rangle = \langle 4x(x^2 + 5), 4y(y^2 - 1) \rangle$. We have $\nabla f(x, y) = \langle 0, 0 \rangle$ for the critical points $(0, -1)$, $(0, 0)$, and $(0, 1)$. We have $f_{xx} = 20$ at all three points and $D(0, 0) = -80, D(0, \pm 1) = 160$. The point $(0, 0)$ is a saddle point and the two critical points $(0, 1), (0, -1)$ are the global minima.

Problem 6) (10 points)

A house is situated at the point $(0, 0)$ in the middle of a mountainous region. The altitude at each point (x, y) is given by the equation $f(x, y) = 4x^2y + y^3$. There is a pathway in the shape of an ellipse around the house, on which the (x, y) coordinates satisfy $2x^2 + y^2 = 6$. Find the highest and lowest points in the closed region bounded by the path.

Solution:

The gradient of f is $\nabla f = \langle 8xy, 4x^2 + 3y^2 \rangle$. From the second equation, the only critical point for the function without constraints is $x = y = 0$. At this critical point, the function f has value 0. If $g(x, y) = 2x^2 + y^2$, then $\nabla g = \langle 4x, 2y \rangle$.

$(0, 0)$ is a critical point with $f(0, 0) = 0$.


For local maxima and minima on the ellipse, the equation $\nabla f = \lambda \nabla g$ must hold, so

$$\langle 8xy, 4x^2 + 3y^2 \rangle = \langle 4x\lambda, 2y\lambda \rangle.$$

Equating the first coordinates gives $8xy = 4x\lambda$, so $x = 0$ or $\lambda = 2y$. In the first case, $y^2 = 6$, so $y = \pm\sqrt{6}$. Therefore the function f has value $\pm 6\sqrt{6}$ in this case. Otherwise, $\lambda = 2y$, and so equating the second coordinates, $2y\lambda = 2y \cdot 2y = 4y^2 = 4x^2 + 3y^2$. Hence $y^2 = 4x^2$. At this point, one can work out $y = \pm 2x$. In any event, $2x^2 + y^2 = 6x^2$, so $6x^2 = 6$. Therefore $x = \pm 1$ and $y = \pm 2$.

In conclusion, there are 6 constraint extrema on the boundary $(0, \sqrt{6}), (0, -\sqrt{6}), (1, 2), (1, -2), (-1, 2), (-1, -2)$

The maximal value of f is +16 and obtained at the points $(1, 2)$ and $(-1, 2)$. The minimal value of f is obtained at the points $(1, -2)$ and $(-1, -2)$.



Problem 7) (10 points)

We are given a function $f(x, y)$ with $x = r \cos(\theta)$ and $y = r \sin(\theta)$ as well as the following data points. Evaluate $\frac{\partial^2 f}{\partial \theta^2}$ at the point $r = 2, \theta = \frac{\pi}{2}$.

(x, y)	$(0, 2)$	$(2, 0)$	$(\pi, 2)$	$(2, \pi)$	$(0, 0)$
$f(x, y)$	2004	2005	2002	2003	2006
$f_x(x, y)$	3	4	6	2	0
$f_y(x, y)$	2	3	4	5	0
$f_{xx}(x, y)$	6	5	4	0	0
$f_{xy}(x, y)$	0	1	0	2	0
$f_{yy}(x, y)$	2	0	2	2	0

Solution:

The first derivative is

$$\begin{aligned}\frac{\partial f}{\partial \theta} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} \\ &= -r \sin \theta f_x + r \cos \theta f_y\end{aligned}$$

So when you take the 2nd derivative you have to use the product rule and the chain rule. The product rule gives you the derivative:

$$\frac{\partial^2 f}{\partial \theta^2} = -r \cos \theta f_x - r \sin \theta \frac{\partial}{\partial \theta}(f_x) - r \sin \theta f_y + r \cos \theta \frac{\partial}{\partial \theta}(f_y)$$

The chain rule is used to find:

$$\frac{\partial}{\partial \theta}(f_x) = \frac{\partial f_x}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f_x}{\partial y} \frac{\partial y}{\partial \theta} = f_{xx} * (-r \sin \theta) + f_{xy} * (r \cos \theta)$$

$$\frac{\partial}{\partial \theta}(f_y) = \frac{\partial f_y}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f_y}{\partial y} \frac{\partial y}{\partial \theta} = f_{yx} * (-r \sin \theta) + f_{yy} * (r \cos \theta)$$

So then plugging these in you get:

$$\begin{aligned}\frac{\partial^2 f}{\partial \theta^2} &= -r \cos \theta f_x - r \sin \theta (f_{xx} * (-r \sin \theta) + f_{xy} * (r \cos \theta)) - r \sin \theta f_y + r \cos \theta (f_{yx} * (-r \sin \theta) + f_{yy} * (r \cos \theta)) \\ &= -r \cos \theta f_x + r^2 \sin^2 \theta f_{xx} - r^2 \sin \theta \cos \theta f_{xy} - r \sin \theta f_y - r^2 \sin \theta \cos \theta f_{yx} + r^2 \cos^2 \theta f_{yy} \\ &= -r \cos \theta f_x - r \sin \theta f_y + r^2 \sin^2 \theta f_{xx} - 2r^2 \sin \theta \cos \theta f_{yx} + r^2 \cos^2 \theta f_{yy}\end{aligned}$$

The point $r = 2, \theta = \pi/2$ is in Cartesian coordinates the $(x, y) = (0, 2)$, so that we have to look at the first column of the table to look up the derivatives of f . Because $\cos(\pi/2) = 0$, two terms of the above formula vanish and the answer is

$$\frac{\partial^2 f}{\partial \theta^2} = 4f_{xx}(0, 2) - 2f_y(0, 2) = 24 - 4 = 20.$$

Problem 8) (10 points)

a) (4 points) Where does the tangent plane at $(1, 1, 1)$ to the surface $z = e^{x-y}$ intersect the z axis?

b) (4 points) Estimate $f(x, y, z) = 1 + \log(1 + x + 2y + z) + 2\sqrt{1+z}$ at the point $(0.02, -0.001, 0.01)$.

c) (2 points) $f(x, y, z) = 0$ defines z as a function $g(x, y)$ of x and y . Find the partial derivative $g_x(x, y)$ at the point $(x, y) = (0, 0)$.

Solution:

a) The tangent plane has the equation $ax + by + cz = d$, where $\langle a, b, c \rangle = \nabla g(1, 1, 1)$, where $g(x, y, z) = z - e^{x-y}$. Because $\nabla g(x, y, z) = \langle -e^{x-y}, e^{x-y}, 1 \rangle$, we have $\nabla g(1, 1, 1) = \langle -1, 1, 1 \rangle$. The plane has the equation $-x + y + z = d$. The number d can be obtained by plugging in the point $(1, 1, 1)$. Therefore the plane is $-x + y + z = 1$. This plane intersects the z axes at $\boxed{z = 1}$.

b) The linearization of $1 + \log(1 + x + 2y + z) + 2\sqrt{1 + z}$ at $(0, 0, 0)$ is $L(x, y, z) = 3 + x + 2y + 2z$. Now $L(0.02, -0.001, 0.01) = 3 + 0.02 - 2 \cdot 0.001 + 2 \cdot 0.01 \boxed{= 3.038}$.

c) The implicit computation formula is $g_x(x, y) = -f_x(x, y, z)/f_z(x, y, z)$. We have $f_z = (1/(1+x+2y+z)+1/\sqrt{1+z})$ and $f_x = 1/(1+x+2y+z)$ so that $g_x(0, 0) = -1/(\sqrt{1+z}+1)$. The value of z would have to be computed numerically.

Problem 9) (10 points)

For each of the following quantities, set up a double or triple integral using any coordinate system you like. You do not have to evaluate the integrals, but the bounds of each single integral must be specified explicitly.

1. (3 points) The volume of the tetrahedron with vertices $(0, 0, 0)$, $(3, 0, 0)$, $(0, 3, 0)$ and $(0, 0, 3)$.
2. (4 points) The surface area of the piece of the paraboloid $z = x^2 + y^2$ lying in the region $z = x^2 + y^2$, where $0 \leq z \leq 1$.
3. (3 points) The volume of the solid bounded by the planes $z = -1$, $z = 1$ and the one-sheeted hyperboloid $x^2 + y^2 - z^2 = 1$.

Solution:

(1) The tetrahedron is bounded by the xy, yz, zx -planes and the plane $z = 3 - x - y$. The triple integral would be: $\int_0^3 \int_0^{3-x} \int_0^{3-x-y} dz dy dx$. Evaluating the inner integral $\int_0^3 \int_0^{3-x} (3 - x - y) dy dx$.

(2) The parameterization $\vec{r}(r, \theta) = \langle r \cos(\theta), r \sin(\theta), r^2 \rangle$ gives

$$\begin{aligned} |\vec{r}_r \times \vec{r}_\theta| &= |\langle \cos(\theta), \sin(\theta), 2r \rangle \times \langle -r \sin(\theta), r \cos(\theta), 0 \rangle| \\ &= |\langle -2r^2 \cos(\theta), -2r^2 \sin(\theta), r \rangle| = \sqrt{4r^4 + r^2}. \end{aligned}$$

The surface area integral is $\int_0^{2\pi} \int_0^1 \sqrt{4r^4 + r^2} dr d\theta$ or $\int_0^1 \int_0^{2\pi} \sqrt{4r^4 + r^2} d\theta dr$.

Parameterization $\vec{r}(x, y) = \langle x, y, x^2 + y^2 \rangle$ gives

$$|\vec{r}_x \times \vec{r}_y| = |\langle 1, 0, 2x \rangle \times \langle 0, 1, 2y \rangle| = |\langle -2x, -2y, 1 \rangle| = \sqrt{4x^2 + 4y^2 + 1},$$

and the integral $\int_{-1}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \sqrt{4x^2 + 4y^2 + 1} dx dy$ or $\int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \sqrt{4x^2 + 4y^2 + 1} dy dx$.

(3) Using cylindrical coordinates, the integral is $\int_{-1}^1 \int_0^{\sqrt{1+z^2}} \int_0^{2\pi} r d\theta dr dz$ which is $2\pi \cdot \int_{-1}^1 \int_0^{\sqrt{1+z^2}} r dr dz$. If the order of integration is changed, then - still in cylindrical coordinates - we have


$$2\pi \left(\int_0^1 \int_{-1}^1 r dz dr + 2 \int_1^{\sqrt{2}} \int_{\sqrt{r^2-1}}^1 r dz dr \right).$$

When using spherical coordinates, one would have to split up the integral into two parts and setting up the integral is harder.

Problem 10) (10 points)

A region R in the xy -plane is given in polar coordinates by $r(\theta) \leq \theta$ for $\theta \in [0, \pi]$. You see the region in the picture to the right. Evaluate the double integral

$$\iint_R \frac{\cos(\sqrt{x^2 + y^2})}{\sqrt{x^2 + y^2}(\pi - \sqrt{x^2 + y^2})} dx dy.$$



Solution:

The region becomes a triangle in polar coordinates. Setting up the integral with $dA = dr d\theta$ does not work. The integral $\int_0^\pi \int_0^\theta \frac{\cos(r)}{r(\pi-r)} r dr d\theta$ can not be solved. We have to change the order of integration:

$$\int_0^\pi \int_r^\pi \frac{\cos(r)}{r(\pi-r)} r d\theta dr$$

Evaluating the inner integral gives $\int_0^\pi \cos(r) dr = \boxed{0}$.

Problem 11) (10 points)

A car drives up a freeway ramp C which is parametrized by

$$\vec{r}(t) = \langle \cos(t), 2 \sin(t), t \rangle, \quad 0 \leq t \leq 3\pi.$$

a) (3 points) Set up an integral which gives the length of the ramp. You do not need to evaluate it.

b) (3 points) Find the unit tangent vector \vec{T} to the curve at the point where $t = 0$.

c) (4 points) Suppose the wind pattern in the area is such that the wind exerts a force $\vec{F} = \langle 4x^2, y, 0 \rangle$ on the car at the position (x, y, z) . What is the total work gain as the car drives up the ramp? In other words, what is the line integral $\int_C \vec{F} \cdot d\vec{r}$.

Solution:

a) $\vec{r}'(t) = \langle -\sin(t), 2 \cos(t), 1 \rangle$ and $|\vec{r}'(t)| = \sqrt{\sin^2(t) + 4 \cos^2(t) + 1}$. The integral is

$$\int_0^{3\pi} \sqrt{\sin^2(t) + 4 \cos^2(t) + 1} dt.$$

=


$$\int_0^{3\pi} \sqrt{2 + 3 \cos^2(t)} dt.$$

b) $\vec{T}(t) = \vec{r}'(t)/|\vec{r}'(t)| = \langle -\sin(t), 2 \cos(t), 1 \rangle / \sqrt{\sin^2(t) + 4 \cos^2(t) + 1}$. At $t = 0$, we have $\vec{T}(0) = \langle 0, 2, 1 \rangle / \sqrt{5}$.

c) The line integral is $\int_0^{3\pi} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$. Note that \vec{F} is a gradient field with potential function $f(x, y, z) = 4x^3/3 + y^2/2$ so that the line integral is $f(\vec{r}(3\pi)) - f(\vec{r}(0))$ by the fundamental theorem of line integrals. Now $f(\vec{r}(3\pi)) = f(-1, 0, 3\pi) = -4/3$ and $f(\vec{r}(0)) = f(1, 0, 0) = 4/3$, so that the result is $\boxed{-8/3}$.

Problem 12) (10 points)

Suppose \vec{F} is an irrotational vector field in the plane (that is, its curl is everywhere zero) that is not defined at the origin $O = (0, 0)$. Suppose the line integral of \vec{F} along the path p from A to B is 5 and the line integral of \vec{F} along the path q from A to B is -4 . Find the line integral of \vec{F} along the following three paths:



- (3 points) The path a from A to B going clockwise below the origin.
- (4 points) The closed path b encircling the origin in a clockwise direction.
- (3 points) The closed path c starting at A and ending in A without encircling the origin.

Solution:

- The result is the same for the path a and the path q . The vector field is conservative in the lower half plane. The result is $\boxed{-4}$.
- The line integral is the same as the difference of the line integral along q and the line integral along p which is $-4 - 5 = \boxed{-9}$. The path $q - p$ encircles the origin in the same direction than the path b . Because the curl is 0 in the region enclosed by these two curves, Greens theorem assures that the line integrals are the same.
- The vector field \vec{F} is conservative in the right half plane. By the fundamental theorem of line integrals or using the closed loop property, the result is $\boxed{0}$.

Problem 13) (10 points)

Let S be the surface which bounds the region enclosed by the paraboloid $z = x^2 + y^2 - 1$ and the xy plane. Let \vec{F} be the vector field $\vec{F}(x, y, z) = \langle x + e^{\sin(z)}, z, -y \rangle$.

a) (5 points) Find the flux of \vec{F} through the surface S .

b) (5 points) Find the flux of \vec{F} through the part of the surface S that belongs to the paraboloid, oriented so that the normal vector points downwards.

Solution:

a) The flux out of the whole surface can be determined by the Divergence Theorem. The flux over the whole surface is the integral of the divergence over the solid region it encloses. The divergence of the vector field is 1. So the flux is the volume of the solid region, namely the integral $\int_D 1 - x^2 - y^2 \, dydx$ over the unit disc D . Converting to polar coordinates, this is $\int_0^{2\pi} \int_0^1 (1 - r^2)r \, dr \, d\theta = \int_0^{2\pi} \int_0^1 r - r^3 \, dr \, d\theta = \int_0^{2\pi} \frac{1}{2} - \frac{1}{4} \, d\theta = \int_0^{2\pi} \frac{1}{4} \, d\theta = \frac{\pi}{2}$. So the flux out of the whole surface is $\boxed{\pi/2}$.

b) There the outward unit normal vector to the top surface which is \vec{k} , so the flux is

$$\int_D -y \, dA = \int_0^{2\pi} \int_0^1 -r \sin \theta \, r \, dr \, d\theta = \int_0^{2\pi} -\frac{1}{3} \sin \theta \, d\theta = 0 .$$

Therefore the flux out of the "roof" D is $\boxed{0}$. The flux through the "floor" the paraboloid part is $\pi/2 - 0 = \boxed{\pi/2}$.

Problem 14) (10 points)

Let \vec{F} be the vector field $\vec{F}(x, y, z) = \langle 4z + \cos(\cos x), y^2, x + 2y \rangle$.

a) (4 points) Let C be the curve given by the parameterization $\vec{r}(t) = \langle \cos t, 0, \sin t \rangle$, for $0 \leq t \leq 2\pi$. Find the line integral of \vec{F} along C .

b) (6 points) Let S be the hemisphere of the unit sphere defined by $y \leq 0$. Find the flux of the curl of \vec{F} out of S . In other words, find

$$\iint_S \text{curl}(\vec{F}) \cdot d\vec{S} .$$

For part b), the surface S is oriented so that the normal vector has a positive y -component.

Solution:

a) The curl of the vector field is $\langle 2, 3, 0 \rangle$. The parameterization describes the circle $x^2 + z^2 = 1$, where $y = 0$. The curve starts at $(1, 0, 0)$ and rotates towards back towards $(0, 0, 1)$. By Stokes theorem, the line integral can be computed as the flux of $\text{curl}(F)$ through the unit disk D in the xz plane which has the normal vector $r_u \times r_v = -\vec{j}$ and $\text{curl}(\vec{F})(x, y, z) \cdot (r_u \times r_v) = -3$. The flux is

$$\iint_D -3 \, dx \, dz = -3\pi .$$

b) The boundary of S is the curve C , oriented differently as in part a). Therefore, the answer is 3π by **Stokes theorem**. Note that the sign is different because the orientation of the curve C and the surface S do not match.