

Name:

MWF 9 Jun-Hou Fung
MWF 9 Koji Shimizu
MWF 10 Matt Demers
MWF 10 Dusty Grundmeier
MWF 10 Erick Knight
MWF 11 Oliver Knill
MWF 11 Kate Penner
MWF 12 Yusheng Luo
MWF 12 YongSuk Moon
TTH 10 Will Boney
TTH 10 Peter Smillie
TTH 10 Chenglong Yu
TTH 11:30 Lukas Brantner
TTH 11:30 Yu-Wen Hsu

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3,8, we need to see **details** of your computation.
- All functions can be differentiated arbitrarily often unless otherwise specified.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) True/False questions (20 points), no justifications needed

- 1) T F There is a function $f(x, y)$ for which the linearization at $(0, 0)$ is $L(x, y) = x^2 + y^2$.
- 2) T F For any two functions f, g and unit vector \vec{u} we have $D_{\vec{u}}(f + g) = D_{\vec{u}}f + D_{\vec{u}}g$.
- 3) T F $\int_0^2 \int_0^{\sqrt{4-x^2}} (x^2 + y^2) dydx = \int_0^2 \int_0^{\pi/2} r^2 d\theta dr$.
- 4) T F If we solve $\sin(y) - xy^2 = 0$ for y , then $y' = -y^2/(\cos(y) - 2xy)$.
- 5) T F If $f(x, 0) = 0$ for all x and $f(0, y) = 0$ for all y , then $g(x, y) = \int_0^x \int_0^y f(s, t) dt ds$ solves $g_{xy}(x, y) = f(x, y)$.
- 6) T F If $|\nabla f| = 1$ at $(0, 0)$, then there exists a direction in which the slope of the graph of f at $(0, 0)$ is 1.
- 7) T F The function $f(x, y) = x^2 + y^2$ satisfies the partial differential equation $f_{xx}f_{yy} - f_{xy}^2 = 4$.
- 8) T F The height of Mount Wachusett is $f(x, y) = 4 - 2x^2 - y^2$. On the trail $x^2 + y^2 = 1$, the point $(1, 0)$ is a maximum.
- 9) T F Mount Wachusett has height $f(x, y) = 4 - 2x^2 - y^2$. Except at the maximum $(0, 0)$, the gradient vector is perpendicular to the graph of the function.
- 10) T F If $f_x(a, b) > 0$ and $f_y(a, b) > 0$ then for any unit vector \vec{u} we must have $D_{\vec{u}}f(a, b) > 0$.
- 11) T F If $f(x, y)$ has two local minima, then f must have at least one local maximum.
- 12) T F If $\vec{r}(t)$ is a curve on the surface $g(x, y, z) = x^2 + y^2 - z^2 = 6$ then $\nabla g(\vec{r}(t)) \cdot \vec{r}'(t) = 0$.
- 13) T F If f and g have the same trace $\{x = 5\}$ then $f_x(5, y) = g_x(5, y)$ for all y .
- 14) T F If f and g have the same trace $\{x = 5\}$ then $f_y(5, y) = g_y(5, y)$ for all y .
- 15) T F The surface area of $\vec{r}_1(u, v) = \langle u \cos(v), u \sin(v), u^2 \rangle$ and $\vec{r}_2(u, v) = \langle \sqrt{u} \cos(v), \sqrt{u} \sin(v), u \rangle$ defined on $\{0 \leq u, v \leq 1\}$ are the same.
- 16) T F If $\vec{r}(t)$ is a curve on a graph $z = f(x, y)$ of a function $f(x, y)$, then the velocity vector of \vec{r} is perpendicular to the vector $\langle f_x, f_y, -1 \rangle$.
- 17) T F A continuous function $f(x, y)$ on the closed disc $R = \{x^2 + y^2 \leq 51^2\}$ (of course, R is called “**area** 51π ”) has a global maximum on R .
- 18) T F Any continuous function $f(x, y)$ has a global minimum and maximum on the curve $y = x^2$.
- 19) T F Fubini’s theorem assures that $\int_a^b \int_c^d f(x, y) dydx = \int_a^b \int_c^d f(x, y) dx dy$.
- 20) T F $\iint_R \sin(x + y) dx dy = 0$ for $R = \{-\pi \leq x \leq \pi, -\pi \leq y \leq \pi\}$.

Problem 2) (10 points)

a) (6 points) Match the integration regions with the integrals. Each integral matches exactly one region $A - F$.



Enter A-F	Integral
	$\int_{-1}^1 \int_{-x^2}^{x^2} f(x, y) dy dx.$
	$\int_{-1}^1 \int_{-y^2}^{y^2} f(x, y) dx dy.$
	$\int_{-1}^1 \int_{y^2-1}^{1-y^2} f(x, y) dx dy.$
	$\int_{-1}^1 \int_0^{\sqrt{1-y^2}} f(x, y) dx dy.$
	$\int_{-1}^1 \int_{x^2-1}^{1-x^2} f(x, y) dy dx.$
	$\int_{-1}^1 \int_{-\sqrt{1-x^2}}^0 f(x, y) dy dx.$


b) (4 points) Fill in one word names (like “Heat”, “Wave” etc) for the partial differential equations:

Enter one word	PDE
	$g_x = g_y$
	$g_{xx} = g_{yy}$
	$g_{xx} = -g_{yy}$
	$g_x = g_{yy}$

Problem 3) (10 points)


(10 points) A function $f(x, y)$ of two variables has level curves as shown in the picture. We also see a constraint in the form of a curve $g(x, y) = 0$ which has the shape of the graph of the cos function. The arrows show the gradient. In this problem, each of the 10 letters $A, B, C, D, E, F, G, H, K, M$ appears exactly once.

Enter A-P	Description
	a local maximum of $f(x, y)$.
	a local minimum of $f(x, y)$.
	a saddle point of $f(x, y)$ where $f_{xx} < 0$.
	a saddle point of $f(x, y)$ where $f_{xx} > 0$.
	a saddle point of $f(x, y)$ where f_{xx} is close to zero
	a point, where $f_x = 0$ and $f_y \neq 0$
	a point, where $f_y = 0$ and $f_x \neq 0$
	the point, where $ \nabla f $ is largest
	a local maximum of $f(x, y)$ under the constraint $g(x, y) = 0$.
	a local minimum of $f(x, y)$ under the constraint $g(x, y) = 0$.



Problem 4) (10 points)


Find and classify all the extrema of the function $f(x, y) = x^5 + y^3 - 5x - 3y$. This function measures “eat temptation” in the x =Easy- y =Tasty plane. Is there a global minimum or global maximum?



The “Easy-Tasty plane” was introduced in the XKCD cartoon titled “F&#% Grapefruits”.

Problem 5) (10 points)

After having watched the latest Disney movie “Tangled”, we want to build a hot air balloon with a cuboid mesh of dimension x, y, z which together with the top and bottom fortifications uses wires of total length $g(x, y, z) = 6x + 6y + 4z = 32$. Find the balloon with maximal volume $f(x, y, z) = xyz$.



Problem 6) (10 points)

a) (8 points) Find the tangent plane to the surface $f(x, y, z) = x^2 - y^2 + z = 6$ at the point $(2, 1, 3)$.

b) (2 points) A curve $\vec{r}(t)$ on that tangent plane of the function $f(x, y, z)$ in a) has constant speed $|\vec{r}'| = 1$ and passes through the point $(2, 1, 3)$ at $t = 0$. What is $\frac{d}{dt}f(\vec{r}(t))$ at $t = 0$?

Problem 7) (10 points)

a) (5 points) Estimate $\sqrt{\sin(0.0004) + 1.001^2}$ using linear approximation.

b) (5 points) We know $f(0, 0) = 1$, $D_{\langle \frac{3}{5}, \frac{4}{5} \rangle}f(0, 0) = 2$ and $D_{\langle -\frac{4}{5}, \frac{3}{5} \rangle}f(0, 0) = -1$. If $L(x, y)$ is the linear approximation to $f(x, y)$ at the point $(0, 0)$, find $L(0.06, 0.08)$.

Problem 8) (10 points)

a) (5 points) Find the following double integral

$$\int_0^1 \int_{x^2}^{\sqrt{x}} \frac{\pi \sin(\pi y)}{y^2 - \sqrt{y}} dy dx .$$

b) (5 points) Evaluate the following double integral

$$\iint_R \frac{\sin(\pi\sqrt{x^2 + y^2})}{\sqrt{x^2 + y^2}} dx dy$$

over the region

$$R = \{x^2 + y^2 \leq 1, x > 0\} .$$

Problem 9) (10 points)

a) (8 points) Find the surface area of the surface parametrized as

$$\vec{r}(u, v) = \langle u - v, u + v, (u^2 - v^2)/2 \rangle ,$$

where (u, v) is in the unit disc $R = \{u^2 + v^2 \leq 1\}$.

b) (2 points) Give a nonzero vector \vec{n} normal to the surface at $\vec{r}(4, 2) = \langle 2, 6, 6 \rangle$.

Problem 10) (10 points)

a) (6 points) Integrate

$$\int_0^{\pi/2} \int_x^{\pi/2} \frac{\cos(y)}{y} dy dx$$

b) (4 points) Find the moment of inertia

$$\iint_R (x^2 + y^2) dy dx ,$$

where R is the ring $1 \leq x^2 + y^2 \leq 9$.