

Name:

MWF 9 Jun-Hou Fung
MWF 9 Koji Shimizu
MWF 10 Matt Demers
MWF 10 Dusty Grundmeier
MWF 10 Erick Knight
MWF 11 Oliver Knill
MWF 11 Kate Penner
MWF 12 Yusheng Luo
MWF 12 YongSuk Moon
TTH 10 Will Boney
TTH 10 Peter Smillie
TTH 10 Chenglong Yu
TTH 11:30 Lukas Brantner
TTH 11:30 Yu-Wen Hsu

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3,8, we need to see **details** of your computation.
- All functions can be differentiated arbitrarily often unless otherwise specified.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) True/False questions (20 points), no justifications needed

- | | | |
|-----|--|--|
| 1) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $f(x, y) = 1$ is a curve, and near $(2, 3)$ one can write y as a function of x , then $y' = -f_y(2, 3)/f_x(2, 3)$. |
| 2) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $\iint_R f(x, y) \, dA = 0$, then the function $f(x, y)$ is everywhere zero on $R = \{x^2 + y^2 \leq 1\}$. |
| 3) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The directional derivative in the direction of the gradient is $ \nabla f $. |
| 4) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The linearization of $f(x, y) = x^3 + y^3$ at $(1, 1)$ is the quadratic function $L(x, y) = 3x^2 + 3y^2$. |
| 5) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The function $f(x, y) = x^2 + y^2$ satisfies the partial differential equation $D = f_{xx}f_{yy} - f_{xy}^2 = 1$. |
| 6) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The function x^2y^2 has no local minimum at $(0, 0)$ because the discriminant function D is zero there. |
| 7) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The double integral $\int_0^{\pi/4} \int_0^2 r^3 \, dr d\theta$ is the volume of the part of a solid cylinder $x^2 + y^2 \leq 4$ which is below the paraboloid $z = x^2 + y^2$ and above the xy plane. |
| 8) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The gradient of $f(x, y, z)$ at (x_0, y_0, z_0) is perpendicular to the level surface of f through (x_0, y_0, z_0) . |
| 9) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $f(x, y, z) = 3x - 4z$, then the minimal possible directional derivative $D_{\vec{u}}f$ at any point in space is -5 . |
| 10) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If (x, y) is not a critical point, then the directional derivative $D_{\vec{v}}f$ can take both positive and negative values for different choices of $\vec{v}$. |
| 11) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | Using linearization of $f(x, y) = x/y$ we can estimate $1.01/1.001 = f(1.01, 1.001) \sim 1 + 0.01 - 0.001 = 1.009$. |
| 12) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $(0, 0)$ is a critical point of $f(x, y)$ with nonzero discriminant $D = f_{xx}f_{yy} - f_{xy}^2$, we know that it is either a saddle, a global maximum or a global minimum. |
| 13) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | For a rectangular region R , Fubini tells that $\int_0^2 \int_0^3 f(x, y) \, dx dy = \int_0^2 \int_0^3 f(x, y) \, dy dx$ for any continuous function $f(x, y)$. |
| 14) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If a function $f(x, y)$ has only one critical point $(0, 0)$ in $G = \{x^2 + y^2 \leq 1\}$ which is a local maximum and $f(0, 0) = 1$, then $\iint_G f(x, y) \, dx dy > 0$. |
| 15) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $\vec{r}(t)$ is a curve in space for which the speed is 1 at all times and $f(x, y, z)$ is a function of three variables, then $d/dt f(\vec{r}(t)) = D_{\vec{r}'(t)}(f)$. |
| 16) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | $\int_0^1 \int_0^1 f_{xy}(x, y) \, dy dx = f(1, 1) - f(1, 0) - f(0, 1) + f(0, 0)$. |
| 17) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | If $f_{yy}(x, y) > 0$ everywhere, then f can not have any local maximum. |
| 18) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The double integral $\int_0^1 \int_0^1 x^2 - y^2 \, dx dy$ is the volume of the solid below the graph of $f(x, y) = x^2 - y^2$ and above the square $0 \leq x \leq 1, 0 \leq y \leq 1$ in the xy -plane. |
| 19) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | For any unit vector $\vec{v}$ and any differentiable function f , one has $D_{\vec{v}}(f) + D_{-\vec{v}}(f) = 0$. |
| 20) | <div style="border: 1px solid black; padding: 2px; display: inline-block;">T</div> <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">F</div> | The surfaces $x + y + z = 0$ and $x^2 + y^2 + z^2 + x + y + z = 0$ have the same tangent plane at $(0, 0, 0)$. |

Problem 2) (10 points)

a) (6 points) Match the regions with the corresponding polar double integrals



Enter A-F	Integral of $f(r, \theta)$	Enter A-F	Integral of $f(r, \theta)$
	$\int_0^{\pi/2} \int_0^{\pi/2} f(r, \theta) r \, dr d\theta$		$\int_0^{\pi/2} \int_{\theta}^{\pi/2} f(r, \theta) r \, dr d\theta$
	$\int_0^{\pi/2} \int_0^{\theta} f(r, \theta) r \, dr d\theta$		$\int_0^{\pi/2} \int_{\pi/2-\theta}^{\pi/2} f(r, \theta) r \, dr d\theta$
	$\int_0^{\pi/2} \int_0^{\pi/2-\theta} f(r, \theta) r \, dr d\theta$		$\int_0^{\pi/2} \int_{\pi/4}^{\pi/2} f(r, \theta) r \, dr d\theta$

b) (4 points) Match the partial differential equations (PDE's) for the functions $u(t, s)$ with their names. No justifications are needed.

Enter A,B,C,D here	PDE	Enter A,B,C,D here	PDE
	$u_t + uu_s - u_{ss} = 0$		$u_{tt} - u_{ss} = 0$
	$u_{tt} + u_{ss} = 0$		$u_t - u_{ss} = 0$

A) Wave equation
 B) Heat equation
 C) Burgers equation
 D) Laplace equation

Problem 3) (10 points)

a) (7 points) Find and classify all the critical points of the function

$$f(x, y) = 5 + 3x^2 + 3y^2 + y^3 + x^3 .$$

b) (3 points) Is there a global maximum or a global minimum for $f(x, y)$?

Problem 4) (10 points)



A **solid bullet** made of a half sphere and a cylinder has the volume $V = 2\pi r^3/3 + \pi r^2 h$ and surface area $A = 2\pi r^2 + 2\pi r h + \pi r^2$. Doctor Manhattan designs a bullet with fixed volume and minimal area. With $g = 3V/\pi = 1$ and $f = A/\pi$ he therefore minimizes

$$f(h, r) = 3r^2 + 2rh$$

under the constraint

$$g(h, r) = 2r^3 + 3r^2 h = 1 .$$

Use the Lagrange method to find a local minimum of f under the constraint $g = 1$.




Problem 5) (10 points)

A region R in the plane shown to the right is called the “**blob of nothingness**”. It does not have any purpose nor meaning. It just sits there. The region is given in polar coordinates as $0 \leq r \leq \theta(\pi - \theta)$ for $0 \leq \theta \leq \pi$. Find the area

$$\int \int_R 1 \, dx dy$$

of this nihilistic object.



Problem 6) (10 points)

a) (4 points) If

$$f(x, y) = y \cos(x - y) ,$$

find equation of plane tangent to $z = f(x, y)$ at the point $(2, 2, 2)$.

b) (3 points) Find the equation of the tangent line to $f(x, y) = 2$ at $(2, 2)$.

c) (3 points) Estimate $f(2.1, 1.9)$ using linear approximation.

Problem 7) (10 points)

A **Harvard robot bee** flies along the curve

$$\vec{r}(t) = \langle t - t^3, 3t^2 - 3t \rangle$$

and measures the temperature $f(x, y)$. It flies over the target point $(0, 0)$ at time $t = 0$ and time $t = 1$. At each time, its sensor measures the temperature change $g'(t)$ where $g(t) = f(\vec{r}(t))$.

a) (5 points) Assume you knew that the gradient of f at $(0, 0)$ is $\langle a, b \rangle$. What are the values of $g'(t) = d/dt f(\vec{r}(t))$ at $t = 0$ and $t = 1$ in terms of a and b ?

b) (5 points) The bee measures $g'(0) = 3$ and $g'(1) = 3$. What is the gradient $\nabla f(0, 0) = \langle a, b \rangle$ of f at $(0, 0)$?




Image source: Harvard Press release on
robobees.seas.harvard.edu

Problem 8) (10 points)

A function $f(x, y)$ of two variables has level curves as shown in the picture. The function values at neighboring level curves differ by 1. [No justifications are needed in this problem. Naturally, since there are less points than boxes, some of the points A-G will appear more than once, but each box will only be filled with one letter.]

Enter A-G	is a point, where ...
	$f_x(x, y) = 0$ and $f_y(x, y) \neq 0$.
	$f_y(x, y) = 0$ and $f_x(x, y) \neq 0$.
	$f(x, y)$ has either a max or a min.
	$f(x, y)$ has a saddle point.
	$f(x, y)$ has no max nor min but is extremal under a constraint $y = c$ for some c .
	$f(x, y)$ has no max nor min but is extremal under a constraint $x = c$ for some c .
	the length of the gradient vector of f is largest among all points A-G.
	$D_{\langle 1/\sqrt{2}, 1/\sqrt{2} \rangle} f(x, y) = 0$ and $D_{\langle 1/\sqrt{2}, -1/\sqrt{2} \rangle} f(x, y) \neq 0$.
	$D_{\langle 1/\sqrt{2}, -1/\sqrt{2} \rangle} f(x, y) = 0$ and $D_{\langle 1/\sqrt{2}, 1/\sqrt{2} \rangle} f(x, y) \neq 0$.
	the tangent line to the curve is $x + y = d$ for some constant d .



Problem 9) (10 points)

Evaluate the following double integral

$$\int_0^1 \int_0^{(1-x)^2} \frac{x^3}{(1-\sqrt{y})^4} dy dx .$$

Problem 10) (10 points)

A mass point with position (x, y) is attached by springs to the points $A_1 = (0, 0)$, $A_2 = (2, 0)$, $A_3 = (0, 2)$, $A_4 = (2, 3)$, $A_5 = (3, 1)$. It has the potential energy

$$f(x, y) = 31 - 14x + 5x^2 - 12y + 5y^2$$

which is the sum of the squares of the distances from (x, y) to the 5 points. Find all extrema of f using the second derivative test. The minimum of f is the position, where the mass point has the lowest energy.

