

Name:

MWF 9 Jun-Hou Fung
MWF 9 Koji Shimizu
MWF 10 Matt Demers
MWF 10 Dusty Grundmeier
MWF 10 Erick Knight
MWF 11 Oliver Knill
MWF 11 Kate Penner
MWF 12 Yusheng Luo
MWF 12 YongSuk Moon
TTH 10 Will Boney
TTH 10 Peter Smillie
TTH 10 Chenglong Yu
TTH 11:30 Lukas Brantner
TTH 11:30 Yu-Wen Hsu

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see details of your computation.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 180 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
14		10
Total:		150

Problem 1) True/False questions (20 points). No justifications are needed.

- 1) T F The functions $e^{x^2+y^3-y}$ and $x^2 + y^3 - y$ have the same critical points.
- 2) T F The line $\vec{r}(t) = \langle t^2, t^2, t^2 \rangle$ hits the plane $x + y + z = 100$ at a right angle.
- 3) T F The quadric $x^2 - 2y^2 + z^2 = 5$ is a one sheeted hyperboloid.
- 4) T F The relation $|\vec{u} \times \vec{v}| = |\vec{u} \cdot \vec{v}|$ is only possible if at least one of the vectors \vec{u} and \vec{v} is the zero vector.
- 5) T F The partial differential equation $u_x = u_{tt}$ is called the **Heat equation**.
- 6) T F The curvature of the curve $\vec{r}(t) = \langle \sin(2t), \cos(2t)/\sqrt{2}, \cos(2t)/\sqrt{2} \rangle$ at $t = 0$ is equal to the curvature of the curve $\vec{s}(t) = \langle 0, \cos(3t), \sin(3t) \rangle$ at $t = 0$.
- 7) T F The space curve $\vec{r}(t) = \langle \sin(t), t^2, \cos(t) \rangle$ for $t \in [0, 10\pi]$ is located on a cylinder.
- 8) T F If a smooth function $f(x, y)$ has a global maximum, then it has a global minimum.
- 9) T F If $L(x, y)$ is the linearization of $f(x, y)$ at (x_0, y_0) and $\vec{s}(t)$ is the line tangent to the curve $\vec{r}(t)$ on $f = c$ at the point $\vec{r}(t_0) = \vec{s}(t_0) = (x_0, y_0)$ so that $|\vec{r}'(t_0)| = |\vec{s}'(t_0)| = 1$, then $|d/dtL(\vec{s}(t))| = |d/dtf(\vec{r}(t))|$ at the time $t = t_0$.
- 10) T F If \vec{F} is a gradient field and $\vec{r}(t)$ is a flow line defined by $\vec{r}'(t) = \vec{F}(\vec{r}(t))$, then the line integral $\int_0^1 \vec{F} \cdot d\vec{r}$ is either positive or zero.
- 11) T F The flux of the vector field $\vec{F} = \nabla f$ through the surface $f(x, y, z) = x^4 + y^4 + z^4 = 1$ is positive if the surface is oriented so that $\vec{r}_u \times \vec{r}_v$ points in the direction of the gradient of f .
- 12) T F If we extremize the function $f(x, y)$ under the constraint $g(x, y) = 1$, and the functions are the same $f = g$, all points on the constraint curve are extrema for f .
- 13) T F If a point (x_0, y_0) is a minimum of $f(x, y)$ under the constraint $g(x, y) = 1$, then it is also a local minimum of the function $f(x, y)$ without constraints.
- 14) T F If a vector field $\vec{F}(x, y)$ is a gradient field, then any line integral along any closed ellipse is zero.
- 15) T F The flux of an irrotational vector field is zero through any surface S in space.
- 16) T F The divergence of a gradient field $\vec{F}(x, y, z) = \nabla f(x, y, z)$ is everywhere zero.
- 17) T F The line integral of the vector field $\vec{F}(x, y, z) = \langle x, y, z \rangle$ along a circle in the xy - plane is zero.
- 18) T F For any solid E , the moment of inertia $\iiint_E x^2 + y^2 dx dy dz$ is always larger than the volume $\iiint_E 1 dx dy dz$ of E .
- 19) T F The curvature of a parametrized curve satisfying $|\vec{r}'(t)| = 1$ is bounded above by the length $|\vec{r}''|$ of the acceleration.
- 20) T F Given a vector field $\vec{F} = \langle P, Q, R \rangle$, the directional derivative of $\text{div}(\vec{F}(x, y, z))$ in the direction $\vec{v} = \langle 1, 0, 0 \rangle$ is $P_{xx} + Q_{xy} + R_{xz}$.

Problem 2) (6 points)


a) (6 points) Match the objects with their definitions



Enter 1-6	Object definition
	$\vec{r}(t) = \langle (2 + \cos(10t)) \cos(t), (2 + \cos(10t)) \sin(t), \sin(10t) \rangle$
	$\vec{F}(x, y, z) = \langle -y, x, 2 \rangle$
	$\vec{r}(t, s) = \langle (2 + \cos(s)) \cos(t), (2 + \cos(s)) \sin(t), \sin(s) \rangle$
	$x^2y^2z^2 = 0$
	$(x - 1)/5 = (y - 2)/10 = (z - 1)/3$
	$\vec{r}(t) = \langle \sin(t) + \cos(5t), \cos(t) + \cos(6t) \rangle$


b) (4 points) Match the solids with the triple integrals:

Enter A-D	3D integral computing volume
	$\int_0^{2\pi} \int_0^{\pi/4} \int_0^{1/\cos(\phi)} \rho^2 \sin(\phi) \, d\rho d\phi d\theta$
	$\int_0^{\pi} \int_{\pi/2}^{\pi} \int_0^{\sin(\phi)} \rho^2 \sin(\phi) \, d\rho d\phi d\theta$
	$\int_0^{\pi} \int_{\pi/2}^{\pi} \int_0^1 \rho^2 \sin(\phi) \, d\rho d\phi d\theta$
	$\int_0^{2\pi} \int_0^{\pi} \int_0^{2\pi-\theta} \rho^2 \sin(\phi) \, d\rho d\phi d\theta$



Problem 3) (10 points)

a) (6 points) The surfaces are given either as a parametrization or implicitly. Match them. Each surface matches one definition.




Enter A-F here	Function or parametrization
	$\vec{r}(u, v) = \langle u^2, v^2, u^2 + v^2 \rangle$
	$\vec{r}(u, v) = \langle (1 + \sin(u)) \cos(v), (1 + \sin(u)) \sin(v), u \rangle$
	$4x^2 + y^2 - 9z^2 = 1$
	$x - 9y^2 + 4z^2 = 1$
	$\vec{r}(u, v) = \langle u, v, \sin(u^2 + v^2) \rangle$
	$4x^2 + 9y^2 = 1$

b) (4 points) If the blank box is replaced by $\nabla f(5, 6)$ the statement becomes true or false. Determine which case we have. The function $f(x, y)$ is an arbitrary nice function like for example $f(x, y) = x - yx + y^2$. The curve $\vec{r}(t)$, wherever it appears, parametrizes the level curve $f(x, y) = f(5, 6)$ and has the property that $\vec{r}'(0) = \langle 5, 6 \rangle$.

True/False	Topic	Statement
	Linearization	$L(x, y) = f(5, 6) + \square \cdot \langle x - 5, y - 6 \rangle$
	Chain rule	$\frac{d}{dt} f(\vec{r}(t)) _{t=0} = \square \cdot \vec{r}'(0)$
	Steepest descent	f decreases at $(5, 6)$ most in the direction of \square
	Estimation	$f(5 + 0.1, 5.99) \sim f(5, 6) + \square \cdot \langle 0.1, -0.01 \rangle$
	Directional derivative	$D_{\vec{v}} f(5, 6) = \square \cdot \vec{v}, \vec{v} = 1$
	Level curve	of f through $(5, 6)$ has the form $\square \cdot \langle x - 5, y - 6 \rangle = 0$
	Vector projection	of $\nabla f(5, 6)$ onto \vec{v} is $\vec{v}(\vec{v} \cdot \square) / \vec{v} ^2$
	Tangent line	of $\vec{r}(t)$ at $(5, 6)$ is parametrized by $\vec{R}(s) = \langle 5, 6 \rangle + s \square$

Problem 4) (10 points)



Two ice cream scoops given by spheres

$$x^2 + y^2 + (z + 1)^2 = 1$$


and

$$(x - 1)^2 + (y - 1)^2 + (z - 2)^2 = 1$$

are enclosed by a cylinder which is tangent to both spheres. Find the equation of the cylinder.

Hint: consider the distance of a general point (x, y, z) to the line passing through the centers of the spheres.

Problem 5) (10 points)



Find a parametrization


$$\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$$

for the line obtained by intersecting the tangent plane Σ to the surface

$$x^2 + y^2 - z = 0$$

at $(-1, -1, 2)$ with the xy -plane.

Problem 6) (10 points)



The vector field


$$\vec{F}(x, y) = \langle P, Q \rangle = \langle y(x^4 - 2x^2), x(y^4 - 4y) \rangle$$

has the curl

$$f(x, y) = \text{curl}(\vec{F})(x, y) = Q_x(x, y) - P_y(x, y) .$$

Find and classify all critical points of f by deciding whether they are local maxima, local minima or saddle points. Is there a global maximum or global minimum of f ?

Problem 7) (10 points)



We want to minimize the volume of the union of a **sphere** of radius x and a **cube** of side length y under the constraint that the sum of the two surface areas is equal to 4. Find the minimal value using the Lagrange method.

Remark: You do not have to show any derivations of the volume and surface area of the sphere.

Problem 8) (10 points)

A solid E in space is determined by the inequalities

$$0 \leq z \leq 9 ,$$


$$z^2 - x^2 - y^2 \geq 4$$

and

$$x^2 + y^2 \leq 1 .$$

Find the volume of E .

Problem 9) (10 points)



A surface S is parametrized by


$$\vec{r}(u, v) = e^{-u^2} \langle 1, \sin(v), \cos(v) \rangle$$

where

$$0 \leq u \leq \sqrt{\pi}, u^2 \leq v \leq \pi .$$

Find its surface area.

Problem 10) (10 points)




What is the line integral $\int_C \vec{F} \cdot d\vec{r}$ of the vector field

$$\vec{F}(x, y) = \langle 1 + y + 2xy, y^2 + x^2 \rangle$$

along the boundary C of the planar “castle region” shown in the picture? Each of the 5 windows is a unit square and the base of the castle has length 9. The boundary consists of 6 curves which are all oriented so that the region is to the left.

Problem 11) (10 points)



Compute the line integral of the vector field

$$\vec{F}(x, y, z) = \langle \cos(x), 2 + \cos(y), e^z + x(y^2 + z^2) \rangle$$

along the curve $\vec{r}(t) = \langle t, \cos(t), \sin(t) \rangle$ with $0 \leq t \leq 3\pi$.

Hint: you might want to find a split $\vec{F} = \vec{G} + \vec{H}$ and compute line integrals of \vec{G} and \vec{H} separately.

Problem 12) (10 points)



A biker in the Harvard Hemenway gym pedals. Assume that the force of a foot is


$$\vec{F} = \langle 0, 0, x^3 - x^2 + \sqrt{2 + \sin(z)} \rangle$$

and that one of the feet moves on a path $C : \vec{r}(t) = \langle 2 \cos(t), 0, 2 \sin(t) \rangle$. How much work

$$\int_C \vec{F} \cdot d\vec{r}$$

is done by this foot, when pedaling 10 times which means $0 \leq t \leq 20\pi$?

Problem 13) (10 points)



X-Rays have intensity and direction and are given by a vector field

$$\vec{F}(x, y, z) = \langle z^7, \sin(z) + y + z^{77}, z + \cos(xy) + \sin(y) \rangle .$$

A **tonsil** is given in spherical coordinates as $\rho \leq \phi$. Find the flux of the X-Ray field \vec{F} through the surface $\rho = \phi$ of the tonsil. The surface is oriented with normal vectors pointing outside. **Remark:** The flux is the amount of **ionizing radiation** absorbed by the tissue. This X-ray exposure is measured in the unit **Gray** which corresponds to the radiation amount to deposit 1 **joule** of energy in 1 **kilogram** of matter and corresponds to about 100 **Rem**. A typical dental X-ray is reported to lead to about one tenth to one half of a Rem.