

Name:

MWF 9 Jameel Al-Aidroos

MWF 9 Dennis Tseng

MWF 10 Yu-Wei Fan

MWF 10 Koji Shimizu

MWF 11 Oliver Knill

MWF 11 Chenglong Yu

MWF 12 Stepan Paul

TTH 10 Matt Demers

TTH 10 Jun-Hou Fung

TTH 10 Peter Smillie

TTH 11:30 Aukosh Jagannath

TTH 11:30 Sebastian Vasey

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see details of your computation.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 180 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
14		10
Total:		150

Problem 1) True/False questions (20 points)

- 1) ☐ T ☒ F For any two nonzero vectors $\vec{v}, \vec{w}$ the vector $((\vec{v} \times \vec{w}) \times \vec{v}) \times \vec{v}$ is parallel to $\vec{w}$.

Solution:

Take $\vec{v} = \langle 1, 0, 0 \rangle, \vec{w} = \langle 0, 1, 0 \rangle$ so that $\vec{v} \times \vec{w} = \langle 0, 0, 1 \rangle$ and $(\vec{v} \times \vec{w}) \times \vec{v} = \langle 0, 1, 0 \rangle$ and $((\vec{v} \times \vec{w}) \times \vec{v}) \times \vec{v} = \langle 0, 0, 1 \rangle$.

- 2) ☐ T ☒ F The cross product satisfies the law $(\vec{u} \times \vec{v}) \times \vec{w} = \vec{u} \times (\vec{v} \times \vec{w})$.

Solution:

Take $\vec{v} = \vec{w}$, then the right hand side is the zero vector while the left hand side is not zero in general (for example if $\vec{u} = \vec{i}, \vec{v} = \vec{j}$).

- 3) ☒ T ☐ F If the curvature of a smooth curve $\vec{r}(t)$ in space is defined and zero for all t , then the curve is part of a line.

Solution:

One can see that with the formula $\kappa(t) = |\vec{r}'(t) \times \vec{r}''(t)|/|\vec{r}'(t)|^3$ which shows that the acceleration $\vec{r}''(t)$ is in the velocity direction at all times. One can also see it intuitively or with the definition $\kappa(t) = |\vec{T}'(t)|/|\vec{T}'(t)|$. If curve is not part of a line, then $\vec{T}'$ has to change which means that κ is not zero somewhere.

- 4) ☒ T ☐ F The curve $\vec{r}(t) = (1 - t)A + tB, t \in [0, 1]$ connects the point A with the point B .

Solution:

The curve is a parameterization of a line and for $t = 0$, one has $\vec{r}(0) = A$ and for $t = 1$ one has $\vec{r}(1) = B$.

- 5) ☒ T ☐ F For every c , the function $u(x, t) = (2 \cos(ct) + 3 \sin(ct)) \sin(x)$ is a solution to the wave equation $u_{tt} = c^2 u_{xx}$.

Solution:

Just differentiate.

- 6)

T	F
---	---

 The length of the curve $\vec{r}(t) = (t, \sin(t))$, where $t \in [0, 2\pi]$ is $\int_0^{2\pi} \sqrt{1 + \cos^2(t)} dt$.

Solution:

The speed at time t is $|\vec{r}'(t)| = \sqrt{1 + \cos^2(t)}$.

- 7)

T	F
---	---

 Let (x_0, y_0) be the maximum of $f(x, y)$ under the constraint $g(x, y) = 1$. Then $f_{xx}(x_0, y_0) < 0$.

Solution:

While this would be true for $g(x, y) = f(y)$, where the constraint is a straight line parallel to the y axis, it is false in general.

- 8)

T	F
---	---

 The function $f(x, y, z) = x^2 - y^2 - z^2$ decreases in the direction $(2, -2, -2)/\sqrt{8}$ at the point $(1, 1, 1)$.

Solution:

It **increases** in that direction.

- 9)

T	F
---	---

 Assume $\vec{F}$ is a vector field satisfying $|\vec{F}(x, y, z)| \leq 1$ everywhere. For every curve $C : \vec{r}(t)$ with $t \in [0, 1]$, the line integral $\int_C \vec{F} \cdot d\vec{r}$ is less or equal than the arc length of C .

Solution:

$$|\vec{F} \cdot \vec{r}'| \leq |\vec{F}| |\vec{r}'| \leq |\vec{r}'|$$

- 10)

T	F
---	---

 Let $\vec{F}$ be a vector field which coincides with the unit normal vector $\vec{N}$ for each point on a curve C . Then $\int_C \vec{F} \cdot d\vec{r} = 0$.

Solution:

The vector field is orthogonal to the tangent vector to the curve.

- 11)

T	F
---	---

 If for two vector fields $\vec{F}$ and $\vec{G}$ one has $\text{curl}(\vec{F}) = \text{curl}(\vec{G})$, then $\vec{F} = \vec{G} + (a, b, c)$, where a, b, c are constants.

Solution:

One can also have $\vec{F} = \vec{G} + \text{grad}(f)$ which are vectorfields with the same curl.

- 12)

T	F
---	---

 If a nonempty quadric surface $g(x, y, z) = ax^2 + by^2 + cz^2 = 5$ can be contained inside a finite box, then $a, b, c \geq 0$.

Solution:

If one or two of the constants a, b, c are negative, we have a hyperboloid which all can not be contained into a finite space. If all three are negative, then the surface is empty.

- 13)

T	F
---	---

 If $\text{div}(\vec{F})(x, y, z) = 0$ for all (x, y, z) , then $\text{curl}(\vec{F}) = (0, 0, 0)$ for all (x, y, z) .

Solution:

There are counter examples: take $(-y, x, 0)$ for example.

- 14)

T	F
---	---

 If in spherical coordinates the equation $\phi = \alpha$ (with a constant α) defines a plane, then $\alpha = \pi/2$.

Solution:

Otherwise, it is would be a cone (or for $\alpha = 0$ or $\alpha = \pi$ a half line).

- 15)

T	F
---	---

 The divergence of the gradient of any $f(x, y, z)$ is always zero.

Solution:

$\text{div}(\text{grad}(f)) = \Delta f$ is the Laplacian of f .

- 16)

T	F
---	---

 For every vector field $\vec{F}$ the identity $\text{grad}(\text{div}(\vec{F})) = \vec{0}$ holds.

Solution:

$F = (x^2, y^2, z^2)$ has $\text{div}(F) = 2x + 2y + 2z$ which has a nonzero gradient $\nabla f = \langle 2, 2, 2 \rangle$.

- 17)

T	F
---	---

 For every function f , one has $\text{div}(\text{curl}(\text{grad}(f))) = 0$.

Solution:

Both because $\text{div}(\text{curl}(F)) = 0$ and $\text{curl}(\text{grad}(f)) = 0$.

- 18)

T

F

 If $\vec{F}$ is a vector field in space then the flux of $\vec{F}$ through any closed surface S is 0.

Solution:

While it is true that the flux of $\text{curl}(F)$ vanishes through every closed surface, this is not true for $\vec{F}$ itself. Take for example $F = (x, y, z)$.

- 19)

T

F

 The flux of the vector field $\vec{F}(x, y, z) = (y + z, y, -z)$ through the boundary of a solid region E is equal to the volume of E .

Solution:

By the divergence theorem, the flux through the boundary is $\int \int \int_E \text{div}(\vec{F}) \, dV$ but $\text{div}(\vec{F}) = 0$. So the flux is zero.

- 20)

T

F


 For every function $f(x, y, z)$, there exists a vector field $\vec{F}$ such that $\text{div}(\vec{F}) = f$.

Solution:


In order to solve $P_x + Q_y + R_z = f$ just take $\vec{F} = (0, 0, \int_0^z f(x, y, w) \, dw)$.

Problem 2) (10 points)


Problem 2a) (5 points) Match the equations with the objects. No justifications are needed.




I




II




III




IV




V



VI



VII



VIII

Enter I,II,III,IV,V,VI,VII,VIII here	Equation
	$g(x, y, z) = \cos(x) + \sin(y) = 1$
	$y = \cos(x) - \sin(x)$
	$\vec{r}(t) = \langle \cos(t), \sin(t) \rangle$
	$\vec{r}(u, v) = \langle \cos(u), \sin(v), \cos(u) \sin(v) \rangle$
	$\vec{F}(x, y, z) = \langle \cos(x), \sin(x), 1 \rangle$
	$z = f(x, y) = \cos(x) + \sin(y)$
	$g(x, y) = \cos(x) - \sin(y) = 1$
	$\vec{F}(x, y) = \langle \cos(x), \sin(y) \rangle$

Solution:

Enter I,II,III,IV,V,VI,VII,VIII here	Equation
V	$g(x, y, z) = \cos(x) + \sin(y) = 1$
VIII	$y = \cos(x) - \sin(x)$
I	$\vec{r}(t) = (\cos(t), \sin(t))$
II	$\vec{r}(u, v) = (\cos(u), \sin(v), \cos(u) \sin(v))$
VI	$\vec{F}(x, y, z) = (\cos(x), \sin(x), 1)$
IV	$z = f(x, y) = \cos(x) + \sin(y)$
VII	$g(x, y) = \cos(x) - \sin(y) = 1$
III	$\vec{F}(x, y) = (\cos(x), \sin(x))$

Problem 2b) (5 points) Mark with a cross in the column below "irrotational" if a vector fields is conservative (that is if $\text{curl}(\vec{F})(x, y, z) = (0, 0, 0)$ for all points (x, y, z)). Similarly, mark the fields which are incompressible (that is if $\text{div}(\vec{F})(x, y, z) = 0$ for all (x, y, z)). No justifications are needed.

Vectorfield	irrotational $\text{curl}(\vec{F}) = \vec{0}$	incompressible $\text{div}(\vec{F}) = 0$
$\vec{F}(x, y, z) = \langle -5, 5, 3 \rangle$		
$\vec{F}(x, y, z) = \langle x, y, z \rangle$		
$\vec{F}(x, y, z) = \langle -y, x, z \rangle$		
$\vec{F}(x, y, z) = \langle x^2 + y^2, xyz, x - y + z \rangle$		
$\vec{F}(x, y, z) = \langle x - 2yz, y - 2zx, z - 2xy \rangle$		

Solution:

Vectorfield	conservative $\text{curl}(\vec{F}) = \vec{0}$	incompressible $\text{div}(\vec{F}) = 0$
$\vec{F}(x, y, z) = \langle -5, 5, 3 \rangle$	X	X
$\vec{F}(x, y, z) = \langle x, y, z \rangle$	X	
$\vec{F}(x, y, z) = \langle -y, x, z \rangle$		
$\vec{F}(x, y, z) = \langle x^2 + y^2, xyz, x - y + z \rangle$		
$\vec{F}(x, y, z) = \langle x - 2yz, y - 2zx, z - 2xy \rangle$		

Problem 3) (10 points)

a) (2 points) What is the area of the triangle A, B, P , where $A = (1, 1, 1), B = (1, 2, 3)$ and $P = (3, 2, 4)$?

b) (2 points) Find the distance between the point P and the line L passing through the points A with B .

Let E be a general parallelogram in three dimensional space defined by two vectors $\vec{u}$ and $\vec{v}$.

c) (3 points) Express the diagonals of the parallelogram as vectors in terms of $\vec{u}$ and $\vec{v}$.

d) (3 points) What is the relation between the length of the crossproduct of the diagonals and the area of the parallelogram?

e) (3 points) Assume that the diagonals are perpendicular. What is the relation between the lengths of the sides of the parallelogram?

Solution:

a) The area is half of the cross product of $\vec{AB}$ and $\vec{AP}$ which is $(0, 1, 2) \times (2, 1, 3)$ which is $|(1, 4, -2)|$ which is $\sqrt{21}$. The triangle has the area $\sqrt{21}/2$.

b) The distance formula is $|\vec{AB} \times \vec{AP}|/|\vec{AB}| = |(1, 4, -2)|/|(0, 1, 2)| = \sqrt{\frac{21}{5}}$.

c) first diagonal $\vec{u} + \vec{v}$, second diagonal $\vec{u} - \vec{v}$.

d) $(\vec{u} + \vec{v}) \times (\vec{u} - \vec{v}) = 2\vec{v} \times \vec{u} = 2$ times area of parallelogram.

e) $(\vec{u} + \vec{v}) \cdot (\vec{u} - \vec{v}) = |\vec{u}|^2 - |\vec{v}|^2 = 0$, so that $|\vec{u}| = |\vec{v}|$.

Problem 4) (10 points)

The height of the ground near the Simplon pass in Switzerland is given by the function

$$f(x, y) = -x - \frac{y^3}{3} - \frac{y^2}{2} + \frac{x^2}{2}.$$

There is a lake in that area as you can see in the photo.

a) (7 points) Find and classify all the critical points of f and tell from each of them, whether it is a local maximum, a local minimum or a saddle point.

b) (3 points) For any pair of two different critical points A, B found in a) let $C_{a,b}$ be the line segment connecting the points, evaluate the line integral $\int_{C_{a,b}} \nabla f \cdot d\vec{r}$.




Photo of the lake in the Swiss alps near the Simplon mountain pass.

Solution:

a) The gradient is $\nabla f(x, y) = \langle x - 1, -y - y^2 \rangle$. This gradient vanishes if $x = 1$ and $y = -1$ or $y = 0$. So, there are two critical points $(1, -1), (1, 0)$. The Hessian matrix is

$$H(x, y) = \begin{bmatrix} 1 & 0 \\ 0 & -1 - 2y \end{bmatrix}.$$

point	discriminant	f_{xx}	nature
$(1, -1)$	$D = 1$	1	min
$(1, 0)$	$D = -1$	1	saddle



b) By the fundamental theorem of line integrals, the line integral between the two points is the difference of the potentials which is $f(1, -1) - f(1, 0) = (-1 + 1/3 - 1/2 + 1/2) - (-1 + 1/2) = -1/6$. Also the answer $1/6$ is correct of course, since we did not specify the direction.

Problem 5) (10 points)

Find the volume of the largest rectangular box with sides parallel to the coordinate planes that can be inscribed in the ellipsoid $\frac{x^2}{4} + \frac{y^2}{9} + \frac{z^2}{25} = 1$.

Solution:

The volume of the box is $8xyz$. The Lagrange equations are

$$\begin{aligned} 8yz &= \lambda x/2 \\ 8xz &= \lambda 2y/9 \\ 8xy &= \lambda 2z/25 \end{aligned}$$

$$\frac{x^2}{4} + \frac{y^2}{9} + \frac{z^2}{25} - 1 = 0$$

We can solve this by solving the first three equations for λ and expressing y, z by x , plugging this into the fourth equation. An other way to solve this is to multiply the first equation with x , the second with y and third with z .

The solution is $\boxed{x = 2/\sqrt{3}, y = \sqrt{3}, z = 5/\sqrt{3}}$. The maximal volume is $8xyz = 80/\sqrt{3}$.

Problem 6) (10 points)

Evaluate


$$\int_0^8 \int_{y^{1/3}}^2 \frac{y^2 e^{x^2}}{x^8} dx dy.$$

Solution:

This type II integral can not be computed as it is. We write it as a type I integral: from the boundary relation $x = y^{1/3}$ we obtain $y = x^3$ and $y = 8$ corresponds to $x = 2$:

$$\begin{aligned} &\int_0^2 \int_0^{x^3} \frac{y^2 e^{x^2}}{x^8} dy dx \\ &\int_0^2 \frac{x^9 e^{x^2}}{3 x^8} dx \\ &\int_0^2 \frac{x}{3} e^{x^2} dx \\ &e^{x^2}/6 \Big|_0^2 = (e^4 - 1)/6 \end{aligned}$$

The result is $\boxed{\frac{e^4 - 1}{6}}$.



Problem 7) (10 points)

In this problem we evaluate $\int \int_D \frac{(x-y)^4}{(x+y)^4} dx dy$, where D is the triangular region bounded by the x and y axis and the line $x + y = 1$.

a) (3 points) Find the region R in the uv -plane which is transformed into D by the change of variables $u = x - y, v = x + y$. (It is enough to draw a carefully labeled picture of R .)

b) (3 points) Find the Jacobian $\frac{\partial(x,y)}{\partial(u,v)}$ of the transformation $(x, y) = (\frac{u+v}{2}, \frac{v-u}{2})$.

c) (4 points) Evaluate $\int \int_D \frac{(x-y)^4}{(x+y)^4} dx dy$ using the above defined change of variables.

Hint. The general topic of change of variables does not appear this semester. You can solve the problem nevertheless, when given the formula $\frac{\partial(x,y)}{\partial(u,v)} = x_u y_v - x_v y_u$ for the integration factor (analogous to r when changing to polar coordinates, or $\rho^2 \sin(\phi)$ when going to spherical coordinates). The integral in c) becomes then $\int \int_R u^4/v^4 du dv$. The region R is the triangle bounded by the edges $(0, 0), (1, 1), (-1, 1)$.

Solution:



a) Take R with the edges $(0, 0), (1, 1)$ and $(-1, 1)$.

The region R is the red triangle shown in the picture. You get that triangle by mapping the points of the triangle D with the map $T(x, y) = (x - y, x + y)$. (This is analogue to the coordinate change $T(r, \theta) = (r \cos(\theta), r \sin(\theta))$.)

$$(0, 0) \Rightarrow (0, 0)$$

$$(1, 0) \Rightarrow (1, 1)$$

$$(0, 1) \Rightarrow (-1, 1)$$



These are the corners of the red triangle.

b) The Jacobian is $x_u y_v - x_v y_u = (1/4 + 1/4) = 1/2$.

(The Jacobian in the case of polar coordinates is r .)

c) The integral is best evaluated as a type II integral:

$$\frac{1}{2} \int_0^1 \int_{-v}^v u^4/v^4 du dv = \frac{1}{2} \int_0^1 2v/5 dv = \int_0^1 v/5 dv = \boxed{1/10}.$$

Problem 8) (10 points)

a) (3 points) Find all the critical points of the function $f(x, y) = -(x^4 - 8x^2 + y^2 + 1)$.

b) (3 points) Classify the critical points.

c) (2 points) Locate the local and absolute maxima of f .

d) (2 points) Find the equation for the tangent plane to the graph of f at each absolute maximum.

Solution:

a) $(\pm 2, 0)$ and $(0, 0)$.

b) $(-2, 0)$ is a local maximum with value 15.

$(0, 0)$ is a saddle with value -1 .

$(2, 0)$ is a maximum with value 15.

c) The local maxima are $(\pm 2, 0)$. They are also the absolute maxima because f decays at infinity.

d) To calculate the tangent plane at the maximum, write the graph of f as a level surface $g(x, y, z) = z - f(x, y)$. The gradient of g is orthogonal to the surface. We have $\nabla g = (0, 0, 1)$ so that the tangent plane has the equation $z = d = \text{const}$. Plugging in the point $(\pm 2, 0, 15)$ shows that $z = 15$ is the equation for the tangent plane for both maxima.

Problem 9) (10 points)


Find the volume of the wedge shaped solid that lies above the xy -plane and below the plane $z = x$ and within the cylinder $x^2 + y^2 = 4$.

Solution:

Use polar coordinates and note that the wedge is above the right side of the unit disc:

$$\int_0^2 \int_{-\pi/2}^{\pi/2} r^2 \cos(\theta) \, d\theta dr = 16/3$$

The solution is $16/3$.



Problem 10) (10 points)

Let the curve C be parametrized by $\vec{r}(t) = \langle t, \sin t, t^2 \cos t \rangle$ for $0 \leq t \leq \pi$. Let $f(x, y, z) = z^2 e^{x+2y} + x^2$ and $\vec{F} = \nabla f$. Find $\int_C \vec{F} \cdot d\vec{r}$.

Solution:

Use the fundamental theorem of line integrals. The result is $f(r(\pi)) - f(r(0)) = f(\pi, 0, -\pi^2) - f(0, 0, 0) = \pi^4 e^\pi + \pi^2 - 0 = \boxed{\pi^4 e^\pi + \pi^2}$.

Problem 11) (10 points)

A cylindrical building $x^2 + (y - 1)^2 = 1$ is intersected with the paraboloid $z = 4 - x^2 - y^2$.

- Parametrize the intersection curve and set up an integral for its arc length.
- Find a parametrization of the surface obtained by intersecting the paraboloid with the solid cylinder $x^2 + (y - 1)^2 \leq 1$ and set up an integral for its surface area.

Solution:

a) $\vec{r}(t) = \langle \cos(t), \sin(t) + 1, 4 - \cos^2(t) - (1 + \sin(t))^2 \rangle$.

Write down $\int_0^{2\pi} |r'(t)| dt = \int_0^{2\pi} \sqrt{3 + 2 \cos(2t)} dt$.

b) $\vec{r}(u, v) = \langle u, v, 4 - u^2 - v^2 \rangle$.

So, the integral is

$$\int \int_{u^2 + (v-1)^2 \leq 1} \sqrt{1 + 4u^2 + 4v^2} du dv = \int_{-1}^1 \int_{-\sqrt{1-u^2}+1}^{\sqrt{1-u^2}+1} \sqrt{1 + 4u^2 + 4v^2} dv du$$

Problem 12) (10 points)

Evaluate the line integral of the vector field $\vec{F}(x, y) = \langle y^2, x^2 \rangle$ in the clockwise direction around the triangle in the xy -plane defined by the points $(0, 0)$, $(1, 0)$ and $(1, 1)$ in two ways:

- (5 points) by evaluating the three line integrals.
- (5 points) using Green's theorem.

Solution:

The problem asks to do this in the clockwise direction. We do it in the counterclockwise direction and change then the sign.

$$\text{a) } \int_0^1 \vec{F}(t, 0) \cdot (1, 0) dt + \int_0^1 \vec{F}(1, t) \cdot (0, 1) dt + \int_0^1 \vec{F}(1-t, 1-t) \cdot (-1, -1) dt = 0 + 1 - 2/3 = 1/3.$$

So, the result for the clockwise direction is $\boxed{-1/3}$.

b) The curl of F is $2x - 2y$.

$$\int_0^1 \int_0^x (2x - 2y) dy dx = \int_0^1 2x^2 - x^2 dx = 1/3$$

So, the result for the clockwise direction is $\boxed{-1/3}$.

Problem 13) (10 points)

Use Stokes theorem to evaluate the line integral of $\vec{F}(x, y, z) = (-y^3, x^3, -z^3)$ along the curve $\vec{r}(t) = \langle \cos(t), \sin(t), 1 - \cos(t) - \sin(t) \rangle$ with $t \in [0, 2\pi]$.

Solution:

The curve is contained in the graph of the function $f(x, y) = 1 - x - y$. That surface is parameterized by $\vec{r}(u, v) = \langle u, v, 1 - u - v \rangle$ and has the normal vector $\vec{r}_u \times \vec{r}_v = \langle 1, 0, -1 \rangle \times \langle 0, 1, -1 \rangle = \langle 1, 1, 1 \rangle$. The curl of F is $\langle 0, 0, 3x^2 + 3y^2 \rangle$ so that $\vec{F}(\vec{r}(u, v)) \cdot (\vec{r}_u \times \vec{r}_v) = 3(x^2 + y^2)$. The surface is parameterized over the region $R = \{u^2 + v^2 \leq 1\}$ and $\int \int_S \vec{F} \cdot d\vec{S} = \int_0^1 \int_0^{2\pi} 3r^3 d\theta dr = \boxed{\frac{3\pi}{2}}$.

Problem 14) (10 points)

Let S be the graph of the function $f(x, y) = 2 - x^2 - y^2$ which lies above the disk $\{(x, y) \mid x^2 + y^2 \leq 1\}$ in the xy -plane. The surface S is oriented so that the normal vector points upwards. Compute the flux $\int \int_S \vec{F} \cdot d\vec{S}$ of the vector field


$$\vec{F} = \left\langle -4x + \frac{x^2 + y^2 - 1}{1 + 3y^2}, 3y, 7 - z - \frac{2xz}{1 + 3y^2} \right\rangle$$

through S using the divergence theorem.

Solution:

We apply the divergence theorem to the region $E = \{0 \leq z \leq f(x, y), x^2 + y^2 \leq 1\}$. Using $\text{div}(F) = -2$, we get

$$\begin{aligned} \iiint \text{div}(F) dV &= \int_0^1 \int_0^{2\pi} \int_0^{2-r^2} (-2) r dr d\theta dz \\ &= (-2) \int_0^1 \int_0^{2\pi} (2-r^2) r d\theta dr \\ &= (-2)(2\pi)(2/2 - 1/4) = -3\pi . \end{aligned}$$



By the divergence theorem, this is the flux of $\vec{F}$ through the boundary of E which consists of the surface S , the cylinder $S_1 : r(u, v) = (\cos(u), \sin(u), v)$ with normal vector $r_u \times r_v = (-\sin(u), \cos(u), 0) \times (0, 0, 1) = (\cos(u), \sin(u), 0)$ plus the flux through the floor $S_2 : \vec{r}(u, v) = (v \sin(u), v \cos(u), 0)$ with normal vector $r_u \times r_v = (0, 0, -v)$. The flux through S_1 is

$$\begin{aligned} \iint_{S_1} \vec{F} \cdot dS &= \int_0^1 \int_0^{2\pi} \vec{F}(\cos(u), \sin(u), v) \cdot (\cos(u), \sin(u), 0) dudv \\ &= \int_0^1 \int_0^{2\pi} (-4 \cos^2(u) + 3 \sin^2(u)) dudv = -\pi . \end{aligned}$$

The flux through S_2 is

$$\begin{aligned} \iint_{S_2} \vec{F} \cdot dS &= \int_0^1 \int_0^{2\pi} F(v \sin(u), v \cos(u), 0) \cdot (0, 0, -v) dudv \\ &= \int_0^1 \int_0^{2\pi} (-7v) dudv = -7\pi . \end{aligned}$$

By the divergence theorem, $\iint_S \vec{F} \cdot dS + \iint_{S_1} \vec{F} \cdot dS + \iint_{S_2} \vec{F} \cdot dS = -3\pi$ so that $\iint_S \vec{F} \cdot dS = -3\pi + \pi + 7\pi = \boxed{5\pi}$.