

Homework 2: Vectors and Dot product

This homework is due Monday, 9/11 respectively Tuesday 9/12 at the beginning of class.

- 1 A **kite surfer** gets pulled with a force $\vec{F} = \langle 7, 1, 4 \rangle$. She moves with velocity $\vec{v} = \langle 4, -2, 1 \rangle$. The dot product of \vec{F} with \vec{v} is **power**. a) Find the angle between the \vec{F} and \vec{v} . b) Find the **vector projection** of the \vec{F} onto \vec{v} .
- 
- 2 Light shines long the vector $\vec{a} = \langle a_1, a_2, a_3 \rangle$ and reflects at the three coordinate planes where the angle of incidence equals the angle of reflection. Verify that the reflected ray is $-\vec{a}$. **Hint.** Reflect first at the xy -plane.
- 3 The **Harvard triple product** is defined in \mathbb{R}^3 as $\vec{u} \star \vec{v} \star \vec{w} = \vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{u}$. a) Are there three unit vectors $\vec{u}, \vec{v}, \vec{w}$ for which the Harvard product is zero? If yes, find an example.
b) Are there three unit vectors $\vec{u}, \vec{v}, \vec{w}$ for which the Harvard product is 3? If yes, find an example.
c) Are there three unit vectors $\vec{u}, \vec{v}, \vec{w}$ for which the Harvard product is -3 ? If yes, find an example.
d) Are there three unit vectors $\vec{u}, \vec{v}, \vec{w}$ for which the Harvard product is 2? If yes, give a reason.
- 4 a) Find the angle between a space diagonal of a cube and the diagonal in one of its faces.
b) The **hypercube** or **tesseract** has vertices $(\pm 1, \pm 1, \pm 1, \pm 1)$. Find the angle between the hyper diagonal connecting $(1, 1, 1, 1)$ with $(-1, -1, -1, -1)$ and the space diagonal connecting $(1, 1, 1, 1)$ with $(-1, -1, -1, 1)$.

- 5 a) Verify that if \vec{a}, \vec{b} are nonzero, then $\vec{c} = |\vec{a}|\vec{b} + |\vec{b}|\vec{a}$ bisects the angle between \vec{a}, \vec{b} if \vec{c} is not zero.
- b) Verify the parallelogram law $|\vec{a} + \vec{b}|^2 + |\vec{a} - \vec{b}|^2 = 2|\vec{a}|^2 + 2|\vec{b}|^2$.

Main definitions

Two points $P = (a, b, c)$ and $Q = (x, y, z)$ define a **vector** $\vec{v} = \langle x - a, y - b, z - c \rangle$. We also write $\vec{v} = \vec{PQ}$. The numbers v_1, v_2, v_3 in $\vec{v} = \langle v_1, v_2, v_3 \rangle$ are the **components** of \vec{v} . The **length** $|\vec{v}|$ of a vector $\vec{v} = \vec{PQ}$ is defined as the distance $d(P, Q)$ from P to Q . A vector of length 1 is called a **unit vector**. The **addition** is $\vec{u} + \vec{v} = \langle u_1, u_2, u_3 \rangle + \langle v_1, v_2, v_3 \rangle = \langle u_1 + v_1, u_2 + v_2, u_3 + v_3 \rangle$. The **scalar multiple** $\lambda\vec{u} = \lambda\langle u_1, u_2, u_3 \rangle = \langle \lambda u_1, \lambda u_2, \lambda u_3 \rangle$. The difference $\vec{u} - \vec{v}$ can be seen as $\vec{u} + (-\vec{v})$.

The **dot product** of two vectors $\vec{v} = \langle a, b, c \rangle$ and $\vec{w} = \langle p, q, r \rangle$ is defined as $\vec{v} \cdot \vec{w} = ap + bq + cr$. The **Cauchy-Schwarz inequality** tells $|\vec{v} \cdot \vec{w}| \leq |\vec{v}||\vec{w}|$.

The **angle** between two nonzero vectors is defined as the unique $\alpha \in [0, \pi]$ satisfying $\vec{v} \cdot \vec{w} = |\vec{v}| |\vec{w}| \cos(\alpha)$. Two vectors are called **orthogonal** or **perpendicular** if $\vec{v} \cdot \vec{w} = 0$. The zero vector $\vec{0}$ is orthogonal to any vector. For example, $\vec{v} = \langle 2, 3 \rangle$ is orthogonal to $\vec{w} = \langle -3, 2 \rangle$. The vector $P(\vec{v}) = \frac{\vec{v} \cdot \vec{w}}{|\vec{w}|^2} \vec{w}$ is called the **projection** of \vec{v} onto \vec{w} . The **scalar projection** $\frac{\vec{v} \cdot \vec{w}}{|\vec{w}|}$ is plus or minus the length of the projection of \vec{v} onto \vec{w} . The vector $\vec{b} = \vec{v} - P(\vec{v})$ is a vector orthogonal to \vec{w} . **Pythagoras tells:** if \vec{v} and \vec{w} are orthogonal, then $|\vec{v} - \vec{w}|^2 = |\vec{v}|^2 + |\vec{w}|^2$.