

## Homework 7: Other coordinates

This homework is due Friday, 9/22 resp Tuesday 9/26.

- 1 Change to cylindrical coordinates: a)  $(x, y, z) = (-3\sqrt{3}, 3, 27)$  b)  $(x, y, z) = (-4, 4, 10)$

Change to spherical coordinates: c)  $(x, y, z) = (-5\sqrt{3}, 0, 5)$  d)  $(x, y, z) = (-1, -1, \sqrt{2})$

- 2 a) Identify the surface  $r^2 = 1 + (z - 5)^2$ .  
 b) Identify the surface given in spherical coordinates as  $\rho \sin(\phi) = 2\rho \cos(\phi)$ .  
 c) Write the surface  $\sin^2(\phi) + \cos^2(\phi)/4 = 1/\rho^2$  in Cartesian coordinates.

- 3 a) Identify the surface  $2r^2 - z^2 = 1$ .  
 b) Identify the surface  $r^2 - z = 1$ .  
 b) Identify the surface  $\cos(\phi) = \rho \sin^2(\phi)$ .

- 4 a) Identify the surface whose equation is given in cylindrical coordinates as

$$\cos(\theta) + \sin(\theta) = 1/r .$$

- b) Identify the surface whose equation is given in spherical coordinates as

$$\rho^2(\sin^2(\phi) \sin^2(\theta) + \cos^2(\phi)) = 16 .$$

- 5 a) Use the Mathematica command "RevolutionPlot3D" to plot the surface which is given by

$$z = (\sin(7\theta) + \cos(7\theta))e^{-r^2} .$$

(See example Mathematica command below). b) Use the Mathematica command "SphericalPlot3D" to plot a bumpy sphere

$$\rho(\phi, \theta) = (2 + \cos(3\phi) + \sin(5\theta)) .$$

with  $0 \leq \phi \leq \pi$  and  $0 \leq \theta < 2\pi$ .

```
RevolutionPlot3D [Cos[r+t], {r, 0, Pi}, {t, 0, 2Pi}]
SphericalPlot3D [s+ t, {s, 0, Pi}, {t, 0, 2Pi}]
```

## Main definitions

A point  $(x, y)$  in the plane has **polar coordinates**  $r = \sqrt{x^2 + y^2} \geq 0, \theta = \arctan(y/x)$ . We have  $(x, y) = (r \cos(\theta), r \sin(\theta))$ . We chose arctan values in  $(-\pi/2, \pi/2]$  and add  $\pi$  if  $x < 0$  or  $x = 0, y < 0$ . This gives a value in  $\theta \in [0, 2\pi)$ . A point  $(x, y, z)$  in space has the **cylindrical coordinates**  $(r, \theta, z)$ , where  $(r, \theta)$  are the polar coordinates of  $(x, y)$ . A curve given in polar coordinates as  $r(\theta) = f(\theta)$  is called a **polar curve**. It can in Cartesian coordinates be described as  $\vec{r}(t) = \langle f(t) \cos(t), f(t) \sin(t) \rangle$ . **Spherical coordinates** use the distance  $\rho$  to the origin as well as two angles  $\theta$  and  $\phi$ . The first angle  $\theta$  is the polar angle in polar coordinates of the  $xy$  coordinates and  $\phi$  is the angle between the vector  $\vec{OP}$  and the  $z$ -axis. The relation is  $(x, y, z) = (\rho \cos(\theta) \sin(\phi), \rho \sin(\theta) \sin(\phi), \rho \cos(\phi))$ . We have

$$\begin{aligned} x &= \rho \cos(\theta) \sin(\phi), \\ y &= \rho \sin(\theta) \sin(\phi), \\ z &= \rho \cos(\phi) \end{aligned}$$