

## Homework 13: Gram Schmidt and QR

This homework is due on Friday, March 4, respectively on Tuesday, March 8, 2016.

- 1 Perform the Gram-Schmidt process on the two vectors  $\left\{ \begin{bmatrix} 5 \\ 4 \\ 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \\ 1 \end{bmatrix} \right\}$ .

- 2 Find an orthonormal basis of the hyper plane  $x_1 + x_2 + x_3 + x_4 = 0$  in  $\mathbf{R}^4$ .

- 3 As in the previous problem, we first find an orthonormal basis of the kernel of

$$A = \begin{bmatrix} 1 & -1 & 1 & -1 & 1 \\ 1 & 1 & -1 & 1 & 1 \end{bmatrix}.$$

Then find an orthonormal basis for the image.

- 4 Find the QR factorization of the following three matrices

$$A = \begin{bmatrix} 0 & -3 & 0 \\ 0 & 0 & 0 \\ 7 & 0 & 0 \\ 0 & 0 & 4 \end{bmatrix}, B = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, C = \begin{bmatrix} 3 & 4 \\ -4 & 3 \end{bmatrix}.$$

- 5 Find the QR factorization of the following matrices:

$$A = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & -1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 6 \end{bmatrix},$$

$$B = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, C = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}.$$

## Gram Schmidt and QR

The **Gram Schmidt orthogonalization process** produces from an arbitrary basis  $\mathcal{B} = \{\vec{v}_j\}$  an orthonormal basis  $\{\vec{u}_j\}$ . This goes as follows:  $\vec{w}_1 = \vec{v}_1$  and  $\vec{u}_1 = \vec{w}_1/|\vec{w}_1|$ . To construct  $\vec{u}_i$  once you've already constructed  $\vec{u}_1, \dots, \vec{u}_{i-1}$  so that they are orthonormal, make the new vector  $\vec{w}_i = \vec{v}_i - \text{proj}_{V_{i-1}}(\vec{v}_i)$ , where  $V_{i-1} = \text{span}(\vec{u}_1, \dots, \vec{u}_{i-1})$ , and then normalize  $\vec{w}_i$  to get  $\vec{u}_i = \vec{w}_i/|\vec{w}_i|$ . Then  $\{\vec{u}_1, \dots, \vec{u}_n\}$  is an orthonormal basis of  $V$  and the formulas

$$\vec{v}_1 = |\vec{v}_1|\vec{u}_1 = r_{11}\vec{u}_1$$

$$\vdots$$

$$\vec{v}_i = (\vec{u}_1 \cdot \vec{v}_i)\vec{u}_1 + \dots + (\vec{u}_{i-1} \cdot \vec{v}_i)\vec{u}_{i-1} + |\vec{w}_i|\vec{u}_i = r_{i1}\vec{u}_1 + \dots + r_{ii}\vec{u}_i$$

$$\vdots$$

$$\vec{v}_n = (\vec{u}_1 \cdot \vec{v}_n)\vec{u}_1 + \dots + (\vec{u}_{n-1} \cdot \vec{v}_n)\vec{u}_{n-1} + |\vec{w}_n|\vec{u}_n = r_{1n}\vec{u}_1 + \dots + r_{nn}\vec{u}_n$$

can be written in matrix form as  $A = QR$ ,

$$\begin{bmatrix} | & | & | \\ \vec{v}_1 & \cdots & \vec{v}_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & | & | \\ \vec{u}_1 & \cdots & \vec{u}_n \\ | & | & | \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ 0 & 0 & \cdots & r_{nn} \end{bmatrix},$$

where  $A$  and  $Q$  are  $m \times n$  matrices and  $R$  is an  $n \times n$  matrix.