

Homework 8: Basis

This homework is due on Wednesday, February 15, respectively on Thursday, February 16, 2017.

1 Which of the following sets are linear spaces? Check in each case the three properties characterizing a linear space. Only a brief explanation is needed (can be a picture too): a) $W = \{(x, y, z) \mid x + 2y + 3z = 0\}$

d) $W = \{(x, y, z) \mid x = y = z = 1\}$

c) $W = \{(x, y, z) \mid x = y = z\}$

d) $W = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$ e) $W = \{(x, y, z) \mid xyz = 0\}$

2 Let V, W be two different linear subspaces of the plane R^2 . Which are linear spaces?

a) (2 points) The union $V \cup W$ of V and W .

b) (2 points) The intersection $V \cap W$ of V and W .

c) (2 points) The set V^\perp of vectors perpendicular to V .

d) (2 points) the intersection $V \cap S$ of V with the unit sphere $S : x^2 + y^2 + z^2 = 1$.

e) (2 points) the difference $V \setminus W$, the points which are in V but not in W .

3 Check whether the given set of vectors is linearly independent

a) $\left\{ \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} \right\}$. b) $\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ 1 \\ -1 \end{bmatrix} \right\}$. c) $\left\{ \begin{bmatrix} 20 \\ 16 \end{bmatrix}, \begin{bmatrix} 2 \\ 18 \end{bmatrix}, \begin{bmatrix} 2 \\ 19 \end{bmatrix} \right\}$.

4 Find a basis for the image as well as as a basis for the kernel of the following matrices

a) $\begin{bmatrix} 7 & 0 & 7 \\ 2 & 3 & 8 \\ 9 & 0 & 9 \\ 5 & 6 & 17 \end{bmatrix}$, b) $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$. c) $\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$.

5 The orthogonal complement of a subspace V of R^n is the set V^\perp of all vectors in R^n that are perpendicular to every single vector

in V . Find a basis for the orthogonal complement in each case:

- The line L in R^5 spanned by $\begin{bmatrix} 1 & 2 & 2 & 1 & 1 \end{bmatrix}^T$, (If v is a row vector v^T denotes the corresponding column vector).
- The plane Σ in R^4 spanned by $\begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}^T$ and $\begin{bmatrix} 1 & -1 & -1 & 1 \end{bmatrix}^T$.
- The space $V = \{(0, 0)\}$ in the two-dimensional plane R^2 .

Basis

V is a **linear space** if 0 is in V , if $v + w$ is in V for all v, w in V and if λv is in V for every v in V and every λ in \mathbb{R} . Examples: kernels $V = \ker(A)$ or images $V = \text{im}(A)$ are linear spaces. If V, W are linear spaces and V is a subset of W , then V is called a **linear subspace** of W . A line through the origin for example is a linear subspace of \mathbb{R}^3 . A set \mathcal{B} of vectors $\{v_1, \dots, v_n\}$ **spans** V if every $v \in V$ is a sum of vectors in \mathcal{B} . A set \mathcal{B} is linear independent if $a_1 v_1 + \dots + a_n v_n = 0$ implies $a_1 = \dots = a_n = 0$. It is a **basis** of V if it both **spans** V and is linearly independent. Example: the standard basis vectors $\{e_1, \dots, e_n\}$ form a basis of \mathbb{R}^n . How do we determine whether a set of vectors is a basis of \vec{R}^n ? Place the vectors of \mathcal{B} as columns in a matrix A , then row reduce A . If every column of a matrix has a leading 1, then the set of column vectors \mathcal{B} are linearly independent and the kernel of A is $\{0\}$. How do we determine whether a set of vectors is linearly independent? Place the vectors as columns of a matrix and row reduce. If there is no free variable, then we have linear independence. Example: three vectors in \mathbb{R}^3 are linearly independent if they are not in a common plane.