

Homework 19: Eigenspaces

This homework is due on Friday, March 24, respectively on Tuesday, March 28, 2017.

- 1 Find all eigenvalues and eigenvectors for the following matrix

$$A = \begin{bmatrix} 2 & -1 & -1 & 0 & 1 \\ -1 & 2 & -1 & 0 & 1 \\ -1 & -1 & 4 & -1 & 0 \\ 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & -1 & 0 & 2 \end{bmatrix}.$$

What are the algebraic and geometric multiplicities? As a hint, we tell you that the eigenvalues 1, 1, 1, 3 appear. Use the trace trick to the 5th eigenvalue.

- 2 Assume that a 2×2 matrix has trace 9 and determinant 14. Find its eigenvalues and find a non-diagonal matrix which realizes the situation.

- 3 a) Verify that for any $n \times n$ matrix A , the matrix A and A^T have the same eigenvalues.

b) Assume A is invertible. What is the relation between the eigenvalues of A and A^{-1} ?



- 4 This is a classic problem from Otto Bretscher. The vector $A^n b$ gives pollution levels in the Silvaplana, Sils and St Moritz lake

n weeks after an oil spill. The matrix is $A = \begin{bmatrix} 0.7 & 0 & 0 \\ 0.1 & 0.6 & 0 \\ 0 & 0.2 & 0.8 \end{bmatrix}$

and $b = \begin{bmatrix} 100 \\ 0 \\ 0 \end{bmatrix}$ is the initial pollution level. Find a closed form solution for the pollution after n weeks.

5 a) Find all the eigenvalues and eigenvectors of the matrix

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 \\ 1 & 6 & 6 & 0 \\ 1 & 14 & 36 & 24 \end{bmatrix}.$$

b) Find the eigenvectors of A^3 , where A is the previous matrix.

c) Find the eigenvectors of $(A^T)^{-1}$, where A is the previous matrix.

Eigenspaces

A nonzero vector \vec{v} is called an **eigenvector**, if $Av = \lambda v$ for some λ . The set of eigenvectors is called the **eigenspace** E_λ . It is the kernel of $A - \lambda I_n$. The dimension of the eigenspace is called the **geometric multiplicity** of λ . There is a general result which tells that the geometric multiplicity of λ is always smaller or equal to the algebraic multiplicity.

Recall that A is similar to B if there exists an invertible S such that $B = S^{-1}AS$. If A and B are similar, then they have the same characteristic polynomial, the same eigenvalues and algebraic multiplicities as well as the same geometric multiplicities. Similar matrices also have the same trace $\text{tr}(A) = \lambda_1 + \dots + \lambda_n$ as well as determinant $\det(A) = \lambda_1 \cdots \lambda_n$. These formulas hold in general if we allow the eigenvalues λ_i to be complex. More on complex eigenvalues next week.