

Homework 21: Complex Eigenvalues

This homework is due on Wednesday, March 29, respectively on Thursday, March 30, 2017.

- 1 a) For $z = 2 + 4i$. Find $z + z^2 + z^3$.
- b) The log of a nonzero complex number $re^{i\theta} = r \cos(\theta) + ir \sin(\theta)$, where $r > 0$ and $0 \leq \theta < 2\pi$, is defined as $\log(re^{i\theta}) = \log r + i\theta$. Find the logarithm of $e \cdot i$. ("Ei" is the product of e and i . It means "Egg" in German.)
- c) Using logarithms we can define $w^z = e^{z \log w}$. What is i^i , the "eye for an eye" number?

- 2 a) First use the identity $(\cos(\theta) + i \sin(\theta))^2 = (e^{i\theta})^2 = e^{i2\theta} = \cos(2\theta) + i \sin(2\theta)$ to get the double angle formulas for $\cos(2\theta)$ and $\sin(2\theta)$.
- b) Express $\cos(4\theta)$ and $\sin(4\theta)$ as polynomials in $\cos(\theta)$, $\sin(\theta)$.

- 3 a) Find all the complex eigenvalues of the matrix $\begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$.

(Note that real numbers are a subset of complex numbers and also complex. Similarly as rational numbers are real numbers too).

- b) Verify that if λ is an eigenvalue, $\vec{v} = \begin{bmatrix} \lambda^3 \\ \lambda^2 \\ \lambda \\ 1 \end{bmatrix}$ is an eigenvector.

- 4 Find the eigenvalues and eigenvectors of the matrix

Hint: Do you recognize $A + 5I_6 + 3A^{-1}$?

$$\begin{bmatrix} 5 & 3 & 0 & 0 & 0 & 1 \\ 1 & 5 & 3 & 0 & 0 & 0 \\ 0 & 1 & 5 & 3 & 0 & 0 \\ 0 & 0 & 1 & 5 & 3 & 0 \\ 0 & 0 & 0 & 1 & 5 & 3 \\ 3 & 0 & 0 & 0 & 1 & 5 \end{bmatrix}$$

5 The matrix $A = \begin{bmatrix} 0.1 & 0.2 & 0.7 \\ 0.8 & 0.3 & 0.3 \\ 0.1 & 0.5 & 0.0 \end{bmatrix}$ is called a Markov matrix: in every column the entries add up to 1, so each column can be interpreted as a probability distribution.

a) Verify that A^T has the eigenvector $\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ with eigenvalue 1.

b) Why does A also have an eigenvalue 1? Find all other eigenvalues.

c) Find all eigenvectors of A .

Complex eigenvalues

Complex numbers are of the form $z = a + ib$. Add and multiply as real numbers, keeping in mind $i^2 = -1$. Euler found $e^{i\theta} = \cos(\theta) + i \sin(\theta)$ which for $\theta = \pi$ gives $e^{i\pi} + 1 = 0$ (the most beautiful formula in whole of mathematics), as it combines e , π , i , 1 and 0. As e is part of analysis and π is part of geometry and 0 is the additive neutral element and 1 the multiplicative neutral element, this identity combines analysis, geometry and algebra. The Euler identity leads to **de Moivre formulas** like $(\cos(\theta) + i \sin(\theta))^3 = (e^{i\theta})^3 = e^{i3\theta} = \cos(3\theta) + i \sin(3\theta)$. So that $\cos^3(\theta) - \cos(\theta) \sin^2(\theta) = \cos(3\theta)$ and $\cos^2(\theta) \sin(\theta) - \sin^3(\theta) = \sin(3\theta)$. Eigenvalues of matrices can become complex as the rotation-dilation matrix $A = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ shows which has the eigenvalues $a \pm ib$. The fundamental theorem of algebra assures that the sum of the algebraic multiplicities of all eigenvalues of a $n \times n$ matrix is equal to n .