1	
2	
3	
4	
5	
6	
7	
8	
9	
10	

# Name:

### LINEAR ALGEBRA AND VECTOR ANALYSIS

 $\mathrm{MATH}\ 22\mathrm{A}$ 

Total:

## Unit 28: Second Hourly Practice

Problems

#### Problem 28P.1 (10 points):

a) (4 points) You know the positive integer  $n^5$  is odd. Prove that n is odd.

b) (3 points) Prove or disprove: if a and b are irrational, then ab is irrational.

c) (3 points) Prove or disprove: if a and b are irrational, then a + b is irrational.

#### Problem 28P.2 (10 points, each sub problem is one point):

- a) What is the name of the differential equation  $f_t = f_{xx}$ ?
- b) What assumptions need to hold so that  $f_{xy} = f_{yx}$  is true?
- c) The gradient  $\nabla f(x_0)$  has a relation to f(x) = c with  $c = f(x_0)$ . Which one?
- d) The linear approximation of f at  $x_0$  is  $L(x) = f(x_0) + \dots$  Complete the formula.
- e) Assume f has a maximum on g = c, then either  $\nabla f = \lambda \nabla g$ , g = c holds or ...
- f) Which mathematician proved the switch the order of integration formula?
- g) True or false: the gradient vector  $\nabla f(x)$  is the same as df(x).

h) The equation  $u_t + uu_x = u_{xx}$  is an example of a differential equation. We have seen two major types (each a three capital letter acronym). Which type is it?

i) What is the formula for the arc length of a curve C?

j) What is the integration factor  $|d\phi|$  when going into polar coordinates?

#### Problem 28P.3 (10 points, 2 points for each sub-problem):

We see the level curves of a Morse function f. Only pick points A-J.

a) Which point is critical with discriminant  $D = \det(d^2 f) < 0$ .

b) At which point is  $f_x > 0, f_y = 0$ ?

c) At which point is  $f_x > 0, f_y > 0$ ?

d) Which  $(x_0, y_0)$  are critical points of f when imposing the constraint  $g(x, y) = y = y_0$ ?

e) Which  $(x_0, y_0)$  are critical points of f when imposing the constraint  $g(x, y) = x = x_0$ ?



#### Problem 28P.4 (10 points):

a) (5 points) Find the tangent plane to the surface  $xyz + x^5y + z = 11$  at (1, 2, 3). b) (5 points) Near (x, y) = (1, 2), we can write z = g(x, y). Find  $g_x(1, 2), g_y(1, 2)$ .

#### Problem 28P.5 (10 points):

a) Find the quadratic approximation of  $f(x, y, z) = 1 + x + y^2 + z^3 + \sin(xyz)$  at (0, 0, 0).

b) Estimate f(0.01, 0.03, 0.05) using linear approximation.

#### Problem 28P.6 (10 points):

a) (8 points) Classify the critical points of the function  $f(x, y) = x^{12} + 12x^2 + y^{12} + 12y^2$ using the second derivative test.

b) (2 points) Does f have a global minimum? Does f have a global maximum?

#### Problem 28P.7 (10 points):

On the top of a MIT building there is a radar dome in the form of a spherical cap. Insiders call it the "**Death star**" radar dome. We know that with the height h and base radius r, we have volume and surface area given by  $V = \pi r h^2 - \pi h^3/3$ ,  $A = 2\pi r h = \pi$ . This leads to the problem to extremize

$$f(x,y) = xy^2 - \frac{y^3}{3}$$

under the constraint

g(x,y) = 2xy = 1.

Find the minimum of f on this constraint using the Lagrange method!

Problem 28P.8 (10 points): Find  $\iint_R 5/(x^2 + y^2) \ dxdy \ ,$ 

where R is the region  $1 \le x^2 + y^2 \le 25$ ,  $y^2 > x^2$ .

Problem 28P.9 (10 points): Integrate f(x, y, z) = z over the solid *E* bound by

and

## $x + y + z = 1 \; .$

z = 0x = 0y = 0

#### Problem 28P.10 (10 points):

What is the surface area of the surface

$$r(x,y) = \begin{bmatrix} 2y\\ x\\ \frac{y^3}{3} + x \end{bmatrix}$$

with  $0 \le y \le 2$  and  $0 \le x \le y^3$ ?

OLIVER KNILL, KNILL@MATH.HARVARD.EDU, MATH 22A, HARVARD COLLEGE, FALL 2018