

REMINDER: INTEGRATION POLAR COORDINATES.

$$\int \int_R f(r, \theta) \boxed{\Gamma} d\theta dr .$$


EXAMPLE 1. Area of a disk of radius R

$$\int_0^R \int_0^{2\pi} r d\theta dr = 2\pi \frac{r^2}{2} \Big|_0^R = R^2\pi .$$

WHERE DOES THE FACTOR "r" COME FROM?


1. EXPLANATION. A small rectangle with dimensions $d\theta dr$ in the (r, θ) plane is mapped to a sector segment in the (x, y) plane. It has approximately the area $r d\theta dr$. It is small for small r .

2. EXPLANATION. The map $(r, \theta) \mapsto (r \cos(\theta), r \sin(\theta))$ which changes from Cartesian coordinates to polar coordinates. The **Jacobian** is defined as the matrix $T' = \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ r \sin(\theta) & r \cos(\theta) \end{bmatrix} = \begin{bmatrix} f_r & f_\theta \\ g_r & g_\theta \end{bmatrix}$. It contains the gradient vectors of $f(r, \theta)$ and $g(r, \theta)$ as rows. The determinant $f_r g_\theta - f_\theta g_r = r$ is the crossproduct of these two gradient vectors and gives the area of the parallelepiped.



CYLINDRICAL COORDINATES. Use polar coordinates in the x - y plane and leave the z coordinate. Take $T(r, \theta, z) = (r \cos(\theta), r \sin(\theta), z)$. The integration factor r is the same as in polar coordinates.

$$\int \int \int_{T(R)} f(x, y, z) dx dy dz = \int \int \int_R f(r, \theta, z) \boxed{\Gamma} dr d\theta dz$$



COORDINATES OF CAMBRIDGE. On the website <http://cello.cs.uiuc.edu/cgi-bin/slamm/ip2ll/> you can enter a host like *www.math.harvard.edu* and get latitude and longitude of the host: $(lat, lon) = (42.365, -71.1)$. Using (r, θ, ϕ) coordinates, we obtain the position $(r, 90 - 42.365, -71.1)$ of the host in spherical coordinates. The site does not give the height, but we are about on sea-level, so that $r = 6365 km$.

EXAMPLE. Calculate the volume bounded by the parabolic $z = 1 - (x^2 + y^2)$ and the $x - y$ plane. In cylindrical coordinates, the paraboloid is $z(r, \phi) = 1 - r^2$:



$$\int_0^1 \int_0^{2\pi} \int_0^{1-r^2} r dz d\phi dr = \int_0^1 \int_0^{2\pi} (r - r^3) d\phi dr = 2\pi (r^2/2 - r^4/4) \Big|_0^1 = \pi .$$

SPHERICAL COORDINATES. Spherical coordinates use the radius ρ as well as two angles: θ the polar angle and ϕ , the angle between the vector and the z axis. The coordinate change is


$$T : (x, y, z) = (\rho \cos(\theta) \sin(\phi), \rho \sin(\theta) \sin(\phi), \rho \cos(\phi)) .$$

The integration factor can be seen from the dimensions of a spherical wedge with dimensions $d\rho, \rho \sin(\phi) d\theta, \rho d\phi = \rho^2 \sin(\phi) d\theta d\phi d\rho$.

$$\int \int \int_{T(R)} f(x, y, z) dx dy dz = \int \int \int_R f(\rho, \theta, \phi) \boxed{\rho^2 \sin(\phi)} d\rho d\theta d\phi$$



$$\begin{aligned} T : (\rho, \theta, \phi) \\ \mapsto \\ (\rho \cos(\theta) \sin(\phi), \\ \rho \sin(\theta) \sin(\phi), \\ \rho \cos(\phi)) \end{aligned}$$



VOLUME OF SPHERE. A sphere of radius R has the volume

$$\int_0^R \int_0^{2\pi} \int_0^\pi \rho^2 \sin(\phi) d\phi d\theta d\rho .$$


The most inner integral $\int_0^\pi \rho^2 \sin(\phi) d\phi = -\rho^2 \cos(\phi)|_0^\pi = 2\rho^2$. The next layer is, because ϕ does not appear: $\int_0^{2\pi} 2\rho^2 d\phi = 4\pi\rho^2$. The final integral is $\int_0^R 4\pi\rho^2 d\rho = 4\pi R^3/3$.

MOMENT OF INERTIA. The moment of inertia of a body G with respect to an axis L is the triple integral $\int \int \int_G r(x, y, z)^2 dz dy dx$, where $r(x, y, z) = R \sin(\phi)$ is the distance from L . Problem: calculate the moment of inertia of a sphere of radius R with respect to the z -axis:

$$I = \int_0^R \int_0^{2\pi} \int_0^\pi \rho^2 \sin^2(\phi) \rho^2 \sin(\phi) d\phi d\theta d\rho = \left(\frac{1}{3} \sin^3(\phi)\right)_0^\pi \left(\int_0^R \rho^4 dr\right) \left(\int_0^{2\pi} d\theta\right) = \frac{4}{3} \cdot \frac{R^5}{5} \cdot 2\pi = \frac{8\pi R^5}{15} = \frac{VR^2}{5} .$$

If a sphere spins around the z -axis with angular velocity ω , then $I\omega^2/2$ is the kinetic energy of that sphere. Example: the moment of inertia of the earth is $810^{37} kgm^2$. The angular velocity is $\omega = 1/day = 1/(86400s)$ so that the energy of the earth rotation is $810^{37} kgm^2 / (7464960000s^2) \sim 10^{28} J = 10^{25} kJ \sim 2.510^{24} kcal$.

DIAMOND. Find the volume and the center of mass of a diamond, the intersection of the unit sphere with the cone given in cylindrical coordinates as $z = \sqrt{3}r$.



Solution: we use spherical coordinates to find the center of mass $(\bar{x}, \bar{y}, \bar{z})$:

$$\begin{aligned} V &= \int_0^1 \int_0^{2\pi} \int_0^{\pi/6} \rho^2 \sin(\phi) d\phi d\theta d\rho = \frac{(1-\frac{\sqrt{3}}{2})}{3} 2\pi \\ \bar{y} &= \int_0^1 \int_0^{2\pi} \int_0^{\pi/6} \rho^3 \sin^2(\phi) \sin(\theta) d\phi d\theta d\rho / V = 0 \\ \bar{x} &= \int_0^1 \int_0^{2\pi} \int_0^{\pi/6} \rho^3 \sin^2(\phi) \cos(\theta) d\phi d\theta d\rho / V = 0 \\ \bar{z} &= \int_0^1 \int_0^{2\pi} \int_0^{\pi/6} \rho^3 \cos(\phi) \sin(\phi) d\phi d\theta d\rho / V = \frac{2\pi}{32V} \end{aligned}$$