

## VECTORS, DOT PRODUCT


O. Knill, Maths21a

**VECTORS.** Two points  $P_1 = (x_1, y_1)$ ,  $Q = P_2 = (x_2, y_2)$  in the plane determine a **vector**  $\vec{v} = \langle x_2 - x_1, y_2 - y_1 \rangle$ . It points from  $P_1$  to  $P_2$  and we can write  $P_1 + \vec{v} = P_2$ .

**COMPONENTS.** Points  $P$  in space are in one to one correspondence to vectors pointing from 0 to  $P$ . The numbers  $\vec{v}_i$  in a vector  $\vec{v} = (v_1, v_2)$  are also called **components** or of the vector.

**REMARKS:** vectors can be drawn **everywhere** in the plane. If a vector starts at 0, then the vector  $\vec{v} = (v_1, v_2)$  points to the point  $(v_1, v_2)$ . That's is why one can identify points  $P = (a, b)$  with vectors  $\vec{v} = \langle a, b \rangle$ . Two vectors which can be translated into each other are considered **equal**. In three dimensions, vectors have three components. In some Encyclopedias like Encyclopedia Britannica define vectors as objects which have "both magnitude and direction". This is unprecise and strictly speaking incorrect because the **zero vector** is a vector with no direction.

### ADDITION SUBTRACTION, SCALAR MULTIPLICATION.



$$\vec{u} + \vec{v} = \langle u_1, u_2 \rangle + \langle v_1, v_2 \rangle = \langle u_1 + v_1, u_2 + v_2 \rangle$$

$$\vec{u} - \vec{v} = \langle u_1, u_2 \rangle - \langle v_1, v_2 \rangle = \langle u_1 - v_1, u_2 - v_2 \rangle$$

$$\lambda \vec{u} = \lambda \langle u_1, u_2 \rangle = \langle \lambda u_1, \lambda u_2 \rangle$$


**BASIS VECTORS.** The vectors  $\vec{i} = \langle 1, 0 \rangle$ ,  $\vec{j} = \langle 0, 1 \rangle$  are called **standard basis vectors** in the plane. In space, one has the basis vectors  $\vec{i} = \langle 1, 0, 0 \rangle$ ,  $\vec{j} = \langle 0, 1, 0 \rangle$ ,  $\vec{k} = \langle 0, 0, 1 \rangle$ . Every vector  $\vec{v} = (v_1, v_2)$  in the plane can be written as a sum of standard basis vectors:  $\vec{v} = v_1 \vec{i} + v_2 \vec{j}$ . Every vector  $\vec{v} = (v_1, v_2, v_3)$  in space can be written as  $\vec{v} = v_1 \vec{i} + v_2 \vec{j} + v_3 \vec{k}$ .

WHERE DO VECTORS OCCUR? Here are some examples:


**Velocity:** if  $(f(t), g(t))$  is a point in the plane which depends on time  $t$ , then  $\vec{v} = \langle f'(t), g'(t) \rangle$  is the **velocity vector** at the point  $(f(t), g(t))$ .




**Forces:** Some problems in statics involve the determination of a forces acting on objects. Forces are represented as vectors




**Fields:** electromagnetic or gravitational fields or velocity fields in fluids are described with vectors.



**Qbits:** in quantum computation, rather than working with bits, one deals with **qbits**, which are vectors.



**Color** can be written as a vector  $\vec{v} = (r, g, b)$ , where  $r$  is red,  $g$  is green and  $b$  is blue. An other coordinate system is  $\vec{v} = (c, m, y) = (1 - r, 1 - g, 1 - b)$ , where  $c$  is cyan,  $m$  is magenta and  $y$  is yellow.



**SVG.** Scalable Vector Graphics is an emerging standard for the web for describing two-dimensional graphics in XML.



**VECTOR OPERATIONS:** The addition and scalar multiplication of vectors satisfy "obvious" properties. There is no need to memorize them. We write  $*$  here for multiplication with a scalar but usually, the multiplication sign is left out.

$$\begin{aligned} \vec{u} + \vec{v} &= \vec{v} + \vec{u} \\ \vec{u} + (\vec{v} + \vec{w}) &= (\vec{u} + \vec{v}) + \vec{w} \\ \vec{u} + \vec{0} &= \vec{0} + \vec{u} = \vec{u} \\ r * (s * \vec{v}) &= (r * s) * \vec{v} \\ (r + s) \vec{v} &= \vec{v} (r + s) \\ r(\vec{v} + \vec{w}) &= r \vec{v} + r \vec{w} \\ 1 * \vec{v} &= \vec{v} \end{aligned}$$

commutativity  
additive associativity  
null vector  
scalar associativity  
distributivity in scalar  
distributivity in vector  
the one element

**LENGTH.** The length  $|\vec{v}|$  of  $\vec{v}$  is the distance from the beginning to the end of the vector.

**EXAMPLES.** 1) If  $\vec{v} = (3, 4)$ , then  $|\vec{v}| = \sqrt{25} = 5$ . 2)  $|\vec{i}| = |\vec{j}| = |\vec{k}| = 1$ ,  $|\vec{0}| = 0$ .

**UNIT VECTOR.** A vector of length 1 is called a **unit vector**. If  $\vec{v} \neq \vec{0}$ , then  $\vec{v}/|\vec{v}|$  is a unit vector.

**EXAMPLE:** If  $\vec{v} = (3, 4)$ , then  $\vec{v} = (2/5, 3/5)$  is a unit vector,  $\vec{i}, \vec{j}, \vec{k}$  are unit vectors.

**PARALLEL VECTORS.** Two vectors  $\vec{v}$  and  $\vec{w}$  are called **parallel**, if  $\vec{v} = r \vec{w}$  with some constant  $r$ .

**DOT PRODUCT.** The **dot product** of two vectors  $\vec{v} = (v_1, v_2, v_3)$  and  $\vec{w} = (w_1, w_2, w_3)$  is defined as

$$\vec{v} \cdot \vec{w} = v_1 w_1 + v_2 w_2 + v_3 w_3$$

Remark: in science, other notations are used:  $\vec{v} \cdot \vec{w} = (\vec{v}, \vec{w})$  (mathematics)  $\langle \vec{v} | \vec{w} \rangle$  (quantum mechanics)  $v_i w^i$  (Einstein notation)  $g_{ij} v^i w^j$  (general relativity). The dot product is also called **scalar product**, or **inner product**.


**LENGTH.** Using the dot product one can express the length of  $\vec{v}$  as  $|\vec{v}| = \sqrt{\vec{v} \cdot \vec{v}}$ .

**CHALLENGE.** Express the dot product in terms of the length alone.

**SOLUTION:**  $(\vec{v} + \vec{w}, \vec{v} + \vec{w}) = (\vec{v}, \vec{v}) + (\vec{w}, \vec{w}) + 2(\vec{v}, \vec{w})$  can be solved for  $(\vec{v}, \vec{w})$ .

**ANGLE.** Because  $|\vec{v} - \vec{w}|^2 = (\vec{v} - \vec{w}, \vec{v} - \vec{w}) = |\vec{v}|^2 + |\vec{w}|^2 - 2(\vec{v}, \vec{w})$  is by the **cos-theorem** equal to  $|\vec{v}|^2 + |\vec{w}|^2 - 2|\vec{v}| \cdot |\vec{w}| \cos(\alpha)$ , where  $\alpha$  is the angle between the vectors  $\vec{v}$  and  $\vec{w}$ , we get the important formula

$$\vec{v} \cdot \vec{w} = |\vec{v}| \cdot |\vec{w}| \cos(\alpha)$$



**CAUCHY-SCHWARZ INEQUALITY:**  $|\vec{v} \cdot \vec{w}| \leq |\vec{v}| |\vec{w}|$  follows from that formula because  $|\cos(\alpha)| \leq 1$ .

**TRIANGLE INEQUALITY:**  $|\vec{u} + \vec{v}| \leq |\vec{u}| + |\vec{v}|$  follows from  $|\vec{u} + \vec{v}|^2 = (\vec{u} + \vec{v}) \cdot (\vec{u} + \vec{v}) = \vec{u}^2 + \vec{v}^2 + 2\vec{u} \cdot \vec{v} \leq \vec{u}^2 + \vec{v}^2 + 2|\vec{u}| \cdot |\vec{v}| \leq (|\vec{u}| + |\vec{v}|)^2$ .


**FINDING ANGLES BETWEEN VECTORS.** Find the angle between the vectors  $(1, 4, 3)$  and  $(-1, 2, 3)$ .

**ANSWER:**  $\cos(\alpha) = 16/(\sqrt{26}\sqrt{14}) \sim 0.839$ . So that  $\alpha = \arccos(0.839..) \sim 33^\circ$ .

**ORTHOGONAL VECTORS.** Two vectors are called **orthogonal** if  $\vec{v} \cdot \vec{w} = 0$ . The zero vector  $\vec{0}$  is orthogonal to any vector. **EXAMPLE:**  $\vec{v} = (2, 3)$  is orthogonal to  $\vec{w} = (-3, 2)$ .

**PROJECTION.** The vector  $\vec{a} = \text{proj}_{\vec{w}}(\vec{v}) = \vec{w}(\vec{v} \cdot \vec{w} / |\vec{w}|^2)$  is called the **projection** of  $\vec{v}$  onto  $\vec{w}$ .

The **scalar projection** is defined as  $\text{comp}_{\vec{w}}(\vec{v}) = (\vec{v} \cdot \vec{w}) / |\vec{w}|$ . (Its absolute value is the length of the projection of  $\vec{v}$  onto  $\vec{w}$ .) The vector  $\vec{b} = \vec{v} - \vec{a}$  is called the **component** of  $\vec{v}$  orthogonal to the  $\vec{w}$ -direction.



**EXAMPLE.**  $\vec{v} = (0, -1, 1)$ ,  $\vec{w} = (1, -1, 0)$ ,  $\text{proj}_{\vec{w}}(\vec{v}) = (1/2, -1/2, 0)$ ,  $\text{comp}_{\vec{w}}(\vec{v}) = 1/\sqrt{2}$ .