

### 3D INTEGRATION

Maths21a, O. Knill

3D INTEGRATION. If  $f(x, y, z)$  is a function of three variables and  $E$  is a solid in space, then  $\iint \int_E f(x, y, z) dx dy dz$  is defined as the limit of Riemann sums  $\frac{1}{n^3} \sum_{(x_i, y_j, z_k) \in R} f(x_i, y_j, z_k)$  for  $n \rightarrow \infty$ , where  $(x_i, y_j, z_k) = (\frac{i}{n}, \frac{j}{n}, \frac{k}{n})$ .

TRIPLE INTEGRALS. As in two dimensions, triple integrals can be evaluated by solving successively one dimensional integrals.

EXAMPLE. Assume  $R$  is the box  $[0, 1] \times [0, 1] \times [0, 1]$  and  $f(x, y, z) = 24x^2y^3z$ .

$$\int_0^1 \int_0^1 \int_0^1 24x^2y^3z dx dy dz$$

CALCULATION. We start from the core  $\int_0^1 24x^2y^3z dz = 12x^3y^3$ , then integrate the middle layer:

$$\int_0^1 12x^3y^3 dy = 3x^2 \text{ and finally handle the outer layer: } \int_0^1 3x^2 dx = 1.$$

WHAT DID WE DO? When we calculate the most inner integral, we fix  $z$  and  $y$ . The integral is the average of  $f(x, y, z)$  along a line intersected with the body. After completing the second integral, we have computed the average on the plane  $z = \text{const}$  intersected with  $R$ . The most outer integral averages all these two dimensional sections.

VOLUME UNDER GRAPH. The volume under the graph of a function  $f(x, y)$  and above a region  $R = [a, b] \times [c, d]$  is the integral  $\int_a^b \int_c^d f(x, y) dx dy$ . Actually, this is a triple integral:

$$V = \int_a^b \int_c^d \int_0^{f(x,y)} 1 dz dx dy$$

An integral of the form  $\int_a^b \int_c^d \int_{g_1(x,y)}^{g_2(x,y)} f(x, y, z) dz dx dy$  is sometimes also called a **type I** triple integral.

VOLUME OF THE SPHERE. (We will do this more elegantly later). The volume is

$$V = \iiint_R dx dy dz = \int_{-1}^1 \left[ \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \left[ \int_{-\sqrt{1-x^2-y^2}}^{\sqrt{1-x^2-y^2}} dz \right] dy \right] dx$$


After computing the inner integral, we have  $V = 2 \int_{-1}^1 \left[ \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} (1-x^2-y^2)^{1/2} dy \right] dx$ .

To resolve the next layer, call  $1-y^2 = a^2$ . The task is to find  $\int_{-a}^a \sqrt{a^2-y^2} dy$ . Make the substitution  $y/a = \sin(u)$ ,  $dy = a \cos(u)$  to write this as  $a \int_0^{\arcsin(a/a)} \sqrt{1-\sin^2(u)} a \cos(u) du = a^2 \int_0^{\pi/2} \cos^2(u) du = a^2 \pi/2$ . We can finish up the last integral

$$V = 2\pi/2 \int_{-1}^1 (1-x^2) dx = 4\pi/3$$

MASS OF A BODY. In general, the mass of a body  $E$  with density  $\rho(x, y, z)$  is  $\iiint_E \rho(x, y, z) dV$ . For bodies with constant density  $\rho$  the mass is  $\rho V$ , where  $V$  is the volume. Compute the mass of a body which is bounded by the parabolic cylinder  $z = 4 - x^2$ , and the planes  $x = 0, y = 0, y = 6, z = 0$  if the density of the body is 1.

$$\begin{aligned} \int_0^2 \int_0^6 \int_0^{4-x^2} dz dy dx &= \int_0^2 \int_0^6 (4-x^2) dy dx \\ &= 6 \int_0^2 (4-x^2) dx = 6(4x - x^3/3)|_0^2 = 32 \end{aligned}$$



CENTER OF MASS. Compute the center of mass of the same body. The center of mass is  $(24/32, 96/32, 256/180) = (3/4, 3, 8/5)$ :

$$\begin{aligned} \int_0^2 \int_0^6 \int_0^{4-x^2} x dz dy dx &= \int_0^2 \int_0^6 x(4-x^2) dy dx = 6 \int_0^2 x(4-x^2) dx = 24x^2/2 - 6x^4/4|_0^2 = 24 \\ \int_0^2 \int_0^6 \int_0^{4-x^2} y dz dy dx &= \int_0^2 \int_0^6 y(4-x^2) dy dx = \int_0^2 18(4-x^2) dx = 18(4x - x^3/3)|_0^2 = 96 \\ \int_0^2 \int_0^6 \int_0^{4-x^2} z dz dy dx &= \int_0^2 \int_0^6 (4-x^2)^2/2 dy dx = 6 \int_0^2 (4-x^2)^2/2 dx = 3(16x - 8x^3/3 + x^5/5)|_0^2 = 256/5 \end{aligned}$$

SOME HISTORY OF COMPUTING VOLUMES. How did people come up calculating the volume  $\iiint_R 1 dx dy dz$  of a body?



**Archimedes ((-287)-(-212)):** Archimedes's method of integration allowed him to find areas, volumes and surface areas in many cases. His method of exhaustion paths the numerical method of integration by Riemann sum. The **Archimedes principle** states that any body submerged in a water is acted upon by an upward force which is equal to the weight of the displaced water. This provides a practical way to compute volumes of complicated bodies.



**Cavalieri (1598-1647):** Cavalieri could determine area and volume using tricks like the **Cavalieri principle**. Example: to get the volume the half sphere of radius  $R$ , cut away a cone of height and radius  $R$  from a cylinder of height  $R$  and radius  $R$ . At height  $z$ , this body has a cross section with area  $R^2\pi - r^2\pi$ . If we cut the half sphere at height  $z$ , we obtain a disc of area  $(R^2 - r^2)\pi$ . Because these areas are the same, the volume of the half-sphere is the same as the cylinder minus the cone:  $\pi R^3 - \pi R^3/3 = 2\pi R^3/3$  and the volume of the sphere is  $4\pi R^3/3$ .



**Newton (1643-1727) and Leibniz (1646-1716):** Newton and Leibniz, developed calculus independently. The new tool made it possible to compute integrals through "anti-derivation". Suddenly, it became possible to find integrals using analytic tools.

MONTE CARLO COMPUTATIONS. Here is an other way to compute integrals: Suppose we want to calculate the volume of some body  $R$  inside the unit cube  $[0, 1] \times [0, 1] \times [0, 1]$ . The **Monte Carlo method** is to shoot randomly  $n$  times onto the unit cube and count the fraction of times, we hit the solid. Here is an experiment with Mathematica and where the body is one eights of the unit ball:

```
R := Random[]; k = 0; Do[x = R; y = R; z = R; If[x^2 + y^2 + z^2 < 1, k + 1], {10000}]; k/10000
```

Assume, we hit 5277 of  $n=10000$  times. The volume so measured is 0.5277. The actual volume of  $1/8$ 'th of the sphere is  $\pi/6 = 0.524$ . For  $n \rightarrow \infty$  the Monte Carlo computation gives the actual volume.

WHERE CAN TRIPLE INTEGRALS OCCUR?

- Calculation of **volumes**  $V = \iiint_R 1 dV$  and **masses**  $M = \iiint_R \rho dV$ .
- Finding **averages**  $\iiint_R f dV / \iiint_R 1 dV$ . Examples: average algae concentration in a swimming pool.
- Determining **probabilities**. Example: quantum probability  $\iiint_R f(x, y, z)^2 dx dy dz$ .
- **Moment of inertia**  $\iiint_R r(x, y, z)^2 \rho(x, y, z) dV$ , where  $r(x, y, z)$  is the distance to the axis of rotation.
- **Center of mass**  $(\iiint_R x \rho dV / M, \iiint_R y \rho dV / M, \iiint_R z \rho dV / M)$ .