

Name:
-------

- Start by printing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work.
- Do not detach pages from this exam packet or unstaple the packet.
- Please try to write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids are allowed.
- Problems 1-3 do not require any justifications. For the rest of the problems you have to show your work. Even correct answers without derivation can not be given credit.
- You have 180 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
14		10
Total:		150

Problem 1) (20 points)

- 1)  T  F The quadratic surface  $x^2 + y - z^2 = -5$  is a hyperbolic paraboloid.

**Solution:**

Write it as  $y - 5 = z^2 - x^2$ , to see it better.

- 2)  T  F There are vectors  $\vec{u}$  and  $\vec{v}$  such that  $|\vec{u} \times \vec{v}| > |\vec{u}||\vec{v}|$ .

**Solution:**

We have a general identity  $|\vec{u} \times \vec{v}| = |\vec{u}||\vec{v}| \sin(\alpha)$  which contradicts the claim.

- 3)  T  F  $\int_0^{2\pi} \int_0^5 r \, d\theta \, dr$  is the area of a disc of radius 5.

**Solution:**

There seems nothing wrong with that. But note that the 0 to 5 integration is paired with the  $d\theta$ .

- 4)  T  F If a vector field  $\vec{F}(x, y)$  satisfies  $\text{curl}(\vec{F})(x, y) = Q_x - P_y = 0$  for all points  $(x, y)$  in the plane, then  $\vec{F}$  is a gradient field.

**Solution:**

True. We have derived this from Green's theorem.

- 5)  T  F The jerk of a parameterized curve  $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$  is parallel to the acceleration if the curve  $\vec{r}(t)$  is a line.

**Solution:**

The velocity, the acceleration and the jerk are all parallel on a line.

- 6)  T  F The curvature of the curve  $\vec{r}(t) = \langle 3 \sin(t), 0, 3 \cos(t) \rangle$  is twice the curvature of the curve  $\vec{s}(t) = \langle 6 + 6 \sin(t), 6 \cos(t), 0 \rangle$ .

**Solution:**

If we scale a curve by a factor 2, its curvature is divided by 2.

- 7)  T  F The curve  $\vec{r}(t) = \langle \sin(t), t^2, \cos(t) \rangle$  for  $t \in [0, 10\pi]$  is located on a cylinder.

**Solution:**

Indeed, one can check that  $x(t)^2 + z(t)^2 = 1$ .

- 8)  T  F If a function  $f(x, y)$  has the property that  $f_x(x, y)$  is zero for all  $x, y$ , then  $f$  is the constant function.

**Solution:**

No, for example  $f(x, y) = y$  is also a solution too and this solution is not constant.

- 9)  T  F If the unit tangent vector  $\vec{T}(t)$  of a curve  $\vec{r}(t)$  is always parallel to a plane  $\Sigma$ , then the curve is contained in a plane parallel to  $\Sigma$ .

**Solution:**

Indeed, we never leave the plane which goes through the initial point  $r(0)$  because also  $r'(t)$  is always parallel to  $\Sigma$  and after integration, the curve  $r(t)$  has to be in the plane.

- 10)  T  F If  $(x_0, y_0)$  is an extremum of  $f(x, y)$  under the constraint  $x^2 + y^2 = 1$ , then the same point is an extremum of  $10f(x, y)$  under the same constraint.

**Solution:**

The point is a solution to the same Lagrange equations.

- 11)  T  F At a critical point  $(x_0, y_0)$  of a function  $f(x, y)$  for which  $f_{xx}(x_0, y_0) > 0$ , the critical point is always a minimum.

**Solution:**

No, we also need  $D > 0$ .

- 12)  T  F If a vector field  $\vec{F}(x, y)$  is a gradient field, and  $C$  is a closed curve which looks like a figure 8, then  $\int_C \vec{F} \cdot d\vec{r}$  is zero.

**Solution:**

This follows from the fundamental theorem of line integrals.

- 13)  T  F If  $C$  is part of a level curve of a function  $f(x, y)$  and  $\vec{F} = \langle f_x, f_y \rangle$  is the gradient field of  $f$ , then  $\int_C \vec{F} \cdot d\vec{r} = 0$ .

**Solution:**

The gradient field is perpendicular to the level curves.

- 14)  T  F The divergence of the gradient vector field  $\vec{F}(x, y, z) = \nabla f(x, y, z)$  is always the zero function.

**Solution:**

The divergence of the gradient of  $f$  is the Laplacian of  $f$

- 15)  T  F The line integral of the vector field  $\vec{F}(x, y, z) = \langle x, y, z \rangle$  along a line segment from  $(0, 0, 0)$  to  $(1, 1, 1)$  is  $3/2$ .

**Solution:**

By the fundamental theorem of line integrals, we can take the difference of the potential  $f(x, y, z) = x^2/2 + y^2/2 + z^2/2$ , which is  $1/2 + 1/2 + 1/2 = 3/2$ .

- 16)  T  F The area of a region  $G$  can be expressed as a line integral along its boundary.

**Solution:**

This is a consequence of Green's theorem and we have seen a few examples.

- 17)  T  F The flux of the vector field  $\vec{F}(x, y, z) = \langle x, y, -z \rangle$  through the boundary  $S$  of a solid ellipsoid  $E$  is equal to the volume the ellipsoid.

**Solution:**

Indeed the divergence of the field is 1 and we can apply the divergence theorem.

- 18)  T  F If  $\vec{F}$  is a vector field in space and  $S$  is a torus surface, then the flux of  $\text{curl}(\vec{F})$  through  $S$  is 0.

**Solution:**

This is true by Stokes theorem.

- 19)  T  F If the divergence and the curl of a vector field  $\vec{F}$  are both zero, then it is a constant field.

**Solution:**

Take  $\vec{F}(x, y, z) = \langle x, -y, 0 \rangle$ . It has zero curl and zero divergence but is not constant.


- 20)  T  F For any function  $f$ , the curl of  $\vec{F} = \text{grad}(f)$  is the zero field  $\langle 0, 0, 0 \rangle$ .

**Solution:**

$\text{curl}(\text{grad}(f)) = \langle 0, 0, 0 \rangle$  is an important identity.


Problem 2) (10 points)

- a) (4 points) Match the regions with the corresponding double integrals




Enter a,b,c,d	Function
	$\int_0^1 \int_{x/2}^x f(x, y) dy dx$
	$\int_0^1 \int_0^y f(x, y) dx dy$
	$\int_0^1 \int_0^{x/2} f(x, y) dy dx$
	$\int_0^1 \int_{y/2}^1 f(x, y) dx dy$


- b) (6 points) Match the parametrized or implicit surfaces with their definitions




A




B




C



D



E



F

Enter A-F here	Function or parametrization
	$\vec{r}(u, v) = \langle \cos(u), \sin(u), v \rangle$
	$\vec{r}(u, v) = \langle u - v, u + 2v, 2u + 3v \rangle$
	$x^2 + y^2/3 + z^2/3 = 1$
	$\vec{r}(u, v) = \langle (\sin(v) + 1) \cos(u), (\sin(v) + 1) \sin(u), v \rangle$
	$z - x + \sin(xy) = 0$
	$x^2 + y^2 - z^2 = 0$



**Solution:**


a) a c d b

b) C E A D B F

Problem 3) (10 points)

a) (4 points) Match the vector fields and curves with the corresponding line integral





Enter I,II,III,IV	Line integral
	$\int_0^{2\pi} \langle \cos(t), \sin(t) \rangle \cdot \langle -\sin(t), \cos(t) \rangle dt$
	$\int_0^{2\pi} \langle -t, t^2 \rangle \cdot \langle 1, 1 \rangle dt$
	$\int_0^{2\pi} \langle t^2, t \rangle \cdot \langle 1, 2t \rangle dt$
	$\int_0^{2\pi} \langle -3 \sin(t), 3 \cos(t) \rangle \cdot \langle -\sin(t), \cos(t) \rangle dt$

b) (6 points) Fill in from following choice: "arc length", "surface area", "chain rule", "volume of parallelepiped", "area of parallelogram", "line integral", "flux integral", "curvature".

Formula	Name of formula or rule or theorem
$\int \int_R  \vec{r}_u \times \vec{r}_v  dudv$	
$\frac{d}{dt} f(\vec{r}(t)) = \nabla f(\vec{r}(t)) \cdot \vec{r}'(t)$	
$\int_a^b  \vec{r}'(t)  dt$	
$\frac{ \vec{r}'(t) \times \vec{r}''(t) }{ \vec{r}'(t) ^3}$	
$ \vec{u} \cdot (\vec{v} \times \vec{w}) $	
$\int_0^1 \int_0^1 \vec{F}(\vec{r}(u, v)) \cdot (\vec{r}_u \times \vec{r}_v) dudv$	

**Solution:**

a) IV, I, II, III

b) surface area, chain rule, arc length, curvature, volume of parallelepiped, flux integral.

Problem 4) (10 points)

Given the line  $x - 1 = y - 2 = z - 3$  and the point  $P = (8, 4, 5)$ . Find the equation

$$ax + by + cz = d$$

of the plane which contains the line and the point.

**Solution:**

From the symmetric equation, we see that the line contains the vector  $\langle 1, 1, 1 \rangle$ . The line also contains the point  $(1, 2, 3)$  so that the vector  $\langle 7, 2, 2 \rangle$  is also in the plane. The cross product  $\vec{n} = \langle 1, 1, 1 \rangle \times \langle 7, 2, 2 \rangle$  is  $\langle 0, 5, -5 \rangle$  so that the equation of the plane is  $5y - 5z = d$ . The constant  $d$  can be obtained by plugging in a point in the plane, for example  $(1, 2, 3)$  so that  $d = -5$ . The equation of the plane is  $5y - 5z = -5$  which can also be written as  $z - y = 1$ .

Problem 5) (10 points)

Find all the critical points of the function  $f(x, y) = y^3 - 3y^2 + 4x + x^2 - 3$  and classify them by telling whether they are local maxima, local minima or saddle points.

**Solution:**

The critical points are  $P = (-2, 0)$  and  $Q = (-2, 2)$ . The discriminant  $D$  at  $P$  is  $D = -12$  so that  $P$  is a saddle point. The Hessian at  $Q$  is 12 and  $f_{xx} = 2$  which is a local minimum.

$P = (-2, 0)$	$D = -12$	irrelevant	saddle point
$Q = (-2, 2)$	$D = 12$	$f_{xx} = 2$	local minimum

Problem 6) (10 points)



The hyperbolic paraboloid  $x^2 - y^2 - 3z = 0$  contains the point  $P = (1, 1, 0)$  and the point  $Q = (3, 0, 3)$ . Find the tangent planes to the surface at  $P$  and  $Q$  and find a parametrization  $\vec{r}(t)$  of the line of intersection of these two planes.

**Solution:**

The gradient of  $g(x, y, z) = x^2 - y^2 - z = 0$  is  $(2x, -2y, -3)$  which is  $(2, -2, -3)$  at the first point and  $(6, 0, -3)$  at the second point. The first tangent plane equation is  $2x - 2y - 3z = 3$ . The second tangent plane equation is  $6x - 3z = 9$ . To get the line of intersection, we find a point in the intersection, like  $(3, 0, 3)$  and the cross product  $\langle 2, -2, -3 \rangle \times \langle 6, 0, -3 \rangle = \langle 6, -12, 12 \rangle$  of the normal vectors to get a direction in the line.  $\vec{r}(t) = (3, 0, 3) + t\langle 1, -1, 2 \rangle$ . Note that there are many solutions since the initial position on the line and the length of the vector in the line can change.

Problem 7) (10 points)

A water reservoir in Burlington, MA (the map to the right is centered there) is bounded by a solid cylinder  $x^2 + y^2 \leq 1$ . It has as the roof the cone  $x^2 + y^2 = (z - 6)^2$  and is bounded from below by the  $xy$ -plane  $z = 0$ . What is the volume of the reservoir?



**Solution:**

If  $R$  denotes the unit disc, we have the triple integral

$$\int \int_R \int_0^{6+\sqrt{x^2+y^2}} 1 \, dz dy dx$$

which of course is integrated best in cylindrical coordinates:

$$\int_0^1 \int_0^{2\pi} \int_0^{6-r} r \, dz \, d\theta \, dr$$

which is  $2\pi \int_0^1 6r - r^2 \, dr = 2\pi(3 - 1/3) = 20\pi/3$ .

Problem 8) (10 points)

Find the maxima and minima of the function  $f(x, y) = x^2 - y^2$  on the parabola  $x + y^2 = 1$  using the Lagrange multiplier method.

**Solution:**

The Lagrange equations are

$$\begin{aligned}2x &= \lambda \\ -2y &= 2\lambda y \\ x + y^2 &= 1\end{aligned}$$

The second equation can be solved by  $y = 0$ , a case which was often forgotten. In that case the third equation gives  $x = 1$ .

If  $y$  is not zero, we can divide the second equation by  $y$ . Dividing now the second by the first equation gives  $-1/x = 2$  so that  $x = -1/2$ . Plugging into the third gives  $y = \pm\sqrt{3/2}$ . The solutions are  $(1, 0), (-1/2, -\sqrt{3/2}), (-1/2, \sqrt{3/2})$ . By plugging in  $f$  values, we see that the first is a maximum, the other two points are minima. As usual with Lagrange methods we can not use the second derivative test to check for maxima and minima. In summary:

$(1, 0)$  is a maximum, and  $(-1/2, \sqrt{3/2})$  as well as  $(-1/2, -\sqrt{3/2})$  are minima.

Problem 9) (10 points)

Compute the surface area of the surface  $\vec{r}(u, v) = \langle u^3, v^3, u^3 - v^3 \rangle$  parametrized so that  $(u, v)$  is in the unit disc.

**Solution:**

$\vec{r}_u \times \vec{r}_v = \langle -9u^2v^2, 9u^2v^2, 9u^2v^2 \rangle$  with length  $|\vec{r}_u \times \vec{r}_v| = \sqrt{39}u^2v^2$ . Integrating this over the unit disc  $R$  leads to a double integral problem

$$9\sqrt{3} \int \int_R u^2v^2 \, dudv .$$

which is best solved in polar coordinates  $u = r \cos(\theta), v = r \sin(\theta)$ . Using  $2^2 \sin^2(\theta)^2 \cos^2(\theta)^2 = \sin^2(2\theta)^2$  we get

$$9\sqrt{3} \int_0^{2\pi} \int_0^1 r^5 (1/4) \sin^2(2\theta) \, d\theta dr$$

Using  $\int_0^{2\pi} \sin^2(2\theta) \, d\theta = \pi$  we end up with  $\sqrt{33}\pi/8$ .

Problem 10) (10 points)

Evaluate the following double integral

$$\int_0^2 \int_{x/2}^1 \cos(y^2) \, dy \, dx .$$

**Solution:**

The integral can not be evaluated as given. Change the order of integration

$$\int_0^1 \int_0^{2y} \cos(y^2) \, dx dy$$

This can now be solved. After evaluating the trivial inner integral we get

$$\int_0^1 2y \cos(y^2) \, dy = \sin(y^2)_0^1 = \sin(1) .$$

The final answer is  $\sin(1)$ .

Problem 11) (10 points)

Find the value of the line integral

$$\int_C \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) \, dt ,$$

where  $\vec{F}(x, y) = \langle y + \sin(\cos(x)), -2x \rangle$  and  $C$  is the boundary of the unit circle traversed in the counter clockwise direction.

**Solution:**


By Greens theorem the line integral is the double integral of  $\text{curl}(\vec{F}) = Q_x - P_y = -3$  over the unit disc, which is  $-3$  times the area  $\pi$  of the disc. The final result is  $\boxed{-6\pi}$ .

Problem 12) (10 points)

Find the value of the flux integral

$$\int \int_S \text{curl}(\vec{F})(\vec{r}(u, v)) \cdot \vec{r}_u \times \vec{r}_v \, dudv$$

where  $\vec{F}(x, y, z) = \langle -y, x, z \rangle$  and  $S$  is the part of the two-sheeted hyperboloid  $x^2 + y^2 - z^2 = -1$  which satisfies  $1 < z < 2$  and which is oriented so that the normal vector points downwards on  $S$ .



**Solution:**


By Stokes theorem, we can compute the line integral of  $\vec{F}$  along the boundary of the surface  $S$  instead. This boundary  $C$  is parametrized by  $\vec{r}(t) = \langle \sqrt{3} \cos(t), \sqrt{3} \sin(t), 2 \rangle$ . The line integral  $\int_0^{2\pi} \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) \, dt$  which is  $\boxed{6\pi}$ .

Problem 13) (10 points)

Let  $E$  be the solid which is bounded on the side by the cone  $S_1 : x^2 + y^2 = z^2, 0 < z < 1$  and on top by the disc  $S_2 = x^2 + y^2 \leq 1, z = 1$ . Let  $\vec{F}(x, y, z) = \langle 1 + 4x, 2 - 5y, 3 + 2z \rangle$ . Find the value of the flux integral

$$\int_S \vec{F}(\vec{r}(u, v)) \cdot \vec{r}_u \times \vec{r}_v \, dudv ,$$

where  $S$  is the union of the two surfaces  $S_1$  and  $S_2$ . The normal vector of  $S$  is oriented outwards on  $S_1 \cup S_2$ .




**Solution:**

By the divergence theorem, the flux is  $\iiint_E 1 \, dx dy dz$ , where  $E$  is the cone. The divergence of  $\vec{F}$  is constant 1 so that we have to integrate 1 over  $E$  which is the volume of the cone. This is best computed in cylindrical coordinates:  $\int_0^{2\pi} \int_0^1 \int_0^z r \, dr dz d\theta =$  which is  $\boxed{\pi/3}$ .

Problem 14) (10 points)

Find the volume of the solid piece of **cheese** bound by the cylinder  $x^2 + y^2 = 1$ , the planes  $y - z = 0$  (bottom boundary) and  $y + z = 0$  (top boundary) which is on the quadrant  $x \geq 0$  and  $y \leq 0$ .



**Solution:**

We use cylindrical coordinates. The base region in the  $xy$  plane is the fourth quadrant. Its roof is  $z = -y$ , its floor is  $z = y$ . We have to integrate  $f(x, y) = -2y = -2r \sin(\theta)$  over the fourth quadrant and get:

$$\int_0^1 \int_{3\pi/2}^{2\pi} -2r \sin(\theta) r \, d\theta dr = 2/3 .$$

The volume of the cheese is  $\boxed{2/3}$ .