

"I affirm my awareness of the standards of the Harvard College Honor Code."

Name:

Please email the PDF as an email attachment to knill@math.harvard.edu. The file needs to have your name capitalized like OliverKnill.pdf. Use **your personal handwriting**, no typing. No books, calculators, computers, or other electronic aids are allowed. (You can use a tablet to write). You can consult with a single page of your own handwritten notes, when writing the exam. The exam needs to arrive on Friday, July 9 at 10 AM. Write clearly and always give details of your computations. If you use separate paper, sign it with the honor code statement, use a page for each problem and copy the structure of the **check boxes**. All your final answers need to be in the boxes.

Problem 1) (20 points) No justifications are needed.

- 1) ☐ T ☐ F The plane $x = 3$ does intersect the yz -plane.
- 2) ☐ T ☐ F The curve $\vec{r}(t) = [1 + 2t, t, 1 + t]$ intersects the z -axis in a point.
- 3) ☐ T ☐ F The Cauchy-Schwartz inequality states $|\vec{v} \cdot \vec{w}| \leq |\vec{v}|$ for any two vectors $\vec{v}, \vec{w}$.
- 4) ☐ T ☐ F The curvature of a circle $\vec{r}(t) = [\cos(2t), 0, \sin(2t)]$ is equal to $1/2$ everywhere.
- 5) ☐ T ☐ F The surface $y^2 - x + y^2 = 2$ is an elliptic paraboloid.
- 6) ☐ T ☐ F The angle between the vectors $0\vec{A}$ and $0\vec{B}$ is positive if the distance between the points A and B is positive.
- 7) ☐ T ☐ F Let $\vec{j} = [0, 1, 0]$. There is a vector $\vec{v}$ for which the vector projection of $\vec{v}$ onto $\vec{j}$ is equal to $-\vec{j}$.
- 8) ☐ T ☐ F Two particles with path $\vec{r}_1(t) = [0, t, -t]$ and $\vec{r}_2(t) = [1 - t, t - 1, 0]$ do collide.
- 9) ☐ T ☐ F In spherical coordinates the surface $\rho^2 \sin^2(\phi) - \rho^2 \cos^2(\phi) = 1$ is a one-sheeted hyperboloid.
- 10) ☐ T ☐ F If $|\vec{u} \times \vec{v}| = 1$, for unit vectors $\vec{u}, \vec{v}$, then $\vec{u}, \vec{v}$ are orthogonal.
- 11) ☐ T ☐ F The curve $r^2 \cos^2(\theta) - r^2 \sin^2(\theta) = 1$ in polar coordinates is a hyperbola.
- 12) ☐ T ☐ F If the arc length of a curve connecting A with B is 0, then $A = B$.
- 13) ☐ T ☐ F The surface parametrized as $\vec{r}(y, z) = [y, z, y^2 - z^2]$ is a hyperbolic paraboloid.
- 14) ☐ T ☐ F The velocity vector and the acceleration are always either parallel or perpendicular.
- 15) ☐ T ☐ F It is possible that a plane and a one-sheeted hyperboloid intersects in two crossing lines.
- 16) ☐ T ☐ F The function $f(x, y) = \log(x^2 + y^2)$ contains as domain all points except the origin $(0, 0)$.


- 17) ☐ T ☐ F The normal vector $\vec{N}$ and the unit tangent vector $\vec{T}$ are perpendicular if $\vec{T}, \vec{T}'$ are both not zero vectors.
- 18) ☐ T ☐ F The distance between two non-parallel lines in three dimensional space can be zero.
- 19) ☐ T ☐ F If $\vec{i}, \vec{j}, \vec{k}$ denote the unit vectors in the x, y and z axis, then $\vec{i} \cdot (\vec{j} \times \vec{k}) = 1$.
- 20) ☐ T ☐ F For any two lines L, M , there are points P on L and Q on M such that $d(P, Q) = 2d(L, M)$, where $d(L, M)$ is the distance between the lines.

Problem 2) (10 points) No justifications are needed in this problem.


In each sub-problem, each of the numbers 0,1,2,3 each occur exactly once.

a) (2 points) Match the surfaces $g(x, y, z) = c$. Enter 0 if there is no match.


1



2




3




Function $g(x, y, z) =$	0,1,2, or 3
$x - y^2 + z = 0$	
$x^2 + y^2 - z^4 = 1$	
$z - e^{-x^2-y^2} = 0$	
$y^2 + z^3 = 1$	

b) (2 points) Match the graphs of the functions $f(x, y)$. Enter 0 if there is no match.


1



2



3




Function $f(x, y) =$	0,1,2, or 3
$\sqrt{ 1 + x^2 - y^2 }$	
$\sin(x^2 + y^2)$	
$ x - y - x + y $	
$y^2 \sin(x^4)$	

c) (2 points) Match the space curves with the parametrizations. Enter 0 if there is no match.


1



2




3




Parametrization $\vec{r}(t) =$	0,1,2, or 3
$[t , t, t]$	
$[\cos(2t), 0, \cos(2t)]$	
$[0, \cos(2t), \sin(2t)]$	
$[t, \sin(t), 0]$	

d) (2 points) Match the functions g with contour plots in the xy-plane. Enter 0 if there is no match.

1



2




3




Function $g(x, y) =$	0,1,2, or 3
$\cos(2x) + \sin(2y)$	
$(x - y)^2$	
$y^2 - x$	
$(2x^2 + 7y^2)^2$	

e) (2 points) Match the parametrized surfaces. Enter 0 if there is no match.


1



2



3



Parametrization $\vec{r}(u, v) =$	0-3
$[u, u^2 + v^2, v]$	
$[\sin(v) \cos(u), \sin(v) \sin(u), \cos(v)]$	
$[u^2 - v^2, u, v]$	
$[u \cos(v), u \sin(v), u]$	

Problem 3) (10 points)

Let $\vec{v} = [2, 2, 1]$ and $\vec{w} = [1, 1, 1]$ and assume θ is the angle between $\vec{v}$ and $\vec{w}$.

a) (5 points) Compute $\vec{n} = \vec{v} \times \vec{w}$ and $a = \vec{v} \cdot \vec{w}$.

$\vec{n} =$

$a =$

b) (5 points) Compute $\sin^2(\theta)$ by using the vector $\vec{n}$ and $\cos^2(\theta)$ by using the scalar a and check that $\sin^2(\theta) + \cos^2(\theta)$ is equal to 1.

$$\sin^2(\theta) =$$

$$\cos^2(\theta) =$$

Check that the sum is 1:

Problem 4) (10 points)

a) (5 points) Parametrize the plane $4x + 4y + 2z = 0$ using parameters s, t :

$$\vec{r}(s, t) =$$

b) (5 points) Find the distance between $4x + 4y + 2z = 0$ and the point $(2, 2, 2)$.

distance =

Problem 5) (10 points)

a) (5 points) Find a parametrization $\vec{r}(t)$ of the intersection of the planes

$$x + y + z = 1, \quad 2x - y + 2z = 2 .$$

$\vec{r}(t) =$

b) (5 points) Find the distance between that line computed in a) and $P = (1, 1, 1)$.

distance =

Problem 6) (10 points)

a) (5 points) Find the arc length of the path

$$\vec{r}(t) = \left[\frac{3t^2}{2}, \frac{4t^2}{2}, \frac{5t^3}{3} \right]$$

with $0 \leq t \leq 1$.

Length =

b) (5 points) Find the curvature of $\vec{r}(t)$ at $t = 1$ using the vectors $\vec{v} = \vec{r}'(1)$, $\vec{w} = \vec{r}''(1)$.

$$\kappa(\vec{r}(1)) =$$

Problem 7) (10 points)

a) (5 points) Given the **jerk**

$$\vec{r}'''(t) = [0, 0, -12]$$

with $\vec{r}(0) = [0, 0, 3]$, $\vec{r}'(0) = [4, 0, 0]$, $\vec{r}''(0) = [0, 2, 0]$, find $\vec{r}(t)$ and especially $\vec{r}(10)$.

$$\vec{r}(10) =$$

b) (5 points) Compute the unit tangent vector $\vec{T}$ of the TNB-frame to the curve $\vec{r}(t)$ at $t = 0$.

$$\vec{T}(0) =$$

Problem 8) (10 points)

We experiment with a **paper air plane**. The wing tips are $C = (0, 3, 4)$ and $D = (0, -3, 4)$ the front is $A = (3, 0, 0)$ the back is $B = (-3, 0, 0)$.

a) (5 points) The sum of the areas of the triangles ABC and ABD is the total wing area. Find the wing area.

$$\text{Area} =$$

b) (5 points) The volume of the tetrahedron with vertices A, B, C, D is known to be 1/6th of the volume of the parallelepiped spanned by AB, AC, AD . Find the volume of the tetrahedron.

Volume =

Problem 9) (10 points) No justifications are needed.

a) (2 points) Parametrize the surface $x^2/9 + y^2/4 + z^2/4 = 1$.

$$\vec{r}(\theta, \phi) = \left[\text{ } \right], \left[\text{ } \right], \left[\text{ } \right]$$

b) (2 points) Parametrize the surface $x^2 = y^2 + z^2$ using an angle θ in the yz -plane.

$$\vec{r}(\theta, x) = \left[\text{ } \right], \left[\text{ } \right], \left[\text{ } \right]$$

c) (2 points) Parametrize the surface $y = x^4 - 6x^2z^2 + z^4$.

$$\vec{r}(x, z) = \left[\text{ } \right], \left[\text{ } \right], \left[\text{ } \right]$$

d) (2 points) Parametrize the surface $x^2 + z^2 = 4$ using an angle θ in the xz -plane.

$$\vec{r}(\theta, y) = \left[\text{ } \right], \left[\text{ } \right], \left[\text{ } \right]$$

e) (2 points) Parametrize the surface $(x - 7)^2 + z^2 - y^2 = 8$.

$$\vec{r}(\theta, y) = \left[\text{ } \right], \left[\text{ } \right], \left[\text{ } \right]$$

The following pictures are here just to be admired and can also be ignored.

