

Tranferring model-theoretic results about $\mathcal{L}_{\infty, \omega}$ to a Grothendieck topos

Nathanael Leedom Ackerman

Harvard University

Harvard and MIT Logic Seminar

March 25, 2013

Definition of $\mathcal{L}_{\infty, \omega}$

Definition

For a language \mathcal{L} , $\mathcal{L}_{\infty, \omega}(\mathcal{L})$ is the smallest collection of formulas which contains the language \mathcal{L} and is closed under:

- \wedge , \vee , \neg , and \Rightarrow .
- $(\forall x)$ and $(\exists x)$.
- Infinite disjunctions and conjunctions, so long as the result only has a finite number of free variables.

Why We Care About $\mathcal{L}_{\infty,\omega}(\mathfrak{L})$

Why We Care:

- $\mathcal{L}_{\infty,\omega}(\mathfrak{L})$ is natural a generalization of first order logic to infinite formulas.
- Many theorems of first order logic have analogs for $\mathcal{L}_{\infty,\omega}(\mathfrak{L})$. These include the downward Löwenheim-Skolem theorem, compactness and completeness.
- The satisfaction relation for $\mathcal{L}_{\infty,\omega}(\mathfrak{L})$ is absolute.

Definition of Grothendieck Topos

Definition

A Grothendieck topos is a category equivalent to the category of sheaves on a site.

In this talk we will restrict our attention to those Grothendieck toposes which are equivalent to sheaves on a countable lattice.

Why We Care About Grothendieck Toposes

Why We Care:

- Grothendieck toposes play an important role in a wide variety of areas ranging from topology to algebraic geometry to higher order logic.
- Grothendieck toposes are a natural generalization of the notion of *space* (as in topology)
- Grothendieck toposes are closely related to Kripke models as well as the notion of forcing.
- Grothendieck toposes are natural models of intuitionistic set theory.
- Grothendieck toposes have enough structure to interpret formulas of $\mathcal{L}_{\infty, \omega}(\mathcal{L})$.

Overall Goal

Overall Goal:

- Fix an (appropriate) Grothendieck topos G .
- Represent \mathfrak{L} -structure in G by models of a sentence in $\mathcal{L}_{\infty, \omega}(\mathfrak{L}')$ in the category of sets.
- Represent sentences of $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ (interpreted in G) by sentences of $\mathcal{L}_{\infty, \omega}(\mathfrak{L}')$ (interpreted in the category of sets).
- Use these representations to prove theorems about $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ interpreted in G that are analogous to those which are known about $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ interpreted in the category of sets.

Main Problem

Main Problem:

- Being a sheaf is a second order property and may not be absolute.
- Therefore being an \mathfrak{L} -structure in a category of sheaves may not be absolute.
- As the satisfaction relation of $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ is absolute, $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ may not be able to capture the class of \mathfrak{L} -structures in a (fixed) category of sheaves.

Absolute Version of a Grothendieck Topos

Lattice with Covers

Definition

Define a **lattice with covers** $\mathfrak{L}at = (L, \preceq, \wedge, \vee, \text{Cov}_{\mathfrak{L}at})$ to consist of the following:

- A distributive lattice $(L, \preceq, \wedge, \vee)$
- For each $p \in L$ a collection of **covering ideals**, $\text{Cov}_{\mathfrak{L}at}(p)$.

such that for each $p \in L$:

- $\{q : q \preceq p\} \in \text{Cov}_{\mathfrak{L}at}(p)$.
- For all $C \in \text{Cov}_{\mathfrak{L}at}(p)$:
 - For all $x \in C$, $x \preceq p$.
 - $p = \bigvee C$.
- If $C \in \text{Cov}_{\mathfrak{L}at}(p)$ then for any $q \in L$,
 $C \wedge q := \{x \wedge q : x \in C\} \in \text{Cov}_{\mathfrak{L}at}(q)$.

In order to simplify things we will assume $\mathfrak{L}at$ is countable.

Definition of Separated Presheaves

Definition

A **separated presheaf** \mathcal{S} on $\mathcal{L}at$ consists of:

- A collection of sets $\{\mathcal{S}(p) : p \in L\}$.
- A collection of commutative functions

$$\{i_{p,q}^{\mathcal{S}} : \mathcal{S}(p) \rightarrow \mathcal{S}(q) \text{ where } q \preceq p\}.$$

such that

- For each $C \in \text{Cov}_{\mathcal{L}at}(p)$ and $x, y \in \mathcal{S}(p)$:

$$\text{If } (\forall q \in C) i_{p,q}(x) = i_{p,q}(y) \text{ then } x = y.$$

Presheaf of Functions

Example

Notice that any $(T, \mathcal{O}(T))$ gives a lattice with covers where the lattice is $(\mathcal{O}(T), \subseteq)$ and $C \in \text{Cov}_T(U)$ if and only if $\bigcup_{V \in C} V = U$.

For each $U \in \mathcal{O}(T)$ let $\mathcal{F}(U) \subseteq X^U$ where $i_{U,V}(f) = f|_V$. Then \mathcal{F} is a separated presheaf.

We call such a collection of functions a **presheaf of functions** on $(T, \mathcal{O}(T))$.

Note that being a presheaf of functions is absolute between transitive models of set theory.

Definition of Compatible Collection

Definition

A **compatible collection** in a separated presheaf \mathcal{S} over $p \in L$ is a sequence $\langle x_q : q \in C \rangle$ where:

- $x_q \in \mathcal{S}(q)$.
- If $r \preceq q$ then $i_{q,r}(x_q) = x_r$.

We can think of a compatible collection as a **hole** which may be missing from \mathcal{S} .

Definition of Sheaf

Definition

A **sheaf** \mathcal{S} is a separated presheaf such that for each compatible collection $\langle x_q : q \in C \rangle$ there is an $x^* \in \mathcal{S}(p)$ where for all $q \in C$, $i_{p,q}(x^*) = x_q$.

A sheaf is then a separated presheaf where all the holes are filled in.

Problem:

Being a sheaf is not absolute. If we move to a larger model of set theory there may be new compatible collections.

Sheaf of Functions

Example

If \mathcal{F} is a presheaf of functions on $(T, \mathcal{O}(T))$ then \mathcal{F} is a sheaf if whenever:

$$\langle f_i : U_i \rightarrow X, i \in I \rangle \text{ is such that } f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$$

there is a function $f : \bigcup_{i \in I} U_i \rightarrow X$ in $\mathcal{F}(\bigcup_{i \in I} U_i)$ which, for each $i \in I$, agrees with f_i on U_i .

Covers

There is a **sheafification** functor, \mathbf{a} , which takes a separated presheaf and returns the *smallest* sheaf containing it.

Definition

We say a separated presheaf A **covers** B if,

- For all $p \in L$, $A(p) \subseteq B(p)$.
- $\mathbf{a}(A) \cong \mathbf{a}(B)$.

In particular **being a cover is absolute**.

Absolute Grothendieck Toposes

Definition

Let $\text{Sh}(\mathcal{L}\text{at})$ be the category where:

- The objects are the separated presheaves on $\mathcal{L}\text{at}$.
- A map from A to B is an (equivalence class of) pairs $\langle A^*, f \rangle$ where
 - A^* covers A
 - $f : A^* \rightarrow B$ is a map of separated presheaves.

where $\langle A_0^*, f_0 \rangle \equiv \langle A_1^*, f_1 \rangle$ if and only if

$$(\forall x \in A_0^* \cap A_1^*) f_0(x) = f_1(x).$$

Lemma

$\text{Sh}(\mathcal{L}\text{at})$ is equivalent to the category of sheaves on $\mathcal{L}\text{at}$.

Encoding With Ordinals

Subobjects

Definition

We say A is **closed** in B if for all $p \in L$ and for all $x \in B(p)$

$x \in A(p)$ if and only if $\{q : i_{p,q}(x) \in A(q)\} \in \text{Cov}_{\mathcal{L}\text{at}}(p)$

Lemma

There is a one-to-one relationship between

- *Closed subpresheaves of B .*
- *Subobjects of B in $Sh(\mathcal{L}\text{at})$.*

Subobjects are the analog of subsets in a Grothendieck topos

In an \mathcal{L} -structure in a Grothendieck topos

Formulas are interpreted by subobjects.

and

The logical operations are operations on the subobjects.

In an \mathcal{L} -structure in a Grothendieck topos

Formulas are interpreted by subobjects.

and

The logical operations are operations on the subobjects.

Example

Suppose each formula $\varphi_i(\mathbf{x})$ is interpreted by closed $X_i \subseteq A$.

Let $X^*(p) := \bigcap_{i \in I} X_i(p)$ for each $p \in L$.

Then $\bigwedge_{i \in I} \varphi_i(\mathbf{x})$ is interpreted by X^* .

Note it is easy to show that if each X_i is closed in A then so is X^* .

Constructing The Closure

The logical operations on presheaves don't always preserve being closed. As such we need the notion of a closure of a subpresheaf.

Definition

Suppose $A \subseteq B$. We define A_α by induction as follows:

- $A_0 := A$.
- $A_{\omega \cdot \beta} := \bigcup_{\gamma < \omega \cdot \beta} A_\gamma$.
- $A_{\alpha+1}(p) := \{x \in B(p) : \{q : i_{p,q}(x) \in A_\alpha\} \in \text{Cov}_{\text{Nat}}(p)\}$.
- $A_\infty := \bigcup_{\gamma \in \text{ORD}} A_\gamma$.

Lemma

For any $A \subseteq B$, A_∞ is the smallest subpresheaf of B which contains A and is closed in B . We call A_∞ the **closure** of A in B .

Disjunction Operator

We can use the construction of the closure to define the other logical operations.

Example

Suppose each formula $\varphi_i(\mathbf{x})$ interpreted by closed $X_i \subseteq A$.

Let $X^*(p) := \bigcup_{i \in I} X_i(p)$ for each $p \in L$.

Then $\bigvee_{i \in I} \varphi_i(\mathbf{x})$ is interpreted by X_∞^* .

Note even if each X_i is closed in A , X^* may not be.

Encoding Separated Presheaves

Definition

Let $\mathcal{L}_{\mathcal{L}at}$ be the language where:

- For each $p \in L$ there is a sort \bar{p} .
- For each $q \preceq p$ there is a function $\overline{i_{p,q}} : \bar{p} \rightarrow \bar{q}$.

Let $Th_{\mathcal{L}at} \in \mathcal{L}_{\omega_1, \omega}(\mathcal{L}_{\mathcal{L}at})$ be the theory which says:

- All $\overline{i_{p,q}}$ commute.
- For each $C \in \text{Cov}_{\mathcal{L}at}(p)$

$$(\forall x, y : \bar{p}) \left[\bigwedge_{q \in C} \overline{i_{p,q}}(x) = \overline{i_{p,q}}(y) \right] \rightarrow x = y.$$

Models of $Th_{\mathcal{L}at}$ then **encode separated presheaves**.

Encoding Ordinals

In order to encode the closure operation we need to be able to represent the ordinals.

Definition

Let \mathfrak{L}_{ω_1} be the language where:

- There is a unique sort \bar{O} .
- For each $\beta \leq \omega_1$ there is a constant c_β of sort \bar{O} .
- There is a binary relation \leq_O on \bar{O} .

Let $Th_{\omega_1} \in \mathcal{L}_{\infty, \omega}(\mathfrak{L}_{\omega_1})$ be the theory which says:

- For each $i \leq j \leq \omega_1$, $c_i \leq_O c_j$ and \leq_O is a linear order.
- $(\forall x : \bar{O}) \bigvee_{\beta \leq \omega_1} x = c_\beta$.

Note Th_{ω_1} is **not** in $\mathcal{L}_{\omega_1, \omega}(\mathfrak{L}_{\omega_1})$ as it needs a disjunction of size ω_1 .

Encoding The Closure

Definition

Let $\mathfrak{L}_{Cl} = \mathfrak{L}_{\text{Nat}} \cup \mathfrak{L}_{\omega_1} \cup \{R_p : p \in L\}$ where R_p is of sort $\bar{p} \times \bar{O}$.

Let T_{Cl} be the theory which says:

- Whenever $q \preceq p$, $(\forall x : \bar{p})(\forall a : \bar{O}) R_p(x, a) \rightarrow R_q(\overline{i_{p,q}}(x), a)$.
- For each $\alpha + 1 \leq \omega_1$ and each $p \in L$,

$$(\forall x : \bar{p}) R_p(x, c_{\alpha+1}) \leftrightarrow \bigvee_{C \in \text{Cov}(p)} \bigwedge_{q \in C} R_q(\overline{i_{q,p}}(x), c_\alpha).$$

- For each limit $\beta \leq \omega_1$ and each $p \in L$,

$$(\forall x : \bar{p}) R_p(x, c_\beta) \leftrightarrow \bigvee_{\gamma < \beta} R_p(x, c_\gamma).$$

Encoding The Closure

Suppose

- The sorts of our language represent a separated presheaf S
- The relations $\{R_p(-, c_0) : p \in L\}$ represent $R \subseteq S$.
- The relations $\{R_p(-, c_{\omega_1}) : p \in L\}$ represent $R^* \subseteq S$.

Then R^* is the closure of R in S .

Encode Disjunctions

We can now show how to use the ordinals to encode a disjunction.

Example

Suppose we have

- For each $i \in \omega$, relations $\{\overline{R}_p^i : p \in L\}$ which encode subpresheaves R^i of S .
- Relations $\{\overline{Q}_p : p \in L\}$ where \overline{Q}_p is of sort $\overline{p} \times \overline{O}$.

and the theory which says:

$$T_{Cl} \text{ and } (\forall x : \overline{p}) \overline{Q}_p(x, c_0) \leftrightarrow \bigvee_{i \in \omega} \overline{R}_p^i(x) \text{ for each } p \in L.$$

Then $\{\overline{Q}_p(-, c_{\omega_1}) : p \in L\}$ represents the formulas $\bigvee_{i \in \omega} R^i$.

Encoding Theorem

Putting these together we have the following theorem.

Theorem

Fix a countable language \mathfrak{L} .

Then there is a language \mathfrak{L}' and a theory $T_{enc} \in \mathcal{L}_{\omega_1, \omega}(\mathfrak{L}')$ such that models $\mathcal{M} \models T_{Enc}$ encode \mathfrak{L} -structures $\mathcal{S}_{\mathcal{M}}$ in $Sh(\mathfrak{L}at)$.

Further, for any sentence $\varphi \in \mathcal{L}_{\omega_1, \omega}(\mathfrak{L})$ there is a sentence $Th_{\varphi} \in \mathcal{L}_{\omega_2, \omega}(\mathfrak{L}')$ such that $\mathcal{M} \models Th_{\varphi}$ if and only if $\mathcal{S}_{\mathcal{M}} \models \varphi$ (in $Sh(\mathfrak{L}at)$).

In this way we can think of Th_{φ} as encoding the sentence φ .

Encoding Theorem

Corollary

Suppose \mathcal{M} is an \mathfrak{L} -structure in $\text{Sh}(\mathfrak{Set})$, and V_0, V_1 are transitive well-founded models of set theory with $\mathfrak{Set}, \mathcal{M}, \omega_1 \in V_0 \cap V_1$.

For each $\varphi \in \mathcal{L}_{\omega_1, \omega}(\mathfrak{L}) \cap V_0 \cap V_1$ we have:

$$(\mathcal{M} \models \varphi)^{V_0} \text{ if and only if } (\mathcal{M} \models \varphi)^{V_1}.$$

In particular this implies the **satisfaction relation for $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ in $\text{Sh}(\mathfrak{Set})$ is absolute.**

Transfer Theorems

Downward Löwenheim-Skolem Theorem

Definition

A separated presheaf S is of **generated size** at most κ if

$$\left| \bigcup_{p \in L} S(p) \right| \leq \kappa$$

Theorem (Downward Löwenheim-Skolem Theorem)

Let \mathcal{M} be an \mathfrak{L} -structure in $Sh(\mathfrak{Lat})$, $X \subseteq \mathcal{M}$, and $\varphi \in \mathcal{L}_{\omega_1, \omega}(\mathfrak{L})$.
 Then there is an \mathfrak{L} -structure in $Sh(\mathfrak{Lat})$, \mathcal{M}_X , where:

- $X \subseteq \mathcal{M}_X \subseteq \mathcal{M}$.
- $|\mathcal{M}_X| = |X|$.
- $\mathcal{M}_X \models \varphi$

Downward Löwenheim-Skolem Theorem

Proof.

Let V be a Σ_n -elementary submodel of the universe of sets (for some sufficiently large n) where $X \in V$ and $|V| = |X|$.

Then there is an \mathcal{L} -structure \mathcal{M}_X in $\text{Sh}(\mathcal{L}\text{at})$ such that $\mathcal{M}_X \in V$, $X \subseteq \mathcal{M}_X$ and $\mathcal{M}_X \models \varphi$.

But then in the actual universe $|\mathcal{M}_X| = |X|$ and $\mathcal{M}_X \models \varphi$ as the satisfaction relation is absolute.



Directed Limits

Theorem (Directed Limits)

Suppose $F \subseteq \mathcal{L}_{\infty, \omega}(\mathfrak{L})$ is a fragment and $\langle \mathcal{M}_i : i \in \kappa \rangle$ is a chain of \mathfrak{L} structures in $Sh(\mathfrak{L}at)$ where for each $i < j$:

- $\mathcal{M}_i \subseteq \mathcal{M}_j$ with inclusion map $\iota_{i,j}$.
- $\iota_{i,j}$ preserves all formulas in F .

Let \mathcal{M}^* be the directed limit of $\langle \mathcal{M}_i : i \in \kappa \rangle$ in $Sh(\mathfrak{L}at)$. Then each inclusion map $\iota_i^* : \mathcal{M}_i \rightarrow \mathcal{M}^*$ preserves each formula in F as well.

Proof.

This follows from the corresponding result for fragments of $\mathcal{L}_{\infty, \omega}(\mathfrak{L})$ in the category of sets. □

Barwise Compactness

Recall the following theorem in the category of sets

Theorem (Barwise Compactness)

Suppose A is a countable admissible set and $T \subseteq \mathcal{L}_{\omega_1, \omega}(\mathcal{L}) \cap A$ where:

- T is Σ_1 definable over A .
- For every $T_0 \subseteq T$ which is Δ_1 definable over A there is an \mathcal{M} with $\mathcal{M} \models T_0$.

Then there is a model which satisfies T .

Sheaf Compactness

Theorem (Sheaf Compactness)

Suppose A is a countable but not locally countable Σ_1 -admissible set for which there is a well-ordering Σ_1 -definable over A .

If $T \subseteq \mathcal{L}_{\omega_1, \omega}(\mathfrak{L}) \cap A$ is such that:

- *T is Σ_1 definable over A .*
- *For every $T_0 \subseteq T$ which is Δ_1 definable over A there is an \mathfrak{L} -structure \mathcal{M} in $Sh(\mathfrak{L}at)$ with $\mathcal{M} \in A$ and $\mathcal{M} \models T_0$.*

Then there is an \mathfrak{L} -structure in $Sh(\mathfrak{L}at)$ which satisfies T .