

LOCAL-GLOBAL COMPATIBILITY OVER FUNCTION FIELDS

SIYAN DANIEL LI-HUERTA

ABSTRACT. We prove that V. Lafforgue’s global Langlands correspondence is compatible with Fargues–Scholze’s semisimplified local Langlands correspondence. As a consequence, we canonically lift Fargues–Scholze’s construction to a non-semisimplified local Langlands correspondence for local fields of characteristic $p \geq 5$. We also deduce that Fargues–Scholze’s construction agrees with that of Genestier–Lafforgue, answering a question of Fargues–Scholze, Hansen, Harris, and Kaletha. The proof relies on a uniformization morphism for moduli spaces of shtukas.

CONTENTS

Introduction	1
1. Recollections on affine Grassmannians	5
2. Formal moduli of local shtukas	8
3. Relative z -adic Hodge theory	16
4. Analytic moduli of local shtukas	23
5. Uniformizing the moduli spaces of global shtukas	33
6. Local-global compatibility	41
7. Applications	50
References	52

INTRODUCTION

The Langlands program predicts a relationship between automorphic forms and Galois representations. More precisely, in the case of a connected reductive group \mathbf{G} over a global function field \mathbf{F} of characteristic $p > 0$, the Langlands program posits a canonical map

$$\mathrm{GLC}_{\mathbf{G}} : \left\{ \begin{array}{l} \text{cuspidal automorphic} \\ \text{representations of } \mathbf{G}(\mathbb{A}_{\mathbf{F}}) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} L\text{-parameters} \\ \text{for } \mathbf{G} \text{ over } \mathbf{F} \end{array} \right\},$$

where $\mathbb{A}_{\mathbf{F}}$ denotes the adèle ring of \mathbf{F} , and all representations are taken with $\overline{\mathbb{Q}}_{\ell}$ -coefficients for some $\ell \neq p$. In a landmark result, such a map $\mathrm{GLC}_{\mathbf{G}}$ was constructed by V. Lafforgue [32].

In the case of a connected reductive group G over a nonarchimedean local field F , the Langlands program predicts a similar map

$$(\dagger) \quad \mathrm{LLC}_G : \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } G(F) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} L\text{-parameters} \\ \text{for } G \text{ over } F \end{array} \right\}.$$

Recent breakthrough work of Fargues–Scholze [11] constructs such a map up to semisimplification; namely, they construct a map

$$(\ddagger) \quad \text{LLC}_G^{\text{ss}} : \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } G(F) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{semisimple } L\text{-parameters} \\ \text{for } G \text{ over } F \end{array} \right\}.$$

Our main result is that V. Lafforgue’s global Langlands correspondence is compatible with Fargues–Scholze’s semisimplified local Langlands correspondence.

Theorem A. *Let v be a place of \mathbf{F} . Then the square*

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{cuspidal automorphic} \\ \text{representations of } \mathbf{G}(\mathbb{A}_{\mathbf{F}}) \end{array} \right\} & \xrightarrow{\text{GLC}_{\mathbf{G}}} & \left\{ \begin{array}{l} L\text{-parameters} \\ \text{for } \mathbf{G} \text{ over } \mathbf{F} \end{array} \right\} \\ \downarrow (-)_v & & \downarrow (-)|_{W_{\mathbf{F}_v}}^{\text{ss}} \\ \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } \mathbf{G}(\mathbf{F}_v) \end{array} \right\} & \xrightarrow{\text{LLC}_{\mathbf{G}_{\mathbf{F}_v}}^{\text{ss}}} & \left\{ \begin{array}{l} \text{semisimple } L\text{-parameters} \\ \text{for } \mathbf{G}_{\mathbf{F}_v} \text{ over } \mathbf{F}_v \end{array} \right\} \end{array}$$

commutes.

Since $\text{GLC}_{\mathbf{G}}$ [32, Théorème 12.3] and LLC_G^{ss} [11, Theorem IX.0.5] are compatible with the Satake isomorphism at unramified places, for a given cuspidal automorphic representation this is already known at *unramified* places.

We actually prove a refinement of Theorem A on the level of *excursion algebras*; see Theorem 6.13.

Remarks.

- (1) V. Lafforgue [32, Théorème 13.2] and Fargues–Scholze [11, Proposition IX.4.1] prove a version of their results with $\overline{\mathbb{F}}_{\ell}$ -coefficients, and the analogous version of Theorem A also holds in this mod- ℓ context. See Theorem 6.15.
- (2) Once one constructs a non-semisimplified local Langlands correspondence as in Equation (\ddagger) (e.g. see Theorem B below), one can ask whether Theorem A holds before semisimplification. The answer is already negative when \mathbf{G} is the units of a quaternion algebra [14, Remarque 0.3]. More generally, Arthur’s conjecture [4] predicts that the answer is negative precisely for global A -packets where a local A -packet intersects more than one local L -packet. For instance, examples of Howe–Piatetski-Shapiro [25] show that the answer is also negative when \mathbf{G} is Sp_4 .

We now turn to some consequences of Theorem A. When $\text{char } F > 0$ is not too small, Theorem A enables us to remove the “up to semisimplification” ambiguity in Fargues–Scholze’s construction.

Theorem B. *Assume that $\text{char } F = p \geq 5$. Then LLC_G^{ss} canonically lifts to a non-semisimplified local Langlands correspondence LLC_G as in Equation (\ddagger) .*

Actually, we only need to assume that p is good for the non-simply laced absolute factors of G ; see Theorem 7.1. The proof that Theorem A implies Theorem B is due to Gan–Harris–Sawin [12]; roughly, the idea is to maneuver into a situation where Theorem A holds even before semisimplification. This uses a globalization result of Beuzart-Plessis [12], work of Heinloth–Ngô–Yun [24] on Kloosterman sheaves, and Deligne’s purity theorem.

Our next result concerns previous work of Genestier–Lafforgue [14], who also constructed a map as in Equation (\ddagger) when $\text{char } F > 0$. Genestier–Lafforgue obtained

a version of Theorem A for their construction, and since this property basically uniquely characterizes such maps, we deduce the following result.

Theorem C. *The Genestier–Lafforgue correspondence agrees with the Fargues–Scholze correspondence.*

This answers a question of Fargues–Scholze [11], Hansen, Harris, and Kaletha [28]. We also prove a refinement of Theorem C on the level of Bernstein centers; see Theorem 6.15.

Remark. Conversely, if we only had Theorem C, then work of Genestier–Lafforgue would imply Theorem A. However, our proof of Theorem A is independent of their results.

We conclude by showing that LLC_G^{ss} satisfies the expected compatibility with the local Jacquet–Langlands correspondence [5], which we denote by JL , when $\text{char } F > 0$ and G is the units of a central simple algebra over F .

Theorem D. *Assume that $\text{char } F > 0$ and G is the units of a central simple algebra over F . For any irreducible essentially L^2 representation π of $G(F)$, we have $\text{LLC}_G^{\text{ss}}(\pi) = \text{LLC}_{\text{GL}_n}^{\text{ss}}(\text{JL}(\pi))$.*

When $\text{char } F > 0$, Theorem D was previously only known when G is GL_n or the units of a central division algebra over F [11, Theorem IX.7.4]. The $\text{char } F = 0$ analogue of Theorem D is due to Hansen–Kaletha–Weinstein [19, Theorem 6.6.1] as a consequence of their work on the local Kottwitz conjecture.

Let us discuss our proof of Theorem A. Elements of our strategy go back to Deligne’s letter to Piatetski-Shapiro [10], which proves local-global compatibility for modular forms. Their associated Galois representations are constructed via the cohomology of modular curves, and one of Deligne’s key ideas was to restrict to the supersingular locus, using the uniformization of the latter by Lubin–Tate space to relate the local and global Langlands correspondences for GL_2 .

Deligne’s proof, as well as subsequent works on local-global compatibility using basic uniformization [9, 21, 40, 35], also crucially relies on arguments specific to the particular group \mathbf{G} in question. However, our proof of Theorem A is uniform in all groups \mathbf{G} .

We begin by observing that, since the correspondences of V. Lafforgue and Fargues–Scholze are constructed via *excursion operators*, it suffices to show that said operators are compatible. Let us recall their definition, which involves moduli spaces of shtukas. For simplicity, assume that \mathbf{G} is split, and write $\widehat{\mathbf{G}}$ for the dual group of \mathbf{G} over $\overline{\mathbb{Q}_\ell}$. For any finite set I and representation V of $\widehat{\mathbf{G}}^I$, write $\text{Sht}_{\mathbf{G}, V}^I$ for the associated moduli space of *global \mathbf{G} -shtukas*,¹ which is a Deligne–Mumford stack. Work of Xue [44] naturally endows the compactly supported intersection cohomology H_V^I of its generic fiber with an action of $W_{\mathbf{F}}^I$, where $W_{\mathbf{F}}$ denotes the absolute Weil group. For any x and ξ in V and V^\vee , respectively, that are fixed by the image of $\Delta : \widehat{\mathbf{G}} \hookrightarrow \widehat{\mathbf{G}}^I$, and any γ_\bullet in $W_{\mathbf{F}}^I$, the associated global excursion operator is

$$(\heartsuit) \quad H_{\mathbf{1}}^* \xrightarrow{x} H_{V|_{\Delta(\widehat{\mathbf{G}})}}^* = H_V^I \xrightarrow{\gamma_\bullet} H_V^I = H_{V|_{\Delta(\widehat{\mathbf{G}})}}^* \xrightarrow{\xi} H_{\mathbf{1}}^*,$$

where $*$ denotes the singleton set, and $\mathbf{1}$ denotes the trivial representation.

¹In the introduction, we ignore convolution data and level structures in our notation.

In the local setting, write $\mathcal{L}\text{ocSht}_{\mathbf{G},V}^I$ for the associated moduli space of *local \mathbf{G} -shtukas*, which is an analytic adic space. Work of Fargues–Scholze [11] naturally endows the intersection homology $H_V^{\text{loc},I}$ of its generic fiber with an action of $W_{\mathbf{F}_v}^I$, so when γ_\bullet lies in $W_{\mathbf{F}_v}^I$, one can form local excursion operators using the same recipe as in Equation (♥).

We compare the local and global excursion operators using a uniformization morphism. To define it, first we construct a formal model $\mathfrak{L}\text{ocSht}_{\mathbf{G},V}^I$ for $\mathcal{L}\text{ocSht}_{\mathbf{G},V}^I$ at hyperspecial level. Stating the formal moduli problem is straightforward, although comparing it with our original definition of local \mathbf{G} -shtukas requires an equicharacteristic version of Kedlaya–Liu’s results [31] on relative p -adic Hodge theory, which we prove. Next, we use Beauville–Laszlo gluing to construct a formally étale morphism of formal stacks

$$\widehat{\Theta} : \mathfrak{L}\text{ocSht}_{\mathbf{G},V}^I \rightarrow \widehat{\text{Sht}}_{\mathbf{G},V}^I$$

when the level is hyperspecial at v , where $\widehat{\text{Sht}}_{\mathbf{G},V}^I$ denotes the formal completion of $\text{Sht}_{\mathbf{G},V}^I$ along v^I , and we assume that $\deg v = 1$ for simplicity. This generalizes results of Arasteh Rad–Hartl [3].

From here, we restrict to a Harder–Narasimhan truncation $\text{Sht}_{\mathbf{G},V}^{I,\leq s}$ of $\text{Sht}_{\mathbf{G},V}^I$ and enlarge the level away from v . This yields a scheme that is locally of finite type, so we can use Huber’s analytification [26, (3.8)] to extend $\widehat{\Theta}$ to a morphism of analytic adic spaces

$$\Theta : \mathcal{L}\text{ocSht}_{\mathbf{G},V}^{I,\leq s} \rightarrow (\text{Sht}_{\mathbf{G},V}^{I,\leq s})_{(\text{Spa } \mathbf{F}_v)^I}.$$

for deeper levels at v . To prove that Θ is étale, it suffices to consider the case of hyperspecial level. There, we prove that $\mathfrak{L}\text{ocSht}_{\mathbf{G},V}^I$ is a formal scheme that is locally formally of finite type, generalizing results of Arasteh Rad–Hartl [2]. After restricting to a Harder–Narasimhan truncation, this lets us upgrade the formal étaleness of $\widehat{\Theta}$ to étaleness, as desired.

Since Θ is étale, we can form the !-pushforward map

$$\Theta_! : H_V^{\text{loc},I,\leq s} \rightarrow H_V^{I,\leq s}.$$

After restricting to a Harder–Narasimhan truncation, this induces a morphism from the composition diagram in Equation (♥) to the analogous composition diagram for $H_V^{\text{loc},I}$. We use this to prove that the global and local excursion operators are compatible, which concludes the proof of Theorem A.

With Theorem A in hand, let us return to the local context and sketch the proofs of Theorem B, Theorem C, and Theorem D. For Theorem B, compatibility with parabolic induction and the Langlands classification reduce us to the case of L^2 representations π . Then the Langlands program predicts $\text{LLC}_G(\pi)$ to be the unique pure L -parameter whose semisimplification is $\text{LLC}_G^{\text{ss}}(\pi)$, if it exists. To construct this L -parameter, we use a globalization result of Beuzart–Plessis [12] to obtain a cuspidal automorphic representation Π that has the same cuspidal support as π at one place and is isomorphic to the cuspidal representation π' considered by Gross–Reeder [16] at another place. Using Theorem A and work of Heinloth–Ngô–Yun [24], we show that the Fargues–Scholze parameter of π' is irreducible. Therefore applying Deligne’s purity theorem to the V . Lafforgue parameter of Π and using Theorem A again yield the desired result.

For Theorem C, we instead reduce to the case of cuspidal representations. Then a classical Poincaré series argument and Theorem A give the desired result. Finally, for Theorem D we use the simple trace formula to construct a cuspidal automorphic representation of GL_n that globalizes $\mathrm{JL}(\pi)$ and transfers to a suitable central division algebra under the global Jacquet–Langlands correspondence [6]. From here, the Chebotarev density theorem and Theorem A imply the desired result.

Outline. In §1, we recall some facts about loop groups and Beilinson–Drinfeld affine Grassmannians. In §2, we define the formal moduli problem and prove that it is a formal scheme that is locally formally of finite type. In §3, we prove the necessary results on z -adic Hodge theory. In §4, we define the analytic moduli problem, compare it with the formal moduli problem, and recall results of Fargues–Scholze [11] on its intersection homology. In §5, we recall the global moduli problem and construct the uniformization morphism. In §6, we use this to prove Theorem A. In §7, we use Theorem A to prove Theorem B, Theorem C, and Theorem D.

Notation. Unless otherwise specified, all products are taken over \mathbb{F}_q . The transition morphisms for our ind-schemes are required to be closed embeddings. We view all functors between derived categories as derived functors.

Starting in §3, we freely use definitions from perfectoid geometry as in [41] and [11]. When viewing an adic space X as a locally ringed space, we take \mathcal{O}_X for its structure sheaf. For any adic space X over \mathbb{Z}_p , write X^\diamond for the associated v -sheaf over \mathbb{F}_p as in [42, Lemma 18.1.1].

Acknowledgements. The author thanks Mark Kisin for his patience and advice. The author would also like to thank Michael Harris for giving a talk on [12] that motivated him to prove Theorem A, and to thank David Hansen for his interest and encouragement.

1. RECOLLECTIONS ON AFFINE GRASSMANNIANS

In this section, we begin by setting up our local context. We then establish some notation on loop groups, Beilinson–Drinfeld affine Grassmannians, and their affine Schubert varieties, as well as recall basic facts about these objects. Nothing in this section is new.

1.1. Let F be a local field of characteristic $p > 0$, and write \mathbb{F}_q for its residue field. Fix a separable closure \bar{F} of F , and write Γ_F for $\mathrm{Gal}(\bar{F}/F)$. Choose a uniformizer z of \mathcal{O}_F , which yields an identification $\mathcal{O}_F = \mathbb{F}_q[[z]]$. Let G be a parahoric group scheme over \mathcal{O}_F as in [8, 5.2.6].

It will be convenient to use the following globalization of our local setup, although we will see that our constructions are independent of this globalization.

Lemma. *There exists a geometrically connected smooth proper curve C over \mathbb{F}_q , a nonempty open subspace $U \subseteq C$, a parahoric group scheme G_C over C as in [38, Definition 2.18], a closed point v of C , and an isomorphism $\widehat{\mathcal{O}}_{C,v} \cong \mathcal{O}_F$ such that*

- a) $G_C|_U$ is reductive over U ,
- b) $G_C|_{\mathcal{O}_v}$ is identified with G as group schemes over $\widehat{\mathcal{O}}_{C,v} \cong \mathcal{O}_F$.

Moreover, there exists an SL_h -bundle \mathcal{V} on C and a closed embedding

$$\iota : G_C \rightarrow \underline{\mathrm{Aut}}(\mathcal{V})$$

of group schemes over C such that $\underline{\text{Aut}}(\mathcal{V})/G_C$ is quasi-affine over C .

Proof. By [38, Lemma 3.1], there exists a connected smooth curve \hat{C} over \mathbb{F}_q , a smooth affine group scheme \hat{G} over \hat{C} with geometrically connected fibers, a closed point v of \hat{C} , and an isomorphism $\hat{\mathcal{O}}_{C,v} \cong \mathcal{O}_F$ such that $\hat{G}|_{\hat{C} \setminus v}$ is reductive over $\hat{C} \setminus v$ and $\hat{G}|_{\hat{\mathcal{O}}_{C,v}}$ is identified with G as group schemes over $\hat{\mathcal{O}}_{C,v} \cong \mathcal{O}_F$. Because \hat{C} has an \mathbb{F}_q -point v , it is geometrically connected. Write C for the associated smooth proper curve over \mathbb{F}_q . Fpqc descent and [8, 5.1.9] yield a parahoric group scheme G_C over C as in [38, Definition 2.18] that extends \hat{G} , so we can take $U = \hat{C} \setminus v$. Finally, the last claim follows from [3, Proposition 2.2(b)]. \square

1.2. Let us recall some facts about loop groups and affine Grassmannians. Let $S = \text{Spec } R$ be an affine scheme over C^I , and for all i in I , write Γ_i for the graph of its i -th projection $S \rightarrow C$, which is a relative effective Cartier divisor on $C \times S$.

Let I_1, \dots, I_k be an ordered partition of I . Write $\hat{\mathcal{O}}_C(S)$ for the ring of global sections of the completion of $\mathcal{O}_{C \times S}$ along $\sum_{i \in I} \Gamma_i$. For all $1 \leq j \leq k$, write $\hat{\mathcal{O}}_C^{j,\circ}(S)$ for the version that is punctured along $\sum_{i \in I_j} \Gamma_i$.

Definition.

- a) Write $L_I^n(G_C)$, $L_I^+(G_C)$, and $L_I^{j,\circ}(G_C)$ for the sheaves over C^I given by sending S to $G_C(\mathcal{O}_{n \sum_{i \in I} \Gamma_i})$, $G_C(\hat{\mathcal{O}}_C(S))$, and $G_C(\hat{\mathcal{O}}_C^{j,\circ}(S))$, respectively.
- b) Write $\text{Gr}_{G_C}^{(I_1, \dots, I_k)}$ for the sheaf over C^I whose S -points parametrize data consisting of
 - i) for all $1 \leq j \leq k$, a G -bundle \mathcal{G}_j on $\text{Spec } \hat{\mathcal{O}}_C(S)$,
 - ii) for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\phi_j : \mathcal{G}_j|_{\text{Spec } \hat{\mathcal{O}}_C^{j,\circ}(S)} \xrightarrow{\sim} \mathcal{G}_{j+1}|_{\text{Spec } \hat{\mathcal{O}}_C^{j,\circ}(S)},$$

where \mathcal{G}_{k+1} denotes the trivial G -bundle.

Write $L_z^+ G$ and $L_z G$ for the fiber at v of $L_*^+(G_C)$ and $L_*^{1,\circ}(G_C)$, respectively, where $*$ denotes the singleton set. Also, write $\text{Gr}_{z,G}^k$ for the fiber at v^I of $\text{Gr}_{G_C}^{\{\{1\}, \dots, \{k\}\}}$.

The proof of [18, Lemma 3.2] shows that $L_I^n(G_C)$ is an affine scheme of finite type over C^I , so $L_I^+(G_C) = \varprojlim_n L_I^n(G_C)$ is an affine scheme over C^I . Recall that $L_I^{j,\circ}(G_C)$ is an ind-affine ind-scheme over C^I [18, Lemma 3.2(i)], and $\text{Gr}_{G_C}^{(I_1, \dots, I_k)}$ is an ind-projective ind-scheme over C^I [3, Proposition 3.12]. Also, note that $L_z^+ G$, $L_z G$, and $\text{Gr}_{z,G}^k$ are independent of the globalization from Lemma 1.1.

1.3. The following lemmas give an alternative description of the Beilinson–Drinfeld affine Grassmannian after completing at a point. Write \mathbb{D} for the formal scheme $\text{Spf } \mathcal{O}_F$, and let I be a finite set. Recall that Spec yields an anti-equivalence from the category of $\mathbb{F}_q[[\zeta_i]]_{i \in I}$ -algebras where the ζ_i are nilpotent to the category of affine schemes over \mathbb{D}^I . Let $S = \text{Spec } R$ be an affine scheme over \mathbb{D}^I .

Lemma. *The direct system $(n \sum_{i \in I} \Gamma_i)_{n \geq 0}$ of schemes over $C \times S$ is naturally isomorphic to $(nv \times S)_{n \geq 0}$. Consequently, $\hat{\mathcal{O}}_C(S)$ is naturally isomorphic to $R[[z]]$, and $\hat{\mathcal{O}}_C^{j,\circ}(S) = \hat{\mathcal{O}}_C(S)[\frac{1}{z-\zeta_i}]_{i \in I_j} = R[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}$ is naturally isomorphic to $R((z))$.*

Proof. As nilpotent thickenings are étale-local and C is smooth at v , it suffices to replace C with $\mathbb{A}_{\mathbb{F}_q}^1 = \text{Spec } \mathbb{F}_q[z]$ and v with the origin. Then $n \sum_{i \in I} \Gamma_i$ is the

vanishing locus of $\prod_{i \in I} (z - \zeta_i)^n$ in $C \times S = \text{Spec } R[z]$, and $nv \times S$ is the vanishing locus of z^n in $C \times S$. Choose positive integers n_i such that $\zeta_i^{n_i} = 0$ in R .

Set $N_1 := \sum_{i \in I} n + n_i - 1$ and $N_2 := n + \max_{i \in I} \{n_i\} - 1$. On $n \sum_{i \in I} \Gamma_i$, we see

$$z^{N_1} = \prod_{i \in I} ((z - \zeta_i) + \zeta_i)^{n+n_i-1} = \prod_{i \in I} \sum_{l=1}^{n+n_i-1} \binom{n+n_i-1}{l} (z - \zeta_i)^l \zeta_i^{n+n_i-1-l} = 0,$$

so $n \sum_{i \in I} \Gamma_i$ lies in $N_1 v \times S$. Conversely, on $nv \times S$, we have

$$\prod_{i \in I} (z - \zeta_i)^{N_2} = \prod_{i \in I} \sum_{l=1}^{N_2} \binom{N_2}{l} z^l \zeta_i^{N_2-l} = 0,$$

so $nv \times S$ lies in $N_2 \sum_{i \in I} \Gamma_i$. \square

1.4. Write $\widehat{\text{Gr}}_G^{(I_1, \dots, I_k)}$ for the formal completion of $\text{Gr}_{G_C}^{(I_1, \dots, I_k)}$ along v^I in C^I .

Lemma. *Our $\widehat{\text{Gr}}_G^{(I_1, \dots, I_k)}$ is an ind-projective ind-scheme over \mathbb{D}^I , and it is naturally isomorphic to $\text{Gr}_{z, G}^k |_{\mathbb{D}^I}$.*

Thus $\widehat{\text{Gr}}_G^{(I_1, \dots, I_k)}$ is independent of the globalization from Lemma 1.1.

Proof. This follows immediately from Lemma 1.3. \square

1.5. We now introduce affine Schubert varieties. Write $\overline{\mathbb{F}_q(C)}$ for the separable closure of $\mathbb{F}_q(C)$ in \overline{F} , and write $\Gamma_{\mathbb{F}_q(C)}$ for $\text{Gal}(\overline{\mathbb{F}_q(C)}/\mathbb{F}_q(C))$. Let T be a maximal subtorus of $G_C |_{\mathbb{F}_q(C)}$, and write $X_*^+(T)$ for the set of dominant cocharacters of $T_{\overline{\mathbb{F}_q(C)}}$ with respect to a fixed Borel subgroup $B \subseteq G_C |_{\overline{\mathbb{F}_q(C)}}$ containing $T_{\overline{\mathbb{F}_q(C)}}$. Identify $X_*^+(T)$ with the set of conjugacy classes of cocharacters of $G_C |_{\overline{\mathbb{F}_q(C)}}$.

Let $\mu_\bullet = (\mu_i)_{i \in I}$ be in $X_*^+(T)^I$. Identify the the field of definition of μ_i with $\mathbb{F}_q(C_i)$ for some finite cover $C_i \rightarrow C$ that is étale over U , and write U_i for the preimage of U . Note that the closure F_i of $\mathbb{F}_q(C_i)$ in \overline{F} equals the completion of $\mathbb{F}_q(C_i)$ at the closed point v_i of C_i above v induced by $\overline{\mathbb{F}_q(C)} \rightarrow \overline{F}$. Write \mathbb{D}_i for $\text{Spf } \mathcal{O}_{F_i}$.

Definition.

- Write $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} U_i} \subseteq \text{Gr}_{G_C}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} U_i}$ for the associated closed affine Schubert variety, and write $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} C_i}$ for its closure in $\text{Gr}_{G_C}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} C_i}$.
- Write $\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$ for the formal completion of $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} C_i}$ along $\prod_{i \in I} v_i$ in $\prod_{i \in I} C_i$.
- When $I = *$, write $\text{Gr}_{z, G, \mu}^1 |_{v_*}$ for the fiber at v_* of $\text{Gr}_{G_C, \mu}^{(*)} |_{C_*}$.

Recall that $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} C_i}$ is a projective scheme over $\prod_{i \in I} C_i$, and the natural $L_I^+(G_C)$ -action on $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} C_i}$ factors through $L_I^n(G_C)$ for large enough n [32, Proposition 1.10]². Therefore $\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$ is a formal scheme that is formally of finite type and adic over $\prod_{i \in I} \mathbb{D}_i$, and its special fiber is projective

²While [32, Proposition 1.10] only treats the case of split G , it extends to the general case. Indeed, this is already implicitly used in [32, (12.10)].

over $\prod_{i \in I} v_i$. Also, the proof of [46, Lemma 3.2] shows that $\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is independent of the globalization from Lemma 1.1.

1.6. Recall that we have an isomorphism

$$\text{Gr}_{z, G}^k \xrightarrow{\sim} (\text{Gr}_{z, G}^1)^k$$

given by $((\mathcal{G}_j)_{j=1}^k, (\phi_j)_{j=1}^k) \mapsto ((\mathcal{G}_k, \phi_k), \dots, (\mathcal{G}_1, \phi_k \circ \dots \circ \phi_1))$.

Definition. Under this identification, write $\text{Gr}_{z, \text{SL}_h, m}^k$ for the closed subsheaf of $\text{Gr}_{z, \text{SL}_h}^k$ corresponding to $(\text{Gr}_{z, \text{SL}_h, m2\rho^\vee}^1)^k \subseteq (\text{Gr}_{z, \text{SL}_h}^1)^k$, where $2\rho^\vee$ denotes the sum of positive coroots in SL_h .

By 1.5, we see that $\text{Gr}_{z, \text{SL}_h, m}^k$ is a projective scheme over \mathbb{F}_q .

1.7. We conclude by showing that, after pulling back to the loop group, affine Schubert varieties are affine. Write $L_I(G_C)_{\mu_\bullet}|_{\prod_{i \in I} C_i}$ for the pullback of

$$\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$$

under the natural morphism $\prod_{j=1}^k L_I^{j, \circ}(G_C) \rightarrow \text{Gr}_{G_C}^{(I_1, \dots, I_k)}$.

Lemma. *Our $L_I(G_C)_{\mu_\bullet}|_{\prod_{i \in I} C_i}$ is affine over $\prod_{i \in I} C_i$.*

Proof. Because $\underline{\text{Aut}}(\mathcal{Y})/G_C$ is quasi-affine over C , the proof of [47, Proposition 1.2.6] shows that $\iota_* : \text{Gr}_{G_C}^{(I_1, \dots, I_k)} \rightarrow \text{Gr}_{\text{SL}_{h, C}}^{(I_1, \dots, I_k)}$ is a locally closed embedding. Now 1.5 indicates that $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ is a quasi-compact scheme, so [23, Lemma 5.4] implies that its image under ι_* lies in $\text{Gr}_{\text{SL}_{h, C}, (m2\rho^\vee)_{i \in I}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ for large enough m . Since $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ is projective over $\prod_{i \in I} C_i$ by 1.5 and ι_* is a monomorphism, we see that $\iota_* : \text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i} \rightarrow \text{Gr}_{\text{SL}_{h, C}, (m2\rho^\vee)_{i \in I}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ is a closed embedding. Combined with the fact that $L_I^+(G_C) \rightarrow L_I^+(\text{SL}_{r, C})$ is a closed embedding, this implies that $L_I(G_C)_{\mu_\bullet}|_{\prod_{i \in I} C_i} \rightarrow L_I(\text{SL}_{h, C})_{(m2\rho^\vee)_{i \in I}}|_{\prod_{i \in I} C_i}$ is also a closed embedding. Now the argument in the proof of [2, Lemma 4.23] shows that $L_I(\text{SL}_{h, C})_{(m2\rho^\vee)_{i \in I}}$ is affine over C^I , so $L_I(G_C)_{\mu_\bullet}|_{\prod_{i \in I} C_i}$ is affine over $\prod_{i \in I} C_i$. \square

2. FORMAL MODULI OF LOCAL SHTUKAS

To define the uniformization morphism via Beauville–Laszlo gluing in §5, we need a formal variant of the moduli of local shtukas. Moreover, to show that the uniformization morphism is étale, we need some finitude properties of this formal moduli. Accomplishing these tasks is the goal of this section.

We start by defining local shtukas and their quasi-isogenies in the formal setting. After proving a rigidity property for quasi-isogenies, we define the formal moduli problem, and we dedicate the rest of this section to proving that it gives a formal scheme that is locally formally of finite type over \mathbb{D}^I .

Our strategy ultimately harks back to Rapoport–Zink’s proof [36] of the analogous property for Rapoport–Zink spaces. The equicharacteristic incarnation of this argument is heavily based on work of Hartl–Viehmann [23] and Arasteh Rad–Hartl [2], although we generalize their results to the case of arbitrarily many legs.

2.1. Later, it will be useful to work in the following generality. Let R be a topological $\mathbb{F}_q[[\zeta_i]]_{i \in I}$ -algebra that is adic with finitely generated ideal of definition, and write $S := \mathrm{Spec} R$. Write $\tau : S \rightarrow S$ for the absolute q -Frobenius endomorphism. By abuse of notation, we also write $\tau : R[[z]] \rightarrow R[[z]]$ for the canonical lift of absolute q -Frobenius. We use ${}^\tau(-)$ to denote pullback by τ .

Write $R[[z, \frac{1}{z}]]$ for the completion of $R((z))$ with respect to the topology induced from R . We now define *local G -shtukas*.

Definition.

- a) A *local G -shtuka* over S consists of
- i) for all $1 \leq j \leq k$, a G -bundle \mathcal{G}_j on $\mathrm{Spec} R[[z]]$,
 - ii) for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\phi_j : \mathcal{G}_j|_{\mathrm{Spec} R[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}} \xrightarrow{\sim} \mathcal{G}_{j+1}|_{\mathrm{Spec} R[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}},$$

where \mathcal{G}_{k+1} denotes the G -bundle ${}^\tau \mathcal{G}_1$.

- b) Suppose that $\mathrm{Spf} R$ lies over $\prod_{i \in I} \mathbb{D}_i$, and let $\mathcal{G} = ((\mathcal{G}_j)_{j=1}^k, (\phi_j)_{j=1}^k)$ be a local shtuka over S . We say that \mathcal{G} is *bounded by μ_\bullet* if, for any affine étale cover $\mathrm{Spf} \tilde{R} \rightarrow \mathrm{Spf} R$ such that ${}^\tau \mathcal{G}_1|_{\mathrm{Spec} \tilde{R}[[z]]}$ is trivial and any trivialization $t : {}^\tau \mathcal{G}_1|_{\mathrm{Spec} \tilde{R}[[z]]} \xrightarrow{\sim} G$, the $\mathrm{Spf} \tilde{R}$ -point of $\widehat{\mathrm{Gr}}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ given by

$$\mathcal{G}_1|_{\mathrm{Spec} \tilde{R}[[z]]} \xrightarrow{(\phi_1)_{\tilde{R}[[z, \frac{1}{z}]}}} \dots \xrightarrow{(\phi_{k-1})_{\tilde{R}[[z, \frac{1}{z}]}}} \mathcal{G}_k|_{\mathrm{Spec} \tilde{R}[[z]]} \xrightarrow{(t \circ \phi_k)_{\tilde{R}[[z, \frac{1}{z}]}}} G$$

lies in $\widehat{\mathrm{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, using the description of $\widehat{\mathrm{Gr}}_G^{(I_1, \dots, I_k)}$ from Lemma 1.4.

It suffices to check Definition 2.1.b) for a single $\mathrm{Spf} \tilde{R} \rightarrow \mathrm{Spf} R$ and t .

2.2. For the rest of this section, assume that R is discrete, so that the ζ_i are nilpotent in R . In this setting, we use the following notion of quasi-isogenies.

Definition. Let \mathcal{G} and \mathcal{G}' be local G -shtukas over S .

- a) A *quasi-isogeny* from \mathcal{G} to \mathcal{G}' consists of, for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\delta_j : \mathcal{G}_j|_{\mathrm{Spec} R((z))} \xrightarrow{\sim} \mathcal{G}'_j|_{\mathrm{Spec} R((z))}$$

such that the diagram

$$\begin{array}{ccc} \mathcal{G}_j|_{\mathrm{Spec} R((z))} & \xrightarrow{\phi_j} & \mathcal{G}_{j+1}|_{\mathrm{Spec} R((z))} \\ \downarrow \delta_j & & \downarrow \delta_{j+1} \\ \mathcal{G}'_j|_{\mathrm{Spec} R((z))} & \xrightarrow{\phi'_j} & \mathcal{G}'_{j+1}|_{\mathrm{Spec} R((z))} \end{array}$$

commutes, where δ_{k+1} denotes the isomorphism ${}^\tau \delta_1$, and we use Lemma 1.3 to identify $R[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}$ with $R((z))$.

- b) Let m be a non-negative integer, and let δ be a quasi-isogeny from \mathcal{G} to \mathcal{G}' . We say that δ is *bounded by m* if, for all $1 \leq j \leq k$, the morphism $\iota_*(\delta_j)$ yields a point of $[L_z^+ \mathrm{SL}_h \backslash \mathrm{Gr}_{z, \mathrm{SL}_h, m2\rho}^1]$.

Since $L_z^+ G$ -bundles on $\mathrm{Spec} R$ are trivial after an étale cover, [23, Lemma 5.4] implies that any quasi-isogeny is bounded by m for large enough m .

2.3. We will need the following quantitative version of the rigidity of quasi-isogenies. Let J be an ideal of R satisfying $J^n = 0$, and write $j : \overline{S} \rightarrow S$ for the associated closed embedding.

Proposition. *For all local G -shtukas \mathcal{G} and \mathcal{G}' over S , pullback yields a bijection*

$$\{\text{quasi-isogenies from } \mathcal{G} \text{ to } \mathcal{G}'\} \xrightarrow{\sim} \{\text{quasi-isogenies from } j^*\mathcal{G} \text{ to } j^*\mathcal{G}'\}.$$

Moreover, suppose that S lies over $\prod_{i \in I} \mathbb{D}_i$ and that \mathcal{G} and \mathcal{G}' are bounded by μ_\bullet . There exists a non-negative integer B such that, if j^δ is bounded by m , then δ is bounded by $m + B \lceil \log_q n \rceil$.*

Proof. By induction, it suffices to consider the case where $n = q$. Then $\tau : S \rightarrow S$ factors as $j \circ \iota$ for a unique morphism $\iota : S \rightarrow \overline{S}$. For any quasi-isogeny δ from \mathcal{G} to \mathcal{G}' , note that ${}^\tau \delta_1 = \iota^* j^* \delta_1$. Therefore the commutative square

$$\begin{array}{ccc} \mathcal{G}_k|_{\text{Spec } R((z))} & \xrightarrow{\phi_k} & {}^\tau \mathcal{G}_1|_{\text{Spec } R((z))} \\ \downarrow \delta_k & & \downarrow {}^\tau \delta_1 \\ \mathcal{G}'_k|_{\text{Spec } R((z))} & \xrightarrow{\phi'_k} & {}^\tau \mathcal{G}'_1|_{\text{Spec } R((z))} \end{array},$$

enables us to recover δ_k from $j^* \delta_1$, where we use Lemma 1.3 to identify $R[[z]][\frac{1}{z-\zeta_i}]_{i \in I_k}$ with $R((z))$. From here, we similarly recover δ_j for $1 \leq j \leq k-1$, showing that pullback by j is injective on quasi-isogenies. By checking the analogous commutative squares over \overline{S} , this also implies that pullback by j is surjective on quasi-isogenies.

Next, suppose that S lies over $\prod_{i \in I} \mathbb{D}_i$ and that \mathcal{G} and \mathcal{G}' are bounded by μ_\bullet . If $j^*\delta$ is bounded by m , then its pullback $\iota^* j^* \delta_1 = {}^\tau \delta_1$ is as well. Because ϕ_k and ϕ'_k are bounded by $(\mu_i)_{i \in I_k}$, where the relative position bound is taken with respect to the $(z - \zeta_i)$ for i in I_k , a quasi-compactness argument shows that there exists a non-negative integer B such that δ_k is bounded by $m + B$. For $1 \leq j \leq k-1$, applying the same argument to δ_j indicates that, after increasing B by an amount depending only on μ_\bullet , our δ_j is also bounded by $m + B$. \square

2.4. We now define the formal moduli of local G -shtukas.

Definition. Write $\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}$ for the sheaf over \mathbb{D}^I whose S -points parametrize data consisting of

- i) a local G -shtuka \mathcal{G} over S ,
- ii) a quasi-isogeny δ from \mathcal{G} to the trivial local G -shtuka $G = ((G)_{j=1}^k, (\text{id})_{j=1}^k)$.

Write $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ for the subsheaf of $\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ whose S -points consist of the (\mathcal{G}, δ) such that \mathcal{G} is bounded by μ_\bullet .

Write $f^\mathfrak{L} : \mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \prod_{i \in I} \mathbb{D}_i$ for the structure morphism.

2.5. **Proposition.** *Our $\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}$ is naturally isomorphic to $\text{Gr}_{z, G}^k|_{\mathbb{D}^I}$ over \mathbb{D}^I , and $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a closed subsheaf of $\widehat{\text{Gr}}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$.*

Proof. In Definition 2.4, note that i) and ii) are uniquely determined by $(\mathcal{G}_j)_{j=1}^k$, $(\phi_j)_{j=1}^{k-1}$, and δ_k . This is precisely the data parametrized by $\text{Gr}_{z, G}^k|_{\mathbb{D}^I}$, which proves the first claim. The second claim follows from the fact that $\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a closed subsheaf of $\widehat{\text{Gr}}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$; see the proof of [2, Proposition 4.11]. \square

2.6. First, we naively stratify $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ by bounding the quasi-isogeny. Write $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ for the subsheaf of $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ whose S -points consist of the (\mathcal{G}, δ) such that δ is bounded by m .

Proposition. *Our $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a formal scheme that is formally of finite type and adic over $\prod_{i \in I} \mathbb{D}_i$, its reduced subscheme is projective over $\prod_{i \in I} v_i$, and $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ equals the direct limit $\varinjlim_m \mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$.*

Proof. Note that we have a Cartesian square

$$\begin{array}{ccc} \mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i} & \longrightarrow & \mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \\ \downarrow \iota_* & & \downarrow \iota_* \\ \mathrm{Gr}_{z,\mathrm{SL}_h,m}^k|_{\mathbb{D}^I} & \longrightarrow & \mathrm{Gr}_{z,\mathrm{SL}_h}^k|_{\mathbb{D}^I}, \end{array}$$

where we use Proposition 2.5 to identify $\mathfrak{LocSht}_{\mathrm{SL}_h}^{(I_1,\dots,I_k)}$ with $\mathrm{Gr}_{z,\mathrm{SL}_h}^k|_{\mathbb{D}^I}$. Because SL_h/G is quasi-affine over \mathcal{O}_F , Proposition 2.5 and [47, Proposition 1.2.6] show that $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \mathrm{Gr}_{z,\mathrm{SL}_h}^k|_{\prod_{i \in I} \mathbb{D}_i}$ is a closed embedding. Therefore its pullback $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \mathrm{Gr}_{z,\mathrm{SL}_h,m}^k|_{\prod_{i \in I} \mathbb{D}_i}$ is as well. Since $\mathrm{Gr}_{z,\mathrm{SL}_h,m}^k|_{\prod_{i \in I} nv_i}$ is projective over $\prod_{i \in I} nv_i$ for any positive integer n by 1.6, the same holds for $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} nv_i}$. Now the underlying topological space of $\mathrm{Gr}_{z,\mathrm{SL}_h,m}^k|_{\prod_{i \in I} nv_i}$ is independent of n , so the $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} nv_i}$ have this property too. From here, [17, (1, 10.6.4)] indicates that $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a noetherian formal scheme that is adic over $\prod_{i \in I} \mathbb{D}_i$. Hence its reduced subscheme equals that of $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} v_i}$, which is projective over $\prod_{i \in I} v_i$, so $\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$. Finally, last statement follows from $\mathrm{Gr}_{z,\mathrm{SL}_h}^k$ equaling the direct limit $\varinjlim_m \mathrm{Gr}_{z,\mathrm{SL}_h,m}^k$. \square

2.7. To obtain a more refined stratification of $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, we need the following algebraization lemma. Briefly, relax our assumption that R is discrete, since we plan to use this lemma later as well. Let $(S_l)_{l \geq 0}$ be a direct system of affine schemes $S_l = \mathrm{Spec} R_l$ over $\prod_{i \in I} \mathbb{D}_i$ such that

- i) the morphisms $S_l \rightarrow S_{l'}$ are closed embeddings,
- ii) the associated ideals $\ker(R_{l'} \rightarrow R_l)$ are nilpotent.

Take R to be the ring $\varprojlim_l R_l$, and endow R with a topological ring structure such that $\mathbb{F}_q[[\zeta_i]]_{i \in I} \rightarrow R$ is continuous, the $R \rightarrow R_l$ are continuous for the discrete topology on R_l , and R is adic with finitely generated ideal of definition.

Lemma. *Pullback yields an equivalence of groupoids*

$$\left\{ \begin{array}{l} \text{local } G\text{-shtukas over} \\ S \text{ bounded by } \mu_\bullet \end{array} \right\} \xrightarrow{\sim} \varprojlim_l \left\{ \begin{array}{l} \text{local } G\text{-shtukas over} \\ S_l \text{ bounded by } \mu_\bullet \end{array} \right\}.$$

Proof. Let $(\mathcal{G}^l)_{l \geq 0}$ be a compatible system of local G -shtukas over S_l bounded by μ_\bullet . We can form the G -bundles $\mathcal{G}_j := \varprojlim_l \mathcal{G}_j^l$ on $\mathrm{Spec} R[[z]]$, so now we just need to form the isomorphisms ϕ_j .

Let $\text{Spec } \tilde{R}_0 \rightarrow S_0$ be an affine étale cover where $\mathcal{G}_j^0|_{\text{Spec } \tilde{R}_0[[z]]}$ is trivial for all $1 \leq j \leq k$, and fix trivializations of the $\mathcal{G}_j^0|_{\text{Spec } \tilde{R}_0[[z]]}$. By ii), there exists a unique affine étale cover $\text{Spec } \tilde{R}_l \rightarrow S_l$ whose pullback to S_0 is $\text{Spec } \tilde{R}_0$, and there also exist compatible systems of trivializations of the $\mathcal{G}_j^l|_{\text{Spec } \tilde{R}_l[[z]]}$ [23, Proposition 2.2(c)]³. Under these identifications, the $(\phi_j^l)_{\tilde{R}_l((z))}$ correspond to compatible systems of b_j^l in $G(\widehat{\mathcal{O}}_C^{j,\circ}(\text{Spec } \tilde{R}_l))$, where we use Lemma 1.3 to identify $\tilde{R}_l((z))$ with $\widehat{\mathcal{O}}_C^{j,\circ}(\text{Spec } \tilde{R}_l)$.

For all i in I , let V_i be an affine neighborhood of v_i in C_i . Because the \mathcal{G}^l are bounded by μ_\bullet , our $(b_j^l)_{j=1}^k$ yield \tilde{R}_l -points of $L_I(G_C)_{\mu_\bullet}|_{\prod_{i \in I} V_i}$. The latter is affine by Lemma 1.7, so the compatible system of $(b_j^l)_{j=1}^k$ yields an $\tilde{R} := \varprojlim_l \tilde{R}_l$ -point $(b_j)_{j=1}^k$ of $L_I(G_C)|_{\prod_{i \in I} V_i}$. By construction, the resulting local G -shtuka $\mathcal{G} := ((\mathcal{G}_j|_{\text{Spec } \tilde{R}[[z]]})_{j=1}^k, (b_j)_{j=1}^k)$ over $\text{Spec } \tilde{R}$ is bounded by μ_\bullet . Since the $(\phi_j^l)_{\tilde{R}_l((z))}$ and thus b_j^l are compatible with the descent data of \mathcal{G}_j^l from $\text{Spec } \tilde{R}_l$ to S_l , we see that the b_j are compatible with the descent data of \mathcal{G}_j from $\text{Spec } \tilde{R}$ to S . Hence \mathcal{G} naturally descends to a local G -shtuka \mathcal{G} over S bounded by μ_\bullet , as desired. \square

2.8. Resume our assumption that R is discrete. The following refined stratification of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ has better closure properties under formal completion. Write $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ for the formal completion of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ along the reduced subscheme of $\mathfrak{LocSht}_{G, \mu_\bullet, m}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$.

Lemma. *Our $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a formal scheme that is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$.*

Proof. Proposition 2.6 and [23, Lemma 5.4] imply that $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ equals the direct limit $\varinjlim_l \mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, where $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ denotes the formal completion of $\mathfrak{LocSht}_{G, \mu_\bullet, m+l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ along the reduced subscheme of $\mathfrak{LocSht}_{G, \mu_\bullet, m}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$. The reduced subscheme of $\mathfrak{LocSht}_{G, \mu_\bullet, m}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is quasi-compact by Proposition 2.6, so it is covered by finitely many affine open subschemes U . Proposition 2.6 indicates that $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a noetherian formal scheme with reduced subscheme equal to that of $\mathfrak{LocSht}_{G, \mu_\bullet, m}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, so we can form the affine open formal subscheme $\mathfrak{U}_l = \text{Spf } A_l$ of $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ with underlying topological space U .

The above shows that $\varinjlim_l \mathfrak{U}_l$ is an open subsheaf of $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$. Thus it suffices to prove that $\varinjlim_l \mathfrak{U}_l$ is an affine formal scheme that is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$. Because the

$$\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l'}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$$

are closed embeddings, the $A_{l'} \rightarrow A_l$ are surjective. Write $A := \varprojlim_l A_l$. Write J_0 for the largest ideal of definition of A_0 , and write J for its preimage in A .

For any positive integer c , we claim that $A_{l'}/J^c \rightarrow A_l/J^c$ is an isomorphism for large enough l and l' . Note that the $A_{l'}/J^c \rightarrow A_l/J^c$ have nilpotent kernels, and the Mittag-Leffler criterion implies that $A/J^c = \varprojlim_l A_l/J^c$. Endow A/J^c with the

³While [23] only treats split reductive G , the proof immediately adapts to any smooth G .

discrete topology. Because the ζ_i vanish in $A/J = A_0/J_0$, we see that the ζ_i are nilpotent in A/J^c . Thus $\mathbb{F}_q[[\zeta_i]]_{i \in I} \rightarrow A/J^c$ is continuous, so we can apply Lemma 2.7 to the local G -shtukas \mathcal{G}^l over $\text{Spec } A_l/J^c$ obtained from the morphism

$$\text{Spec } A_l/J^c \rightarrow \text{Spf } A_l \rightarrow \mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$$

to get a local G -shtuka \mathcal{G} over $\text{Spec } A/J^c$ bounded by μ_\bullet . Next, consider the quasi-isogeny δ^0 obtained from $\text{Spec } A_0/J_0 \rightarrow \mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, 0}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$. Proposition 2.3 uniquely lifts δ^0 to a quasi-isogeny δ from \mathcal{G} to G , which implies that the resulting A/J^c -point (\mathcal{G}, δ) of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$ lies in $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$. Therefore [23, Lemma 5.4] indicates that (\mathcal{G}, δ) lies in $\mathfrak{LocSht}_{G, \mu_\bullet, \widehat{m}, l}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i}$ for large enough l . Pulling back to $\text{Spec } A_0/J_0$ shows that (\mathcal{G}, δ) even lies in $\text{Spf } A_l$. The uniqueness of Proposition 2.3 implies that the pullback of (\mathcal{G}, δ) to $\text{Spec } A_{l'}/J^c$ equals $(\mathcal{G}^{l'}, \delta^{l'})$, so $A_{l'} \rightarrow A_l \rightarrow A/J^c \rightarrow A_{l'}/J^c$ equals the quotient map. Quotienting by the image of J^c in A_l shows that $A_{l'}/J^c \rightarrow A_l/J^c$ is an isomorphism, which concludes our proof of the claim.

Write $\mathfrak{a}_l := \ker(A \rightarrow A_l)$. The claim indicates that the ideals $\mathfrak{a}_l + J^c$ of A stabilize for any positive integer c , and because the A_l are noetherian, we see that the $\text{im}(J/J^2 \rightarrow A_l/J^2) = J/(J^2 + \mathfrak{a}_l)$ are finite over A . Therefore [36, proposition (2.5)] shows that A with the inverse limit topology is noetherian and J -adic, which implies that $\varinjlim_l \mathfrak{a}_l = \text{Spf } A$. Finally, the reduced subscheme of $\text{Spf } A$ is of finite type over $\prod_{i \in I} v_i$ by Proposition 2.6, so $\text{Spf } A$ is formally of finite type over \mathbb{D}_i . \square

2.9. We can use the quasi-isogeny to define the following distance function.

Definition. Let K be a field over \mathbb{F}_q , and let $x = (\mathcal{G}, \delta)$ and $x' = (\mathcal{G}', \delta')$ be K -points of $\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}$. Write $d(x, x')$ for the smallest non-negative integer m such that the quasi-isogeny $\delta^{-1} \circ \delta'$ of local G -shtukas over $\text{Spec } K((z))$ is bounded by m .

2.10. **Lemma.** *As K runs over all fields over \mathbb{F}_q , the maps d induce a metric on the underlying set $|\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}|$. For any x in $|\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}|$ and non-negative integer r , the associated closed ball $B_r(x)$ of radius r centered at x is closed with respect to the Zariski topology on $|\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}|$.*

Proof. We immediately see that d is insensitive to field extensions, so d induces a map $|\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}| \times |\mathfrak{LocSht}_G^{(I_1, \dots, I_k)}| \rightarrow \mathbb{Z}_{\geq 0}$. Since relative position bounds along the same divisor are sub-additive under composition, d satisfies the triangle inequality, and because $2\rho^\vee$ is fixed by the Chevalley involution, d is symmetric. Next, if $d(x, x') = 0$, then $\iota_*(\delta_j^{-1} \circ \delta'_j)$ extends to an isomorphism of SL_r -bundles on $\text{Spec } K[[z]]$ for all $1 \leq j \leq k$. Since ι is a monomorphism, this implies that the $\delta_j^{-1} \circ \delta'_j$ extend to isomorphisms of G -bundles on $\text{Spec } K[[z]]$, so $x = x'$. For the last statement, note that $B_r(x)$ equals, on the level of topological spaces, the preimage of the closed substack $[L_z^+ \text{SL}_h \backslash \text{Gr}_{z, \text{SL}_h, r2\rho^\vee}^1]^k$ under the morphism

$$\mathfrak{LocSht}_G^{(I_1, \dots, I_k)} \rightarrow [L_z^+ \text{SL}_h \backslash \text{Gr}_{z, \text{SL}_h}^1]^k$$

given by $(\mathcal{G}', \delta') \mapsto (\iota_*(\delta_j^{-1} \circ \delta'_j))_{j=1}^k$. \square

2.11. All points of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ are close enough to one defined over a fixed finite field in the following sense.

Lemma. *There exists a finite extension $\mathbb{F}_{q'}$ of \mathbb{F}_q and a non-negative integer D such that, for every x in $|\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$, there exists an $\mathbb{F}_{q'}$ -point y of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ satisfying $d(x, y) \leq D$.*

Proof. Suppose that x corresponds to a K -point (\mathcal{G}, δ) , where we can assume that K is an algebraically closed field over \mathbb{F}_q . Then \mathcal{G}_j is trivial for all $1 \leq j \leq k$, and after fixing trivializations of the \mathcal{G}_j , our δ_j correspond to g_j in $G(K((z)))$. The commutativity of the diagram

$$\begin{array}{ccccc} \mathcal{G}_1|_{\mathrm{Spec} K((z))} & \xrightarrow{-\phi_1} & \cdots & \xrightarrow{-\phi_{k-1}} & \mathcal{G}_k|_{\mathrm{Spec} K((z))} & \xrightarrow{-\phi_k} & \tau \mathcal{G}_1|_{\mathrm{Spec} K((z))} \\ \downarrow \delta_1 & & & & \downarrow \delta_k & & \downarrow \tau \delta_1 \\ G & \xlongequal{\quad} & \cdots & \xlongequal{\quad} & G & \xlongequal{\quad} & G \end{array}$$

implies that $\tau \delta_1^{-1} \circ \delta_1 = \phi_k \circ \cdots \circ \phi_1$, so the image of $\tau(g_1)^{-1}g_1$ in $\mathrm{Gr}_{z, G}^1|_{v_*}$ lies in $\mathrm{Gr}_{z, G, \sum_{i \in I} \mu_i}^1|_{v_*}$. Now 1.5 indicates that $\mathrm{Gr}_{z, G, \sum_{i \in I} \mu_i}^1|_{v_*}$ is a quasi-compact scheme, so [23, Lemma 5.4] shows that its image under ι_* lies in $\mathrm{Gr}_{z, \mathrm{SL}_h, m}^1$ for large enough m . Therefore [33, 2.2.1 (ii)] and [37, (2.1)] yield a non-negative integer D such that, for all such g_1 , there exists h_1 in $G(\mathbb{F}_q((z)))$ such that the image of $g_1 h_1^{-1}$ in $\mathrm{Gr}_{z, \mathrm{SL}_h}^1$ lies in $\mathrm{Gr}_{z, \mathrm{SL}_h, D2\rho^v}^1$.

If $\sum_{i \in I} \mu_i$ is not a coroot, then $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is empty, and the result vacuously holds. So assume that $\sum_{i \in I} \mu_i$ is a coroot. Then the image of 1 in $\mathrm{Gr}_G^{(I)}|_{\prod_{i \in I} v_i}$ lies in $\mathrm{Gr}_{G, \sum_{i \in I} \mu_i}^{(I)}|_{\prod_{i \in I} v_i}$. Since the convolution morphism $\mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} v_i} \rightarrow \mathrm{Gr}_{G, \sum_{i \in I} \mu_i}^{(I)}|_{\prod_{i \in I} v_i}$ is of finite type by 1.5 and surjective, its fiber at 1 has an $\mathbb{F}_{q'}$ -point b for some finite extension $\mathbb{F}_{q'}$ of \mathbb{F}_q . Next, identify $\mathrm{Gr}_G^{(I_1, \dots, I_k)}|_{v^I}$ with $\mathrm{Gr}_{z, G}^k|_{v^I}$. Because the fiber of $(L_z G)^k \rightarrow \mathrm{Gr}_G^k|_{\prod_{i \in I} v_i}$ at b is an $(L_z^+ G)^k$ -bundle on $\mathrm{Spec} \mathbb{F}_{q'}$, Lang's lemma indicates that it has an $\mathbb{F}_{q'}$ -point $(b_j)_{j=1}^k$. By construction, the local G -shtuka $\mathcal{H} := ((G)_{j=1}^k, (b_j)_{j=1}^k)$ over $\mathrm{Spec} \mathbb{F}_{q'}$ is bounded by μ_\bullet , and $b_k \cdots b_1$ equals 1 up to right $G(\mathbb{F}_{q'}[[z]])$ -translation. By replacing b_1 with a right $G(\mathbb{F}_{q'}[[z]])$ -translate, we can assume that $b_k \cdots b_1 = 1$. Combined with the fact that $h_1 = \tau(h_1)$, this shows that the diagram

$$\begin{array}{ccccc} G & \xrightarrow{-b_1} & \cdots & \xrightarrow{-b_{k-1}} & G & \xrightarrow{-b_k} & G \\ \downarrow h_1 & & & & \downarrow h_k & & \downarrow \tau h_1 \\ G & \xlongequal{\quad} & \cdots & \xlongequal{\quad} & G & \xlongequal{\quad} & G \end{array}$$

commutes for uniquely determined h_2, \dots, h_k in $G(\mathbb{F}_{q'}((z)))$. Since b_j and ϕ_j are bounded by $\sum_{i \in I_j} \mu_j$ for $1 \leq j \leq k-1$, where the relative position bound is taken with respect to z , a quasi-compactness argument as before shows that, after increasing D by an amount depending only on μ_\bullet , the image of $g_j h_j^{-1}$ in $\mathrm{Gr}_{z, \mathrm{SL}_h}^1$ lies in $\mathrm{Gr}_{z, \mathrm{SL}_h, D2\rho^v}^1$. Therefore the quasi-isogeny $h := (h_j)_{j=1}^k$ from \mathcal{H} to G yields an $\mathbb{F}_{q'}$ -point $y := (\mathcal{H}, h)$ of $\mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ with $d(x, y) \leq D$, as desired. \square

2.12. The following theorem is the main result of this section. Write $B_r(x)_{\mu_\bullet}$ for the intersection of $B_r(x)$ and $|\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$, and write $\mathbf{1}$ for the \mathbb{F}_q -point $(G, (\text{id})_{j=1}^k)$ of $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}$. Note that $B_m(\mathbf{1})_{\mu_\bullet}$ equals $|\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$.

Theorem. *Our $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a formal scheme that is locally formally of finite type over $\prod_{i \in I} \mathbb{D}_i$.*

Proof. Let $\mathbb{F}_{q'}$ and D be as in Lemma 2.11. Write Z_m^s for the union

$$\bigcup_y B_D(y)_{\mu_\bullet} \cap B_m(\mathbf{1})_{\mu_\bullet},$$

where y runs over $\mathbb{F}_{q'}$ -points of $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ satisfying $d(\mathbf{1}, y) \geq s$. The triangle inequality implies that it suffices to take y also satisfying $d(\mathbf{1}, y) \leq m + D$. Because $B_{m+D}(\mathbf{1})_{\mu_\bullet}$ equals $|\mathfrak{LocSht}_{G,\mu_\bullet,m+D}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$, Proposition 2.6 implies that there are finitely many such y . Hence Lemma 2.10 indicates that Z_m^s is a finite union of Zariski closed subsets of $|\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$.

Write \mathfrak{U}_m^s for the open formal subscheme of $\mathfrak{LocSht}_{G,\mu_\bullet,\widehat{m}}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ with underlying topological space given by the complement of Z_m^s . By Lemma 2.8, \mathfrak{U}_m^s is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$. Note that $\mathfrak{LocSht}_{G,\mu_\bullet,\widehat{m}}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ equals the formal completion of $\mathfrak{LocSht}_{G,\mu_\bullet,m+1}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ along the reduced subscheme of $\mathfrak{LocSht}_{G,\mu_\bullet,\widehat{m}}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, so \mathfrak{U}_{m+1}^s equals the formal completion of \mathfrak{U}_m^s along the reduced subscheme of \mathfrak{U}_m^s .

For any non-negative integer s , we claim that \mathfrak{U}_m^s stabilizes. The above indicates that it suffices to check this on underlying sets, so suppose that there exists x in $|\mathfrak{U}_{m+1}^s| \setminus |\mathfrak{U}_m^s|$. Lemma 2.11 yields an $\mathbb{F}_{q'}$ -point y of $\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ satisfying $d(x, y) \leq D$. As x does not lie in Z_{m+1}^s , we have $d(\mathbf{1}, y) < s$, so the triangle inequality yields $m + 1 = d(\mathbf{1}, x) < s + D$. Hence \mathfrak{U}_m^s stabilizes for $m \geq s + D - 1$, which concludes our proof of the claim.

Set $\mathfrak{U}^s := \varinjlim_m \mathfrak{U}_m^s$. Proposition 2.6 implies that \mathfrak{U}^s is an open subsheaf of

$$\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}.$$

The claim shows that \mathfrak{U}^s equals \mathfrak{U}_m^s for large enough m , so \mathfrak{U}^s is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$. Now we just need to prove $\varinjlim_s \mathfrak{U}^s = \mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$. It suffices to check this on underlying sets, so take x in $|\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$. Proposition 2.6 indicates that x lies in

$$|\mathfrak{LocSht}_{G,\mu_\bullet,m}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$$

for large enough m , so for all y in $|\mathfrak{LocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}|$ such that x lies in $B_D(y)_{\mu_\bullet}$, the triangle inequality yields $d(\mathbf{1}, y) \leq m + D$. Therefore x lies in $|\mathfrak{U}^{m+D+1}|$. \square

2.13. Using representations of the dual group, we can index relative position bounds as follows. Let \widetilde{F} be the finite Galois extension of F such that $\text{Gal}(\widetilde{F}/F)$ equals the image of the Γ_F -action on $X_*^+(T)$. Write $\widetilde{\mathbb{D}}$ for $\text{Spd } \mathcal{O}_{\widetilde{F}}$. Let E be a finite extension of $\mathbb{Q}_\ell(\sqrt{q})$, write \widehat{G} for the dual group of G_F over \mathcal{O}_E , and write ${}^L G$ for $\widehat{G} \rtimes \text{Gal}(\widetilde{F}/F)$.

Let V be an object of $\text{Rep}_E({}^L G)^I$. Note that $\coprod_{\mu_\bullet} \mathfrak{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\mathbb{D}^I}$ naturally descends to a sheaf $\mathfrak{LocSht}_{G, V}^{(I_1, \dots, I_k)}$ over \mathbb{D}^I , where μ_\bullet runs over highest weights appearing in $V_{\overline{\mathbb{Q}}_\ell}|_{\widehat{G}^I}$. Theorem 2.12 and descent imply that $\mathfrak{LocSht}_{G, V}^{(I_1, \dots, I_k)}$ is a formal scheme that is locally formally of finite type over \mathbb{D}^I .

2.14. Finally, we define partial Frobenii for the formal moduli of local G -shtukas.

Definition. Write $\mathfrak{Fr}^{(I_1, \dots, I_k)} : \mathfrak{LocSht}_{G, V}^{(I_1, \dots, I_k)} \rightarrow \mathfrak{LocSht}_{G, V}^{(I_2, \dots, I_k, I_1)}$ for the morphism given by sending

$$\begin{array}{ccccccc} \mathcal{G}_1 & \xrightarrow{\phi_1} & \cdots & \xrightarrow{\phi_{k-1}} & \mathcal{G}_k & \xrightarrow{\phi_k} & \tau \mathcal{G}_1 & & \mathcal{G}_2 & \xrightarrow{\phi_2} & \cdots & \xrightarrow{\phi_k} & \tau \mathcal{G}_1 & \xrightarrow{\phi_1} & \tau \mathcal{G}_2 \\ | & & & & | & & | & \text{to} & | & & & & | & & | \\ \downarrow \delta_1 & & & & \downarrow \delta_k & & \downarrow \delta_1 & & \downarrow \delta_2 & & & & \downarrow \delta_1 & & \downarrow \delta_2 \\ G & \xlongequal{\quad} & \cdots & \xlongequal{\quad} & G & \xlongequal{\quad} & G & & G & \xlongequal{\quad} & \cdots & \xlongequal{\quad} & G & \xlongequal{\quad} & G \end{array}$$

Note that $\mathfrak{Fr}^{(I_1, \dots, I_k)}$ lies above the endomorphism of \mathbb{D}^I given by geometric q -Frobenius on the i -th factor for i in I_1 and the identity on all other factors.

3. RELATIVE z -ADIC HODGE THEORY

The local shtukas defined in §2 are (formal) algebraic, while the local shtukas used by Fargues–Scholze [11] are (non-archimedean) analytic in nature. To compare them, we need an equicharacteristic version of Kedlaya–Liu’s results [31] on relative p -adic Hodge theory. Our goal in this section is to prove the necessary results on *relative z -adic Hodge theory*, in the spirit of work of Hartl [22].

We begin by recalling the equicharacteristic version of Fontaine’s period ring A_{inf} . Using a result of Anschütz [1], we prove an algebraization theorem for G -bundles on A_{inf} , at least pro-étale locally on the base. Finally, we relate $G(\mathcal{O}_F)$ -local systems to G -bundles on the equicharacteristic version of the (relative integral) Robba ring equipped with a Frobenius automorphism.

Our arguments closely follow those of Kedlaya–Liu [31] and Scholze–Weinstein [42]. However, we have streamlined and simplified the presentation, both because we only prove what we need as well as because the arithmetic of formal power series is easier than that of Witt vectors.

3.1. Let $S = \text{Spa}(R, R^+)$ be an affinoid perfectoid space over \mathbb{F}_q , and choose a pseudouniformizer ϖ of R . Write \mathcal{Y}_S for the complement of the vanishing locus of ϖ and z in $\text{Spa } R^+[[z]]$, and note that \mathcal{Y}_S is the analytic locus of the pre-adic space $\text{Spa } R^+[[z]]$. We have a continuous map $\text{rad} : |\mathcal{Y}_S| \rightarrow [0, \infty]$ given by

$$x \mapsto \frac{\log |\varpi(\tilde{x})|}{\log |z(\tilde{x})|},$$

where \tilde{x} denotes the unique rank-1 generalization of x in \mathcal{Y}_S . For any closed interval \mathcal{I} in $[0, \infty]$ with rational endpoints, write $\mathcal{Y}_{S, \mathcal{I}} = \text{Spa}(B_{S, \mathcal{I}}, B_{S, \mathcal{I}}^+)$ for the associated rational open subspace of $\text{Spa } R^+[[z]]$, which lies in \mathcal{Y}_S . More generally, for any subset \mathcal{I} of $[0, \infty]$, write $\mathcal{Y}_{S, \mathcal{I}}$ for the open subspace $\bigcup_{\mathcal{I}'} \mathcal{Y}_{S, \mathcal{I}'}$ of \mathcal{Y}_S , where \mathcal{I}' runs over closed intervals in \mathcal{I} with rational endpoints. Note that $\mathcal{Y}_{S, \mathcal{I}} \subseteq \text{rad}^{-1}(\mathcal{I})$. We see that $\mathcal{Y}_{S, [0, \infty)}$ and $\mathcal{Y}_{S, (0, \infty)}$ are naturally isomorphic to $\mathbb{D} \times S$ and $\text{Spa } F \times S$, respectively.

Write $\tau : S \rightarrow S$ for the absolute q -Frobenius automorphism, and by abuse of notation, write $\tau : R[[z]] \rightarrow R[[z]]$ for the canonical lift of absolute q -Frobenius. Note that $\text{rad} \circ \tau = q \cdot \text{rad}$. Finally, write X_S for the quotient $\mathcal{Y}_{S,(0,\infty)}/\tau^{\mathbb{Z}}$.

3.2. When \mathcal{I} contains ∞ , we can describe $B_{S,\mathcal{I}}$ using the following lemma. For any positive r in $\mathbb{Z}[\frac{1}{p}]$, write $R^+[[z, \frac{\varpi^r}{z}]]$ for the ϖ -adic completion of $R^+[[z]][\frac{\varpi^r}{z}]$.

Lemma. *We can identify*

$$R^+[[z, \frac{\varpi^r}{z}]] = \left\{ \sum_{m=-\infty}^{\infty} a_m z^m \mid \text{the } a_m \in R^+ \text{ and } \lim_{m \rightarrow -\infty} a_m \varpi^{rm} = 0 \right\}.$$

If we give $R^+[[z, \frac{\varpi^r}{z}]]$ the (ϖ, z) -adic topology, then $B_{S,[1/r,\infty]}$ equals $R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]$.

Proof. The above description of $R^+[[z, \frac{\varpi^r}{z}]]$ follows immediately from the definition. This description shows that $R^+[[z, \frac{\varpi^r}{z}]]$ is z -adically complete as a ring, so $R^+[[z, \frac{\varpi^r}{z}]]$ equals the (ϖ, z) -adic completion of $R^+[[z]][\frac{\varpi^r}{z}]$ as rings. Since $\mathcal{Y}_{S,[1/r,\infty]}$ equals the rational open subspace $\{|\varpi^r| \leq |z| \neq 0\}$ of $\text{Spa } R^+[[z]]$, this identifies $B_{S,[1/r,\infty]}$ with $R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]$ if we give $R^+[[z, \frac{\varpi^r}{z}]]$ the (ϖ, z) -adic topology. \square

3.3. Sometimes, it will be convenient to ignore the topology induced from R as follows. Write $A'(R^+)$ for $R^+[[z]]$ with the z -adic topology.

Lemma. *Our $\text{Spa}(A'(R^+)[\frac{1}{z}], A'(R^+))$ is a sousperfectoid adic space.*

Proof. The natural map $A'(R^+)[\frac{1}{z}] \rightarrow R^+[[z^{\pm 1/p^\infty}]]$ is a split injection of topological $A'(R^+)[\frac{1}{z}]$ -modules, where we give $R^+[[z^{\pm 1/p^\infty}]]$ the z -adic topology. \square

3.4. **Proposition.** *Our \mathcal{Y}_S is a sousperfectoid adic space.*

Proof. Note that \mathcal{Y}_S is covered by $\mathcal{Y}_{S,[0,\infty)}$ and $\mathcal{Y}_{S,[1,\infty)}$. Now $\mathcal{Y}_{S,[0,\infty)}$ is a sousperfectoid adic space by [11, Proposition II.1.1], so it suffices to prove that $\mathcal{Y}_{S,[1,\infty)}$ is a sousperfectoid adic space. By Proposition 3.2, $B_{S,[1,\infty)}$ equals $R^+[[z, \frac{\varpi}{z}]][\frac{1}{z}]$, where $R^+[[z, \frac{\varpi}{z}]]$ has the (ϖ, z) -adic topology.

Now z divides ϖ in $R^+[[z]][\frac{\varpi}{z}]$, so the (ϖ, z) -adic topology on $R^+[[z]][\frac{\varpi}{z}]$ equals the z -adic topology. This enables us to identify $\mathcal{Y}_{S,[1,\infty)}$ with the rational open subspace $\{|\varpi| \leq |z| \neq 0\}$ of $\text{Spa}(A'(R^+)[\frac{1}{z}], A'(R^+))$. The latter is sousperfectoid by Lemma 3.3, so $\mathcal{Y}_{S,[1,\infty)}$ is as well. \square

3.5. Since a power of ϖ divides a power of z in $R^+[[z]][\frac{z}{\varpi^r}]$, the (ϖ, z) -adic topology on $R^+[[z]][\frac{z}{\varpi^r}]$ equals the ϖ -adic topology. Therefore $B_{S,[0,1/r]}$ equals the Tate algebra $R\langle \frac{z}{\varpi^r} \rangle$. This argument lets us similarly identify

$$B_{S,[1,1]} = \left\{ \sum_{m=-\infty}^{\infty} a_m z^m \mid \text{the } a_m \in R \text{ and } \lim_{m \rightarrow \pm\infty} a_m \varpi^m = 0 \right\}.$$

We will use the following result with the Tannakian description of G -bundles.

Proposition. *Pullback yields a fully faithful functor*

$$\{\text{vector bundles on } \text{Spec } R^+[[z]]\} \hookrightarrow \{\text{vector bundles on } \mathcal{Y}_S\}.$$

Proof. Let $f : M \rightarrow M'$ be a map of finite projective $R^+[[z]]$ -modules, and consider its pullback g to \mathcal{Y}_S . Now Proposition 3.4 and [31, Theorem 2.7.7] indicate that $g|_{\mathcal{Y}_{S,[0,1]}}$, $g|_{\mathcal{Y}_{S,[1,\infty]}}$, and $g|_{\mathcal{Y}_{S,[1,1]}}$ correspond to maps of finite projective modules over $B_{S,[0,1]}$, $B_{S,[1,\infty]}$, and $B_{S,[1,1]}$, respectively, which are given by tensoring with f over $R^+[[z]]$. Lemma 3.2 indicates that $B_{S,[1,\infty]}$ equals $R^+[[z, \frac{\varpi}{z}]][\frac{1}{z}]$ as rings, so we see that $B_{S,[0,1]}$ and $B_{S,[1,\infty]}$ inject into $B_{S,[1,1]}$. Note that their intersection equals $R^+[[z]]$. Therefore the flatness of M yields a Cartesian square

$$\begin{array}{ccc} M & \hookrightarrow & M \otimes_{R^+[[z]]} B_{S,[0,1]} \\ \downarrow & & \downarrow \\ M \otimes_{R^+[[z]]} B_{S,[1,\infty]} & \hookrightarrow & M \otimes_{R^+[[z]]} B_{S,[1,1]}, \end{array}$$

and the same holds for M' . In particular, we recover f as the restriction of $g|_{\mathcal{Y}_{S,[0,1]}}$ (or of $g|_{\mathcal{Y}_{S,[1,\infty]}}$) to the intersection of $M \otimes_{R^+[[z]]} B_{S,[0,1]}$ and $M \otimes_{R^+[[z]]} B_{S,[1,\infty]}$ in $M \otimes_{R^+[[z]]} B_{S,[1,1]}$. \square

3.6. We turn to the first main result of this section, which algebraizes G -bundles on \mathcal{Y}_S when S is a product of points as in [15, Definition 1.2].

Recall that Spa yields an anti-equivalence from the category of perfectoid Huber pairs over $\mathbb{F}_q[[\zeta_i]]_{i \in I}$ to the category of affinoid perfectoid spaces over \mathbb{D}^I . Let $S = \text{Spa}(R, R^+)$ be an affinoid perfectoid space over \mathbb{D}^I , and for all i in I , write Γ_i for the graph of its i -th projection $S \rightarrow \mathbb{D}$, which is a closed effective Cartier divisor on \mathcal{Y}_S [11, Proposition VI.1.2 (i)].

Theorem. *Suppose that S is a product of points as in [15, Definition 1.2], and let $1 \leq j \leq k$ be an integer. Then pullback yields an equivalence of groupoids*

$$\{G\text{-bundles on } \text{Spec } R^+[[z]]\} \xrightarrow{\sim} \{G\text{-bundles on } \mathcal{Y}_S\},$$

where morphisms on the left-hand side are given by isomorphisms of their pullbacks to $\text{Spec } R^+[[z]][\frac{1}{z-\zeta_i}]_{j \in I_j}$, and morphisms on the right-hand side are given by isomorphisms of their pullbacks to $\mathcal{Y}_S \setminus \sum_{i \in I_j} \Gamma_i$ that are meromorphic along $\sum_{i \in I_j} \Gamma_i$.

Proof. First, we tackle full faithfulness. Write $\mathcal{O}(\sum_{i \in I_j} \Gamma_i)$ for the line bundle on \mathcal{Y}_S associated with the closed effective Cartier divisor $\sum_{i \in I_j} \Gamma_i$, and let \mathcal{G} and \mathcal{G}' be G -bundles on \mathcal{Y}_S . The Tannakian description of G -bundles implies that an isomorphism $\mathcal{G}|_{\mathcal{Y}_S \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \mathcal{G}'|_{\mathcal{Y}_S \setminus \sum_{i \in I_j} \Gamma_i}$ that is meromorphic along $\sum_{i \in I_j} \Gamma_i$ corresponds to a family of morphisms of vector bundles over \mathcal{Y}_S

$$\mathcal{G}(V) \rightarrow \mathcal{G}'(V) \otimes \mathcal{O}(\sum_{i \in I_j} \Gamma_i)^{\otimes n(V)}$$

that is functorial in V , compatible with tensor products, and compatible with duals, where V runs over objects of $\text{Rep}_{\mathcal{O}_F}(G)$ and $n(V)$ is a large enough integer. Hence full faithfulness follows immediately from Proposition 3.5.

As for essential surjectivity, let \mathcal{G} be a G -bundle on \mathcal{Y}_S . By [31, Theorem 2.7.7], $\mathcal{G}|_{\mathcal{Y}_{S,[0,1]}}$ and $\mathcal{G}|_{\mathcal{Y}_{S,[1,\infty]}}$ correspond to G -bundles N_0 and N_∞ on $\text{Spec } B_{S,[0,1]}$ and $\text{Spec } B_{S,[1,\infty]}$, respectively. Note that the z -adic completion of $R^+[[z]][\frac{z}{\varpi}]$ equals $R^+[\langle \frac{z}{\varpi} \rangle]$ as rings, so the global sections of the rational open subspace $\{|z| \leq |\varpi| \neq 0\}$ of $\text{Spa}(A'(R^+)[\frac{1}{z}], A'(R^+))$ equals $R[\langle \frac{z}{\varpi} \rangle][\frac{1}{z}] = B_{S,[0,1]}[\frac{1}{z}]$ as rings. We have seen in the proof of Proposition 3.4 that the global sections of the rational open subspace

$\{|\varpi| \leq |z| \neq 0\}$ of $\mathrm{Spa}(A'(R^+)[\frac{1}{z}], A'(R^+))$ equals $B_{S,[1,\infty]}$. Because these two rational open subspaces cover $\mathrm{Spa}(A'(R^+)[\frac{1}{z}], A'(R^+))$, Lemma 3.3 and [31, Theorem 2.7.7] enable us to glue $N_0[\frac{1}{z}]^4$ and N_∞ into a G -bundle $N_{\frac{1}{z}}$ on $\mathrm{Spec} A'(R^+)[\frac{1}{z}] = \mathrm{Spec} R^+(\!(z)\!)$.

Note that the z -adic completion of $R^+[[z]][\frac{1}{\varpi}]$ equals $R[[z]]$. Since

$$N_{\frac{1}{z}} \otimes_{R^+(\!(z)\!)} B_{S,[0,1]}[\frac{1}{z}] = N_0[\frac{1}{z}],$$

we see that $N_{\frac{1}{z}}[\frac{1}{\varpi}] \otimes_{R^+(\!(z)\!)[\frac{1}{\varpi}]} R((z)) = N_0 \otimes_{B_{S,[0,1]}} R((z))$. Therefore we can apply Beauville–Laszlo to the vanishing locus of z in $\mathrm{Spec} R^+[[z]][\frac{1}{\varpi}]$ to glue $N_{\frac{1}{z}}[\frac{1}{\varpi}]$ and $N_0 \otimes_{B_{S,[0,1]}} R[[z]]$ into a G -bundle $N_{\frac{1}{\varpi}}$ on $\mathrm{Spec} R^+[[z]][\frac{1}{\varpi}]$. As $N_{\frac{1}{\varpi}}[\frac{1}{z}] = N_{\frac{1}{z}}[\frac{1}{\varpi}]$, we can glue $N_{\frac{1}{\varpi}}$ and $N_{\frac{1}{z}}$ into a G -bundle \mathring{N} on the complement of the vanishing locus of ϖ and z in $\mathrm{Spec} R^+[[z]]$. Finally, because S is a product of points, [1, Proposition 11.5] uniquely extends \mathring{N} to a G -bundle N on $\mathrm{Spec} R^+[[z]]$.

Let us verify that the pullback of N to \mathcal{Y}_S equals \mathcal{G} . Because $N[\frac{1}{z}] = \mathring{N}[\frac{1}{z}] = N_{\frac{1}{z}}$, we see that $N \otimes_{R^+[[z]]} B_{S,[1,\infty]} = N_\infty$. Thus we just need to show $N \otimes_{R^+[[z]]} B_{S,[0,1]} = N_0$. We have $N[\frac{1}{\varpi}] = \mathring{N}[\frac{1}{\varpi}] = N_{\frac{1}{\varpi}}$, so

$$N \otimes_{R^+[[z]]} B_{S,[0,1]}[\frac{1}{z}] = N_{\frac{1}{z}} \otimes_{R^+(\!(z)\!)} B_{S,[0,1]}[\frac{1}{z}] = N_0[\frac{1}{z}].$$

Note that the z -adic completion of $B_{S,[0,1]} = R\langle \frac{z}{\varpi} \rangle$ equals $R[[z]]$, and

$$N \otimes_{R^+[[z]]} R[[z]] = N_{\frac{1}{\varpi}} \otimes_{R^+[[z]][\frac{1}{\varpi}]} R[[z]] = N_0 \otimes_{B_{S,[0,1]}} R[[z]].$$

Hence the desired result follows from applying the uniqueness of Beauville–Laszlo gluing to the vanishing locus of z in $\mathrm{Spec} B_{S,[0,1]}$. \square

3.7. We have the following version of non-abelian Artin–Schreier–Witt theory for \mathcal{O}_F . Recall the terminology of τ -modules as in [42, Definition 12.3.3], and let n be a positive integer. For any \mathcal{O}_F/z^n -local system \mathbb{L} on $\mathrm{Spec} R$, write $M(\mathbb{L})$ for the τ -module over $\mathrm{Spec} R[[z]]/z^n$ given by $\mathbb{L} \otimes_{\mathcal{O}_F/z^n} (\mathcal{O}_{\mathrm{Spec} R[[z]]/z^n}, \mathrm{id})$. Conversely, for any τ -module (M, ϕ) over $\mathrm{Spec} R[[z]]/z^n$, write $\mathbb{L}(M, \phi)$ for the \mathcal{O}_F/z^n -sheaf over $\mathrm{Spec} R$ given by $\underline{\mathrm{Hom}}_{\tau\text{-mod}}((\mathcal{O}_{\mathrm{Spec} R[[z]]/z^n}, \mathrm{id}), (M, \phi))$.

Proposition. *Our $M(-)$ yields an exact tensor equivalence of categories*

$$\{\mathcal{O}_F/z^n\text{-local systems on } \mathrm{Spec} R\} \xrightarrow{\sim} \{\tau\text{-modules over } \mathrm{Spec} R[[z]]/z^n\}.$$

Consequently, $\mathbb{L} \mapsto \mathbb{L} \otimes_{\mathcal{O}_F} (\mathcal{O}_{\mathrm{Spec} R[[z]]}, \mathrm{id})$ is an exact tensor equivalence of categories

$$\{\mathcal{O}_F\text{-local systems on } S\} \xrightarrow{\sim} \{\tau\text{-modules over } \mathrm{Spec} R[[z]]\}.$$

Proof. Note that $M(-)$ is left adjoint to $\mathbb{L}(-)$, and the unit $\mathrm{id} \rightarrow \mathbb{L}(M(-))$ is an isomorphism. So we just need to prove that $M(-)$ is essentially surjective. Because \mathcal{O}_F/z^n -local systems are trivial after a finite étale cover, it suffices to prove that the same holds for τ -modules over $\mathrm{Spec} R[[z]]/z^n$.

So let (M, ϕ) be a τ -module over $\mathrm{Spec} R[[z]]/z^n$ such that M has rank h . When $n = 1$, the desired result is [31, Lemma 3.2.7]. For $n \geq 2$, by induction there exists a finite étale cover $\mathrm{Spec} R' \rightarrow \mathrm{Spec} R$ such that the pullback of (M, ϕ) to $\mathrm{Spec} R'[[z]]/z^{n-1}$ has a basis fixed by $\phi_{R'[[z]]/z^{n-1}}$. Nakayama’s lemma shows that any lift of this basis to $R'[[z]]/z^n$ yields a basis of $M \otimes_R R'$. In these coordinates,

⁴By abuse of notation, we apply notation for pullbacks of vector bundles to G -bundles.

we see that $\phi_{R'[z]/z^n}^{-1}$ acts by $A \circ \tau$, where A in $\mathrm{GL}_h(R'[z]/z^n)$ satisfies $A \equiv 1 \pmod{z^{n-1}}$.

Write $\mathrm{Spec} \tilde{R}$ for the vanishing locus in $\mathrm{Spec} R'[u_{ab}]_{1 \leq a, b \leq h}$ of the matrix

$$\tau(U) - U - \frac{1}{z^{n-1}}(A - 1),$$

where U denotes the matrix with entries u_{ab} . Examining entrywise shows that \tilde{R} is finite over R' , the Jacobian criterion shows that \tilde{R} is étale over R' , and checking on fibers shows that $\mathrm{Spec} \tilde{R} \rightarrow \mathrm{Spec} R'$ is surjective. Finally, on $\tilde{R}[z]/z^n$ we have $(1 + z^{n-1}U)A\tau(1 + z^{n-1}U)^{-1} = (1 + z^{n-1}U)(1 + A - 1)(1 - z^{n-1}U - (A + 1)) = 1$, so the basis of $M \otimes_R \tilde{R}$ given by $1 + z^{n-1}U$ is fixed by $\phi_{\tilde{R}}^{-1}$. Therefore the pullback of (M, ϕ) to $\mathrm{Spec} \tilde{R}[z]/z^n$ is trivial, as desired. \square

3.8. We can upgrade Proposition 3.7 to G -bundles as follows. Briefly, let X be a scheme or a sousperfectoid adic space over \mathcal{O}_F , and let $\tau : X \rightarrow X$ be an endomorphism over \mathcal{O}_F . By a τ - G -bundle over X , we mean a G -bundle \mathcal{G} on X along with an isomorphism of G -bundles $\phi : \mathcal{G} \xrightarrow{\sim} \tau^*\mathcal{G}$.

Let n be a positive integer or ∞ , and define z^∞ to be 0. For any $G(\mathcal{O}_F/z^n)$ -bundle \mathbb{P} on S , by abuse of notation write $M(\mathbb{P})$ for the τ - G -bundle over $\mathrm{Spec} R[[z]]/z^n$ given by $\mathbb{P} \times_{G(\mathcal{O}_F/z^n)} (G, \mathrm{id})$.

Proposition. *Our $M(-)$ yields an equivalence of groupoids*

$$\{G(\mathcal{O}_F/z^n)\text{-bundles on } S\} \xrightarrow{\sim} \{\tau\text{-}G\text{-bundles over } \mathrm{Spec} R[[z]]/z^n\}.$$

Proof. The assignment $\mathbb{P} \mapsto (V \mapsto \mathbb{P} \times_{G(\mathcal{O}_F/z^n)} V(\mathcal{O}_F/z^n))$ yields a functor

$$\{G(\mathcal{O}_F/z^n)\text{-bundles on } S\} \longrightarrow \left\{ \begin{array}{c} \mathcal{O}_F\text{-linear exact tensor functors} \\ \mathrm{Rep}_{\mathcal{O}_F}(G) \rightarrow \{\underline{\mathcal{O}_F/z^n}\text{-local systems on } S\} \end{array} \right\}.$$

By Proposition 3.7 and the Tannakian description of G -bundles, the right-hand side is equivalent to the groupoid of τ - G -bundles over $\mathrm{Spec} R[[z]]/z^n$. Now we just need to prove that the above functor is an equivalence of groupoids. Because $G(\mathcal{O}_F/z^n)$ -bundles are trivial after a pro-étale cover, it suffices to prove that the same holds for objects of the right-hand side.

So let $\rho : \mathrm{Rep}_{\mathcal{O}_F}(G) \rightarrow \{\underline{\mathcal{O}_F/z^n}\text{-local systems on } S\}$ be an \mathcal{O}_F -linear exact tensor functor, and let $\tilde{S} \rightarrow S$ be a pro-étale cover such that \tilde{S} is strictly totally disconnected. Then $\underline{\mathcal{O}_F/z^n}$ -local systems on \tilde{S} are equivalent to finite projective $\mathrm{Cont}(|\tilde{S}|, \underline{\mathcal{O}_F/z^n})$ -modules, so $\rho|_{\tilde{S}}$ corresponds to a G -bundle $\tilde{\mathcal{G}}$ on

$$\mathrm{Spec} \mathrm{Cont}(|\tilde{S}|, \underline{\mathcal{O}_F/z^n}).$$

Note that $\mathrm{Cont}(|\tilde{S}|, \underline{\mathcal{O}_F/z^n}) = \mathrm{Cont}(\pi_0(\tilde{S}), \underline{\mathcal{O}_F/z^n})$. For any s be in $\pi_0(\tilde{S})$, [31, Lemma 2.2.3] indicates that $\varinjlim_U \mathrm{Cont}(U, \underline{\mathcal{O}_F/z^n})$ is Henselian with respect to the kernel of evaluation at s , where U runs over neighborhoods of s in $\pi_0(\tilde{S})$. Lang's lemma shows that the pullback of $\tilde{\mathcal{G}}$ to $\mathrm{Spec} \mathrm{Cont}(s, \underline{\mathcal{O}_F/z^n}) = \mathrm{Spec} \underline{\mathcal{O}_F/z^n}$ is trivial, so Hensel lifting implies that the pullback of $\tilde{\mathcal{G}}$ to $\mathrm{Spec} \mathrm{Cont}(U, \underline{\mathcal{O}_F/z^n})$ is trivial for some U . Therefore $\rho|_{\tilde{U}}$ is isomorphic to the canonical fiber functor, where \tilde{U} denotes the preimage of U in $|\tilde{S}|$. As s varies, this yields a pro-étale cover of S where ρ is trivial, as desired. \square

3.9. Let us recall the equicharacteristic version of the (*relative integral*) *Robba ring*. Write $\|-\|$ for the spectral norm on R , normalized such that $\|\varpi\| = \frac{1}{q}$. For any positive rational b , we have a map $\|-\|_b : R[[z]] \rightarrow [0, \infty]$ given by

$$\sum_{m=0}^{\infty} a_m z^m \mapsto \sup_{m \geq 0} \{q^{-m} \|a_m\|_b^m\}.$$

Evidently $\|\tau(-)\|_b = \|-\|_{qb}$. When $1/b$ lies in $\mathbb{Z}[\frac{1}{p}]$, 3.5 shows that the restriction of $\|-\|_b$ to $B_{S,[0,b]} \subseteq R[[z]]$ is a norm and induces the usual topology on $B_{S,[0,b]}$. Moreover, $\sum_{m=0}^{\infty} a_m z^m$ lies in $B_{S,[0,b]}$ if and only if $\|a_m z^m\|_b \rightarrow 0$.

Write $\widetilde{\mathcal{R}}_R^{\text{int}}$ for $\varinjlim_b B_{S,[0,b]}$, where b runs over positive rationals. Note that any multiple f of z in $\widetilde{\mathcal{R}}_R^{\text{int}}$ satisfies $\|f\|_b < 1$ for small enough b , so the completeness of $B_{S,[0,b]}$ implies that z lies in the Jacobson radical of $\widetilde{\mathcal{R}}_R^{\text{int}}$.

3.10. Just like \mathcal{O}_F -local systems, we show that τ -modules over the Robba ring are trivial after a pro-finite étale cover.

Lemma. *Let $(\widetilde{M}, \widetilde{\phi})$ be a τ -module over $\text{Spec } \widetilde{\mathcal{R}}_R^{\text{int}}$ such that \widetilde{M} is free of rank h . Then there exists a pro-finite étale cover $\text{Spa}(\widetilde{R}, \widetilde{R}^+) \rightarrow S$ such that the pullback of $(\widetilde{M}, \widetilde{\phi})$ to $\text{Spec } \widetilde{\mathcal{R}}_R^{\text{int}}$ is trivial.*

Proof. Proposition 3.7 enables us to assume that the pullback of $(\widetilde{M}, \widetilde{\phi})$ to $\text{Spec } R$ has a basis fixed by ϕ_R . Now 3.9 and Nakayama's lemma show that any lift of this basis yields a basis of \widetilde{M} , and in these coordinates, we see that ϕ^{-1} acts by $A \circ \tau$, where A in $\text{GL}_h(\widetilde{\mathcal{R}}_R^{\text{int}})$ satisfies $A \equiv 1 \pmod{z}$. Proposition 3.7 yields a pro-finite étale cover $\text{Spa}(\widetilde{R}, \widetilde{R}^+) \rightarrow \text{Spa}(R, R^+)$ such that the pullback of $(\widetilde{M}, \widetilde{\phi})$ to $\text{Spec } \widetilde{R}[[z]]$ has a basis fixed by $(\widetilde{\phi})_{\widetilde{R}[[z]]}$. Since the pullback of $(\widetilde{M}, \widetilde{\phi})$ to $\text{Spec } R$ is already trivial, we can choose this basis of $\widetilde{M} \otimes_{\widetilde{\mathcal{R}}_R^{\text{int}}} \widetilde{R}[[z]]$ such that its matrix U in $\text{GL}_h(\widetilde{R}[[z]])$ satisfies $U \equiv 1 \pmod{z}$. Now we just need to prove that U lies in $\text{GL}_h(\widetilde{\mathcal{R}}_R^{\text{int}})$.

As $A-1$ is divisible by z , we have $\|A-1\|_b < 1$ for small enough positive rational b . Write $C := \max\{q^{-1}, \|A-1\|_b\} < 1$, write U_n for the mod- z^n truncation of U , and write X_n for the z^n -coefficient of U . For any positive integer n , we claim that

$$\|z^n X_n\|_{qb}, \|U_n - 1\|_b, \text{ and } \|U_n - 1\|_{qb} \leq C.$$

When $n = 1$, the last two bounds hold because $U_1 = 1$. For general n , we have

$$\begin{aligned} U_n + z^n X_n &\equiv U \equiv A\tau(U) \equiv A(\tau(U_n) + z^n \tau(X_n)) \pmod{z^{n+1}} \\ \implies z^n (X_n - A\tau(X_n)) &\equiv (A-1)\tau(U_n) + (\tau(U_n) - 1) - (U_n - 1) \pmod{z^{n+1}} \\ \implies X_n - \tau(X_n) &\equiv \frac{1}{z^n} [(A-1)\tau(U_n) + (\tau(U_n) - 1) - (U_n - 1)] \pmod{z}. \end{aligned}$$

By evaluating this equation at rank-1 points of S and considering the Newton polygon of its entries, induction on n implies that

$$\begin{aligned} \|X_n\|_b &\leq \max\{1, (q^n \|(A-1)\tau(U_n) + \tau(U_n) - 1) - (U_n - 1)\|_b\}^{1/q} \\ &\leq \max\{1, (q^n C)^{1/q}\} \leq (q^n C)^{1/q}. \end{aligned}$$

Therefore $\|z^n X_n\|_{qb} \leq C$, so $\|U_{n+1} - 1\|_{qb} \leq C$. Since $C \geq q^{-n}$, we also get

$$\|U_{n+1} - 1\|_b \leq \max\{\|z^n X_n\|_b, \|U_n - 1\|_b\} \leq \max\{q^{-n} (q^n C)^{1/q}, C\} \leq C,$$

which concludes our proof of the claim.

By 3.9, the claim implies that U has coefficients in $B_{S,[0,b']}$ for any positive rational $b' < qb$ such that $1/b'$ lies in $\mathbb{Z}[\frac{1}{p}]$. After decreasing b' such that $b' < b$, the claim also implies that U is invertible over $B_{S,[0,b']}$. Therefore U indeed lies in $\mathrm{GL}_h(\tilde{\mathcal{R}}_R^{\mathrm{int}})$, as desired. \square

3.11. Vector bundles on the Robba ring are local on S in the following sense. Let $(S_\alpha)_\alpha$ be a finite cover of S by rational open subspaces, where $S_\alpha = \mathrm{Spa}(R_\alpha, R_\alpha^+)$. Write $S_{\alpha\beta} = \mathrm{Spa}(R_{\alpha\beta}, R_{\alpha\beta}^+)$ for their pairwise intersections, and write $S_{\alpha\beta\gamma} = \mathrm{Spa}(R_{\alpha\beta\gamma}, R_{\alpha\beta\gamma}^+)$ for their triple intersections.

Lemma. *Pullback yields an equivalence from the category of vector bundles on $\mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}$ to the category of vector bundles on the $\mathrm{Spec} \tilde{\mathcal{R}}_{R_\alpha}^{\mathrm{int}}$ with transition morphisms on the $\mathrm{Spec} \tilde{\mathcal{R}}_{R_{\alpha\beta}}^{\mathrm{int}}$ whose pullbacks to $\mathrm{Spec} \tilde{\mathcal{R}}_{R_{\alpha\beta\gamma}}^{\mathrm{int}}$ satisfy the cocycle condition. Moreover, for any vector bundle M on $\mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}$, there exists $(S_\alpha)_\alpha$ as above such that $M|_{\mathrm{Spec} \tilde{\mathcal{R}}_{R_\alpha}^{\mathrm{int}}}$ is trivial for all α .*

Proof. Because $\tilde{\mathcal{R}}_R^{\mathrm{int}} = \varinjlim_b B_{S,[0,b]}$, we have an equivalence of categories

$$\varinjlim_b \{\text{vector bundles on } \mathrm{Spec} B_{S,[0,b]}\} \xrightarrow{\sim} \{\text{vector bundles on } \mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}\}.$$

When $1/b$ lies in $\mathbb{Z}[\frac{1}{p}]$, the $B_{S,[0,b]}$ are Tate algebras over R . Hence $S \mapsto B_{S,[0,b]}$ commutes with rational localization on S . Applying [31, Theorem 2.7.7] to the resulting open cover of $\mathcal{Y}_{S,[0,b]}$ by $(\mathcal{Y}_{S_\alpha,[0,b]})_\alpha$ shows that vector bundles on $\mathrm{Spec} B_{S,[0,b]}$ are equivalent to vector bundles on the $\mathrm{Spec} B_{S_\alpha,[0,b]}$ with transition morphisms on the $\mathrm{Spec} B_{S_{\alpha\beta},[0,b]}$ whose pullbacks to $\mathrm{Spec} B_{S_{\alpha\beta\gamma},[0,b]}$ satisfy the cocycle condition. Because there are finitely many α , taking the directed limit over b yields the first claim.

For the second claim, [31, Theorem 2.7.7] shows that there exists $(S_\alpha)_\alpha$ as above such that the pullback of M to $\mathrm{Spec} R_\alpha$ is trivial for all α . Since z lies in the Jacobson radical of $\tilde{\mathcal{R}}_{R_\alpha}^{\mathrm{int}}$, any trivialization lifts to $\tilde{\mathcal{R}}_{R_\alpha}^{\mathrm{int}}$ by Nakayama's lemma. \square

3.12. We conclude by showing that τ -modules on $R[[z]]$ uniquely descend to the Robba ring.

Theorem. *Pullback yields an exact tensor equivalence of categories*

$$\{\tau\text{-modules over } \mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}\} \xrightarrow{\sim} \{\tau\text{-modules over } \mathrm{Spec} R[[z]]\}.$$

Consequently, pullback induces an equivalence of groupoids

$$\{\tau\text{-}G\text{-bundles over } \mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}\} \xrightarrow{\sim} \{\tau\text{-}G\text{-bundles over } \mathrm{Spec} R[[z]]\}.$$

Proof. First, we tackle full faithfulness. By considering internal homs for τ -modules, it suffices to prove that, for any τ -module $(\tilde{M}, \tilde{\phi})$ over $\mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}$, any m in $\tilde{M} \otimes_{\tilde{\mathcal{R}}_R^{\mathrm{int}}} R[[z]]$ that is fixed by $\tilde{\phi}_{R[[z]]}$ lies in \tilde{M} . Lemma 3.11 implies that it suffices to prove this after passing to an open cover of S , so we can assume that \tilde{M} is free of rank h . Then Lemma 3.10 yields a pro-finite étale cover $\mathrm{Spa}(\tilde{R}, \tilde{R}^+) \rightarrow S$ such that the pullback of $(\tilde{M}, \tilde{\phi})$ to $\mathrm{Spec} \tilde{\mathcal{R}}_{\tilde{R}}^{\mathrm{int}}$ has a basis fixed by $\tilde{\phi}_{\tilde{\mathcal{R}}_{\tilde{R}}^{\mathrm{int}}}$. In these coordinates, the entries of m lie in $(\tilde{R}^\tau)[[z]]$, which lies in $\tilde{\mathcal{R}}_{\tilde{R}}^{\mathrm{int}}$ by 3.9. Note that the intersection of $R[[z]]$ and $\tilde{\mathcal{R}}_{\tilde{R}}^{\mathrm{int}}$ equals $\tilde{\mathcal{R}}_R^{\mathrm{int}}$, so the flatness of \tilde{M} shows that m lies in \tilde{M} .

As for essential surjectivity, let (M, ϕ) be a τ -module over $\text{Spec } R[[z]]$. By passing to a clopen cover of S , we can assume that M has rank h . Proposition 3.7, full faithfulness, and finite étale descent enable us to assume that the pullback of (M, ϕ) to $\text{Spec } R$ has a basis fixed by ϕ_R . Nakayama's lemma shows that any lift of this basis yields a basis of $M \otimes_{\tilde{\mathcal{R}}_R^{\text{int}}} R[[z]]$, and in these coordinates, we see that $\phi_{R[[z]]}^{-1}$ acts by $A \circ \tau$, where A in $\text{GL}_h(R[[z]])$ satisfies $A \equiv 1 \pmod{z}$.

Let n be a positive integer. We inductively construct certain C_n , B_n , and U_n in $\text{GL}_h(R[[z]])$ such that $C_n - B_n$ is divisible by z^n . First, set $C_1 := A$ and $B_1 := 1$. For general n , write X_n for the z^n -coefficient of $C_n - B_n$. There exists Y_n in $\text{Mat}_h(R)$ satisfying $\|X_n + Y_n - \tau(Y_n)\|_1 < q^{n/2}$ [31, Lemma 8.5.2], which we use to define

$$U_n := 1 + z^n Y_n, \quad C_{n+1} := U_n C_n \tau(U_n)^{-1}, \quad \text{and} \quad B_{n+1} := B_n + z^n (X_n + Y_n - \tau(Y_n)).$$

By induction, we have

$$\begin{aligned} C_{n+1} &\equiv (1 + z^n Y_n) C_n (1 - z^n \tau(Y_n)) \\ &\equiv B_n + z^n (X_n + Y_n - \tau(Y_n)) \equiv B_{n+1} \pmod{z^{n+1}}, \end{aligned}$$

as desired.

We see from 3.9 that the B_n converge to a matrix B in $\text{GL}_h(B_{S,[0,1]})$. Now the C_n converge to a matrix C in $\text{GL}_h(R[[z]])$, and because $C_n - B_n$ is divisible by z^n , we have $C = B$. Moreover, the infinite product $U := U_1 U_2 \cdots$ converges to a matrix U in $\text{GL}_h(R[[z]])$, and the above shows that $U A \tau(U)^{-1} = C = B$. Thus the basis of $M \otimes_{\tilde{\mathcal{R}}_R^{\text{int}}} R[[z]]$ given by U descends (M, ϕ) to a τ -module over $\text{Spec } \tilde{\mathcal{R}}_R^{\text{int}}$, as desired.

Finally, we show that pullback has an exact tensor quasi-inverse. Note that we have a commutative triangle

$$\begin{array}{ccc} & \{\mathcal{O}_F\text{-local systems on } S\} & \\ & \swarrow & \searrow^{M(-)} \\ \{\tau\text{-modules over } \text{Spec } \tilde{\mathcal{R}}_R^{\text{int}}\} & \longrightarrow & \{\tau\text{-modules over } \text{Spec } R[[z]]\}. \end{array}$$

Every arrow is an exact tensor functor, and $M(-)$ is an exact tensor equivalence by Proposition 3.7. Hence its quasi-inverse $\mathbb{L}(-)$ postcomposed with the left arrow yields an exact tensor quasi-inverse to pullback. \square

4. ANALYTIC MODULI OF LOCAL SHTUKAS

In this section, we define local shtukas in the analytic setting and compare them with the formal variant from §2. We start by giving an algebraic version of local shtukas over a perfectoid space, which is the equicharacteristic version of Breuil–Kisin–Fargues modules. This mediates between the formal variant and more analytic variants. Next, we define an analytic version of local shtukas, as well as the corresponding moduli problem. Using results from §3, we show that the analytic moduli problem agrees with the formal moduli problem from §2.

From here, we define the covering tower for our analytic moduli problem. We conclude by recalling the moduli of local shtukas appearing in Fargues–Scholze [11], which is defined purely in terms of the Fargues–Fontaine curve. While this subtly differs from our analytic moduli problem, their intersection homology complexes are naturally isomorphic, which is all we need.

4.1. Let $S = \mathrm{Spa}(R, R^+)$ be an affinoid perfectoid space over \mathbb{D}^I . For any i in I , if ζ_i is an $R^{\circ\circ}$ -multiple of ϖ^r , then

$$\frac{1}{z - \zeta_i} = \frac{1}{z} \sum_{n=0}^{\infty} \left(\frac{\zeta_i}{z} \right)^n$$

lies in $R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]$. As ζ_i is topologically nilpotent, this always holds for small enough r .

Recall the μ_i and \mathbb{D}_i from 1.5, and recall Definition 2.1. We use Definition 2.1 to define an algebraic version of local G -shtukas over S .

Definition.

- An *algebraic local G -shtuka* over S is a local G -shtuka over $\mathrm{Spec} R^+$.
- Suppose that S lies over $\prod_{i \in I} \mathbb{D}_i$, and let \mathcal{G} be an algebraic local shtuka over S . We say that \mathcal{G} is *bounded by μ_\bullet* if the corresponding local G -shtuka over $\mathrm{Spec} R^+$ is bounded by μ_\bullet .
- Let \mathcal{G} and \mathcal{G}' be algebraic local G -shtukas over S . A *quasi-isogeny* from \mathcal{G} to \mathcal{G}' consists of, for some small enough positive r in $\mathbb{Z}[\frac{1}{p}]$ and all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\delta_j : \mathcal{G}_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} \xrightarrow{\sim} \mathcal{G}'_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]}$$

such that the diagram

$$\begin{array}{ccc} \mathcal{G}_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} & \xrightarrow{(\phi_j)_{R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]}} & \mathcal{G}_{j+1}|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} \\ \downarrow \delta_j & & \downarrow \delta_{j+1} \\ \mathcal{G}'_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} & \xrightarrow{(\phi'_j)_{R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]}} & \mathcal{G}'_{j+1}|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} \end{array}$$

commutes, where δ_{k+1} denotes the isomorphism ${}^{\tau} \delta_1$.

4.2. Let n be a non-negative integer, and note that R^+/ϖ^n is a discrete $\mathbb{F}_q[[\zeta_i]]_{i \in I}$ -algebra. For any algebraic local shtuka \mathcal{G} over S , write \mathcal{G}^n for the local shtuka over $S_n := \mathrm{Spec} R^+/\varpi^n$ given by pullback. Since $R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]/\varpi^n$ equals $(R^+/\varpi^n)((z))$, quasi-isogenies of algebraic local G -shtukas over S pull back to quasi-isogenies of local G -shtukas over S_n .

Lemma 2.7 shows that bounded algebraic local G -shtukas are all captured by this limit process. The following lemma shows that quasi-isogenies between them are also all captured by this limit process.

Lemma. *Suppose that S lies over $\prod_{i \in I} \mathbb{D}_i$, and let \mathcal{G} and \mathcal{G}' be algebraic local G -shtukas over S bounded by μ_\bullet . Then pullback yields a bijection*

$$\{\text{quasi-isogenies from } \mathcal{G} \text{ to } \mathcal{G}'\} \xrightarrow{\sim} \varprojlim_n \{\text{quasi-isogenies from } \mathcal{G}^n \text{ to } \mathcal{G}'^n\}.$$

Proof. Let $(\delta^n)_{n \geq 0}$ be a compatible system of quasi-isogenies from \mathcal{G}^n to \mathcal{G}'^n . Because $\varprojlim_n (R^+/\varpi^n)((z))$ equals $R^+[[z, \frac{1}{z}]]$, we see that $\delta_j := \varprojlim_n \delta_j^n$ yields an isomorphism of G -bundles $\mathcal{G}_j|_{\mathrm{Spec} R^+[[z, \frac{1}{z}]]} \xrightarrow{\sim} \mathcal{G}'_j|_{\mathrm{Spec} R^+[[z, \frac{1}{z}]]}$ for all $1 \leq j \leq k$. Now δ^0 is bounded by m for some non-negative integer m as in Definition 2.2.b), so Proposition 2.3 yields a non-negative integer B such that δ^n is bounded by $m +$

$B[\log_q n]$. From here, the Tannakian description of G -bundles implies that δ_j naturally descends to an isomorphism of G -bundles

$$\mathcal{G}_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} \xrightarrow{\sim} \mathcal{G}'_j|_{\mathrm{Spec} R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]}$$

for any positive r in $\mathbb{Z}[\frac{1}{p}]$. By taking r small enough such that $\frac{1}{z-\zeta_i}$ lies in $R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]$ for all i in I , the commutativity of the square in Definition 4.1.c) follows from the commutativity of the analogous square in Definition 2.2.a). \square

4.3. Before introducing the analytic version of local G -shtukas, we need some notation on the B_{dR} -affine Grassmannian. Write $B_{\mathrm{dR}}^+(S)$ for the ring of global sections of the completion of $\mathcal{O}_{\mathcal{Y}_S}$ along $\sum_{i \in I} \Gamma_i$, and write $B_{\mathrm{dR}}^j(S)$ for the version that is punctured along $\sum_{i \in I_j} \Gamma_i$.

Definition.

- a) Write $\mathcal{L}_I^n G$ and $\mathcal{L}_I^+ G$ for the small v-sheaves over $(\mathbb{D}^I)^\diamond$ given by sending S to $G(\mathcal{O}_{\sum_{i \in I} \Gamma_i})$ and $G(B_{\mathrm{dR}}^+(S))$, respectively.
- b) Write $\mathcal{G}_G^{(I_1, \dots, I_k)}$ for the small v-sheaf over $(\mathbb{D}^I)^\diamond$ whose S -points parametrize data consisting of
 - i) for all $1 \leq j \leq k$, a G -bundle \mathcal{G}_j on $\mathrm{Spec} B_{\mathrm{dR}}^+(S)$,
 - ii) for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\phi_j : \mathcal{G}_j|_{\mathrm{Spec} B_{\mathrm{dR}}^+(S)} \xrightarrow{\sim} \mathcal{G}_{j+1}|_{\mathrm{Spec} B_{\mathrm{dR}}^+(S)},$$

where \mathcal{G}_{k+1} denotes the trivial G -bundle.

4.4. In certain cases, we can describe the functor of points of (generalized) analytifications without analytically sheafifying. Briefly, let A be a noetherian ring, and let X be a scheme locally of finite type over $Z := \mathrm{Spec} A$. Let $J \subseteq A$ be an ideal, write \hat{A} for the completion of A with respect to J , and write \hat{Z} for the adic space $\mathrm{Spa} \hat{A}$. Write $X_{\hat{Z}}$ for the fiber product as in [26, (3.8)].

Lemma. *Suppose that X is quasi-projective over Z . For any analytic affinoid adic space $S = \mathrm{Spa}(R, R^+)$, the S -points of $X_{\hat{Z}}$ consist of the R -points of X such that the resulting ring homomorphism $A \rightarrow R$ is continuous for the J -adic topology on A .*

Proof. The universal property of $X_{\hat{Z}}$ [26, (3.8)] indicates that an S -point of $X_{\hat{Z}}$ is equivalent to a morphism $S \rightarrow \hat{Z}$ of adic spaces along with a morphism $S \rightarrow X$ of locally ringed spaces such that, in the category of locally ringed spaces, the square

$$\begin{array}{ccc} S & \longrightarrow & X \\ \downarrow & & \downarrow \\ \hat{Z} & \longrightarrow & Z \end{array}$$

commutes. The Spec-global sections adjunction shows that $S \rightarrow X \rightarrow Z$ yields a ring homomorphism $A \rightarrow R$, and note that the commutativity of this square is equivalent to $A \rightarrow R$ being continuous for the J -adic topology on A .

Now assume that $X = \mathbb{P}_Z^N$. Since Z is affine, the Spec-global sections adjunction implies that $S \rightarrow X$ is equivalent to the data of a line bundle \mathcal{L} on S along with sections s_0, \dots, s_N that generate \mathcal{L} . By [30, Theorem 1.4.2], this is equivalent

to a finite projective R -module M of rank 1 along with elements r_0, \dots, r_N that generate M , which is precisely the data of an R -point of X .

In general, X is a locally closed subscheme of \mathbb{P}_Z^N . Because Z is noetherian, there exist finitely many homogeneous polynomials f_1, \dots, f_l and g_1, \dots, g_m in $A[t_0, \dots, t_N]$ such that $X \subseteq \mathbb{P}_Z^N$ is the locus where $f_a(s_0, \dots, s_N)$ vanishes for all $1 \leq a \leq l$ and $g_b(s_0, \dots, s_N)$ does not vanish for all $1 \leq b \leq m$. These properties are preserved by [30, Theorem 1.4.2], so we see that $S \rightarrow X$ is equivalent to an R -point of X . \square

4.5. We check that the B_{dR} -affine Grassmannian and its affine Schubert varieties are the analytifications of their algebraic counterparts. Write S^{alg} for the R -point of C^I given by $\text{Spec } R \rightarrow \text{Spec } \mathbb{F}_q[\zeta_i]_{i \in I} \rightarrow C^I$, and write Γ_i^{alg} for the resulting relative effective Cartier divisor on $C \times S$ as in 1.2. Recall the F_i from 1.5.

Lemma. *We have a natural isomorphism of rings $\mathcal{O}_{n \sum_{i \in I} \Gamma_i^{\text{alg}}} \cong \mathcal{O}_{n \sum_{i \in I} \Gamma_i}$. Consequently, we obtain natural isomorphisms from $(L_I^n(G_C))_{\mathbb{D}^I}^\diamond$ and $(L_I^+(G_C))_{\mathbb{D}^I}^\diamond$ to $\mathcal{L}_I^n(G)$ and $\mathcal{L}_I^+(G)$, respectively, and we may view $(\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i})^\diamond$ as a closed subsheaf*

$$\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond} \subseteq \mathcal{G}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}.$$

Finally, the S -points of $\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \text{Spd } F_i}$ consist of the $((\mathcal{G}_j)_{j=1}^k, (\phi_j)_{j=1}^k)$ such that, for all geometric points \bar{s} of S and $1 \leq j \leq k$, the relative position of $\phi_{j, \bar{s}}$ at $\Gamma_{i, \bar{s}}$ is bounded by $\sum_{i'} \mu_{i'}$, where i' runs over elements of I satisfying $\Gamma_{i', \bar{s}} = \Gamma_{i, \bar{s}}$.

Proof. The first claim is immediate, which identifies $(L_I^n(G_C))_{\mathbb{D}^I}^\diamond$ with $\mathcal{L}_I^n(G)$. The first claim also induces isomorphisms $\widehat{\mathcal{O}}_C(S^{\text{alg}}) \cong B_{\text{dR}}^+(S)$ and $\widehat{\mathcal{O}}_C^{j, \circ}(S^{\text{alg}}) \cong B_{\text{dR}}(S)$, which identifies $(L_I^+(G_C))_{\mathbb{D}^I}^\diamond$ with $\mathcal{L}_I^+(G)$. This also shows that, for any presentation of $\text{Gr}_{G_C}^{(I_1, \dots, I_k)}$ as a directed limit $\varinjlim_l X_l$ of projective schemes X_l over C^I , we have

$$\mathcal{G}_G^{(I_1, \dots, I_k)}(S) = \text{Gr}_{G_C}^{(I_1, \dots, I_k)}(S^{\text{alg}}) = (\varinjlim_l X_l)(S^{\text{alg}}) = \varinjlim_l X_l(S^{\text{alg}}) = \varinjlim_l (X_l)_{\mathbb{D}^I}^\diamond(S),$$

where the last two equalities follow from [23, Lemma 5.4] and Lemma 4.4, respectively. Now 1.5 indicates that $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ is a closed subscheme of $X_l|_{\prod_{i \in I} C_i}$ for large enough l . Since $\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ is projective over $\prod_{i \in I} C_i$, the natural morphism of adic spaces $\widehat{\text{Gr}}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow (\text{Gr}_{G_C, \mu_\bullet}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathbb{D}_i}$ is an isomorphism [26, (4.6.iv.d)]. Hence taking $(-)^{\diamond}$ yields the desired closed subsheaf

$$\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond} \subseteq \mathcal{G}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}.$$

Finally, the description of $\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \text{Spd } F_i}$ follows from 1.5. \square

4.6. Now, we can define an analytic version of local G -shtukas over S . Let a in $\mathbb{Z}[\frac{1}{p}]$ be non-negative. For any i in I , if ζ_i^a is an R° -multiples of ϖ , then $\text{rad}(\Gamma_i)$ lie in $[0, a)$. As ζ_i is topologically nilpotent, this always holds for large enough a .

Definition.

- a) An *analytic local G -shtuka* over S consists of
- i) for all $1 \leq j \leq k$, a G -bundle \mathcal{G}_j on $\mathcal{Y}_{S, [0, \infty)}$,

ii) for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\phi_j : \mathcal{G}_j|_{\mathcal{Y}_{S,[0,\infty)} \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \mathcal{G}_{j+1}|_{\mathcal{Y}_{S,[0,\infty)} \setminus \sum_{i \in I_j} \Gamma_i},$$

that is meromorphic along $\sum_{i \in I_j} \Gamma_i$, where \mathcal{G}_{k+1} denotes the G -bundle ${}^\tau \mathcal{G}_1$.

b) Suppose that S lies over $\prod_{i \in I} \mathbb{D}_i$, and let \mathcal{G} be an analytic local G -shtuka over S . We say that \mathcal{G} is *bounded by μ_\bullet* if, for any affinoid perfectoid étale cover $\mathrm{Spa}(\tilde{R}, \tilde{R}^+) \rightarrow S$ where ${}^\tau \mathcal{G}_1|_{\mathcal{Y}_{\mathrm{Spa}(\tilde{R}, \tilde{R}^+), [0,\infty)}}$ is trivial and any trivialization $t : {}^\tau \mathcal{G}_1|_{\mathcal{Y}_{\mathrm{Spa}(\tilde{R}, \tilde{R}^+), [0,\infty)}} \xrightarrow{\sim} G$, the $\mathrm{Spa}(\tilde{R}, \tilde{R}^+)$ -point of $\mathcal{G}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$ given by

$$\mathcal{G}_1|_{\mathrm{Spec} B_{\mathrm{dR}}^+(\tilde{R})} \overset{(\phi_1)_{B_{\mathrm{dR}}^+(\tilde{R})}}{\dashrightarrow} \cdots \overset{(\phi_{k-1})_{B_{\mathrm{dR}}^{k-1}(\tilde{R})}}{\dashrightarrow} \mathcal{G}_k|_{B_{\mathrm{dR}}^+(\tilde{R})} \overset{(t \circ \phi_k)_{B_{\mathrm{dR}}^+(\tilde{R})}}{\dashrightarrow} G$$

lies in $\mathcal{G}_G^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$.

c) Let \mathcal{G} and \mathcal{G}' be analytic local G -shtukas over S . A *quasi-isogeny* from \mathcal{G} to \mathcal{G}' consists of, for some large enough rational a and all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\delta_j : \mathcal{G}_j|_{\mathcal{Y}_{S,[a,\infty)}} \xrightarrow{\sim} \mathcal{G}'_j|_{\mathcal{Y}_{S,[a,\infty)}}$$

such that the diagram

$$\begin{array}{ccc} \mathcal{G}_j|_{\mathcal{Y}_{S,[a,\infty)}} & \xrightarrow{(\phi_j)_{\mathcal{Y}_{S,[a,\infty)}}} & \mathcal{G}_{j+1}|_{\mathcal{Y}_{S,[a,\infty)}} \\ \downarrow \delta_j & & \downarrow \delta_{j+1} \\ \mathcal{G}'_j|_{\mathcal{Y}_{S,[a,\infty)}} & \xrightarrow{(\phi'_j)_{\mathcal{Y}_{S,[a,\infty)}}} & \mathcal{G}'_{j+1}|_{\mathcal{Y}_{S,[a,\infty)}} \end{array}$$

commutes, where δ_{k+1} denotes the isomorphism ${}^\tau \delta_1$.

It suffices to check Definition 4.6.b) for a single $\mathrm{Spa}(\tilde{R}, \tilde{R}^+) \rightarrow S$ and t .

4.7. We now define the analytic moduli of local G -shtukas.

Definition. Write $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$ for the small v -sheaf over $\prod_{i \in I} \mathbb{D}_i^\diamond$ whose S -points parametrize data consisting of

- i) an analytic local G -shtuka over S bounded by μ_\bullet ,
- ii) a quasi-isogeny δ from \mathcal{G} to the trivial analytic local G -shtuka G .

Write $f^\mathcal{L} : \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond} \rightarrow \prod_{i \in I} \mathbb{D}_i^\diamond$ for the structure morphism.

4.8. Let us compare the formal and analytic moduli of local G -shtukas. Recall $\mathcal{L}\mathrm{oc}\mathfrak{Sht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ from Definition 2.4.

Proposition. *Our $(\mathcal{L}\mathrm{oc}\mathfrak{Sht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i})^\diamond$ is the analytic sheafification of the presheaf over $\prod_{i \in I} \mathbb{D}_i^\diamond$ whose S -points parametrize data consisting of*

- i) an algebraic local G -shtuka \mathcal{G} over S bounded by μ_\bullet ,
- ii) a quasi-isogeny δ from \mathcal{G} to the trivial algebraic local G -shtuka G .

In particular, we have a canonical morphism of v -sheaves over $\prod_{i \in I} \mathbb{D}_i^\diamond$

$$\underline{\mathrm{an}} : (\mathcal{L}\mathrm{oc}\mathfrak{Sht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i})^\diamond \rightarrow \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$$

given by pulling back (\mathcal{G}, δ) to $\mathcal{Y}_{S,[0,\infty)}$.

Proof. Theorem 2.12 shows that $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ is a locally noetherian formal scheme, so as an adic space it is the analytic sheafification of the presheaf

$$\text{Spa}(A, A^+) \mapsto \text{Hom}(\text{Spa}(A^+, A^+), \mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}).$$

Because R^+ is adic with ideal of definition generated by ϖ , we have

$$\begin{aligned} & \text{Hom}(S, \mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}) \\ &= \text{Hom}(\text{Spf } R^+, \mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}) \\ &= \varprojlim_n \text{Hom}(\text{Spec } R^+/\varpi^n, \mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}). \end{aligned}$$

From here, Lemma 2.7 and Lemma 4.2 yield the first claim. The second claim follows from the fact that $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$ is already a sheaf in the analytic topology, so pulling back (\mathcal{G}, δ) induces a morphism $\underline{\text{an}}$ as desired. \square

4.9. Theorem. *Our $\underline{\text{an}}$ is an isomorphism. Consequently, $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \text{Spd } F_i}$ is a locally spatial diamond.*

Proof. First, we prove that $\underline{\text{an}}$ is an isomorphism. Because products of points as in [15, Definition 1.2] form a basis for the v-topology [15, Example 1.1]⁵ and both $(\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i})^\diamond$ and $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$ are v-sheaves, it suffices to check this on S -points when S is a product of points. Products of points are totally disconnected [15, Proposition 1.6], so we do not need to analytically sheafify when evaluating $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ on them.

So assume that S is a product of points, and let (\mathcal{G}, δ) be an S -point of

$$\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}.$$

For large enough rational a and all $1 \leq j \leq k$, we can use $\delta_j|_{\mathcal{Y}_{S,[a,a]}}$ to glue $\mathcal{G}_j|_{\mathcal{Y}_{S,[0,a]}}$ and $G|_{\mathcal{Y}_{S,[a,\infty]}}$ into a G -bundle $\overline{\mathcal{G}}_j$ on \mathcal{Y}_S . The commutativity of the square in Definition 4.6.c) imply that ϕ_j and id glue into an isomorphism of G -bundles

$$\overline{\phi}_j : \overline{\mathcal{G}}_j|_{\mathcal{Y}_S \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \overline{\mathcal{G}}_{j+1}|_{\mathcal{Y}_S \setminus \sum_{i \in I_j} \Gamma_i},$$

where $\overline{\mathcal{G}}_{k+1}$ denotes the G -bundle ${}^\tau \overline{\mathcal{G}}_1$. Then Theorem 3.6 indicates that $\overline{\mathcal{G}}_j$ and $\overline{\phi}_j$ are uniquely pulled back from a G -bundle $\mathcal{G}_j^{\text{alg}}$ on $\text{Spec } R^+[[z]]$ and an isomorphism of G -bundles $\phi_j^{\text{alg}} : \mathcal{G}_j^{\text{alg}}|_{\text{Spec } R^+[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}} \xrightarrow{\sim} \mathcal{G}_{j+1}^{\text{alg}}|_{\text{Spec } R^+[[z]][\frac{1}{z-\zeta_i}]_{i \in I_j}}$, where $\mathcal{G}_{k+1}^{\text{alg}}$ denotes the G -bundle ${}^\tau \mathcal{G}_1^{\text{alg}}$.

Altogether $\mathcal{G}^{\text{alg}} := ((\mathcal{G}_j^{\text{alg}})_{j=1}^k, (\phi_j^{\text{alg}})_{j=1}^k)$ is an algebraic local G -shtuka over S . Since \mathcal{G} is bounded by μ_\bullet , Lemma 4.5 shows that \mathcal{G}^{alg} is too. Finally, take a for which $r := 1/a$ lies in $\mathbb{Z}[\frac{1}{p}]$. Applying Lemma 3.2, Proposition 3.4, and [31, Theorem 2.7.7] to the canonical isomorphism $\mathcal{G}_j|_{\mathcal{Y}_{S,[a,\infty]}} \xrightarrow{\sim} G$ yields an isomorphism of G -bundles $\delta_j^{\text{alg}} : \mathcal{G}_j^{\text{alg}}|_{\text{Spec } R^+[[z, \frac{\varpi^r}{z}]][\frac{1}{z}]} \xrightarrow{\sim} G$, and we see that $\delta^{\text{alg}} := (\delta_j^{\text{alg}})_{j=1}^k$ is a quasi-isogeny from \mathcal{G}^{alg} to G . The uniqueness of Theorem 3.6 and [31, Theorem 2.7.7] imply that (\mathcal{G}, δ) is uniquely the image of $(\mathcal{G}^{\text{alg}}, \delta^{\text{alg}})$ under $\underline{\text{an}}$. Hence $\underline{\text{an}}$ is bijective on S -points, as desired. Finally, the last statement follows from $\mathcal{L}\text{ocSht}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \text{Spa } F_i}$ being an analytic adic space and [41, Lemma 15.6]. \square

⁵However, in [15, Example 1.1] one must replace the $k(x)$ with its completed algebraic closure $C(x)$ and $k(x)^+$ with its integral closure in $C(x)$.

4.10. Next, we turn to level structures. Let n be a non-negative integer.

Definition. Suppose that S lies over $(\mathrm{Spa} F)^I$, and let \mathcal{G} be an analytic local G -shtuka over S . A *level- n structure* on \mathcal{G} consists of, for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\psi_j : \mathcal{G}_j|_{\mathrm{Spec} R[[z]]/z^n} \xrightarrow{\sim} G$$

such that the diagram

$$\begin{array}{ccc} \mathcal{G}_j|_{\mathrm{Spec} R[[z]]/z^n} & \xrightarrow{(\phi_j)_{R[[z]]/z^n}} & \mathcal{G}_{j+1}|_{\mathrm{Spec} R[[z]]/z^n} \\ \downarrow \psi_j & & \downarrow \psi_{j+1} \\ G & \xlongequal{\hspace{2cm}} & G \end{array}$$

commutes, where \mathcal{G}_{k+1} denotes ${}^\tau \mathcal{G}_1$, and ψ_{k+1} denotes ${}^\tau \psi_1$.

Since S lies over $(\mathrm{Spa} F)^I$, the $(\phi_j)_{R[[z]]/z^n}$ are isomorphisms. Therefore ψ_1 uniquely determines ψ_j for $2 \leq j \leq k$.

4.11. We now define the covering tower of the generic fiber of $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$.

Definition. Write $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$ for the small v-sheaf over $\prod_{i \in I} \mathrm{Spd} F_i$ whose S -points parametrize data consisting of

- i) an analytic local G -shtuka \mathcal{G} over S bounded by μ_\bullet ,
- ii) a quasi-isogeny δ from \mathcal{G} to the trivial analytic local G -shtuka,
- iii) a level- n structure $\psi = (\psi_j)_{j=1}^k$ on \mathcal{G} .

Write $f^\mathcal{L} : \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \rightarrow \prod_{i \in I} \mathrm{Spd} F_i$ for the structure morphism.

4.12. For $n' \geq n$, we have morphisms

$$\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \rightarrow \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$$

given by pulling back ψ_j to $\mathrm{Spec} R[[z]]/z^n$ for all $1 \leq j \leq k$. Write $K_{n', n}$ for the kernel of $G(\mathcal{O}_F/z^{n'}) \rightarrow G(\mathcal{O}_F/z^n)$, and note that $K_{n', n}$ acts on $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$ over $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$ via postcomposition with ψ_j for all $1 \leq j \leq k$.

Proposition. *The morphism*

$$\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \rightarrow \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$$

is finite Galois, where the Galois action is given by that of $K_{n', n}$. Consequently, $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$ is a locally spatial diamond.

Proof. First, take $n = 0$, so that

$$\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} = \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}.$$

For any S -point (\mathcal{G}, δ) of $\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$, form the Cartesian square

$$\begin{array}{ccc} S' & \longrightarrow & \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \\ \downarrow & & \downarrow \\ S & \longrightarrow & \mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \end{array}$$

Then S' parametrizes level- n' structures ψ on \mathcal{G} . Because ψ_1 uniquely determines ψ_j for $2 \leq j \leq k$, we see that level- n' structures on \mathcal{G} are equivalent to trivializations of the τ - G -bundle $(\mathcal{G}_1|_{\mathrm{Spec} R[[z]]/z^{n'}}, (\phi_k \circ \cdots \circ \phi_1)_{R[[z]]/z^{n'}})$ over $\mathrm{Spec} R[[z]]/z^{n'}$. Thus Proposition 3.8 and [41, Proposition 9.7] imply that $S' \rightarrow S$ is finite Galois with the desired Galois action.

For general n , the result follows from the commutative triangle

$$\begin{array}{ccc} \mathrm{LocSht}_{G, \mu_\bullet, n'v}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} & \longrightarrow & \mathrm{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} \\ & \searrow & \swarrow \\ & \mathrm{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i} & \end{array}$$

and compatibility of the $K_{n', n}$ -action with changing n' and n . Finally, the last statement follows from Theorem 4.9 and [41, Lemma 11.21]. \square

4.13. The covering tower enjoys the following Hecke correspondences. Write

$$\mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} := \varprojlim_n \mathrm{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_v},$$

and write K_n for the kernel of $G(\mathcal{O}_F) \rightarrow G(\mathcal{O}_F/z^n)$.

Proposition. *We have a canonical $G(F)$ -action on $\mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)}$ over $\prod_{i \in I} \mathrm{Spd} F_i$ that extends the $G(\mathcal{O}_F)$ -action from 4.12. Consequently, for any g in $G(F)$, we have a canonical finite étale correspondence $\mathbf{1}_{K_n g K_n}$ from $\mathrm{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$ to itself.*

Proof. Let (\mathcal{G}, δ) be an S -point of $\mathrm{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$, and let $(\psi^n)_{n \geq 0}$ be a compatible system of level- n structures ψ^n on \mathcal{G} . For all $1 \leq j \leq k$, we see that $\psi_j := \varprojlim_n \psi_j^n$ yields an isomorphism of G -bundles $\mathcal{G}_j|_{\mathrm{Spec} R[[z]]} \xrightarrow{\sim} G$. For any g in $G(F)$, we get an isomorphism of G -bundles $g \circ (\psi_j)_{R((z))} : \mathcal{G}_j|_{\mathrm{Spec} R((z))} \xrightarrow{\sim} G$, which we use with Beauville–Laszlo to glue $G|_{\mathrm{Spec} R[[z]]}$ and $\mathcal{G}_j|_{\mathcal{Y}_{S, (0, \infty)}}$ into a G -bundle $g \cdot \mathcal{G}_j$ on $\mathcal{Y}_{S, [0, \infty)}$.

Since $(g \cdot \mathcal{G}_j)|_{\mathcal{Y}_{S, (0, \infty)} \setminus \sum_{i \in I_j} \Gamma_i}$ is canonically isomorphic to $\mathcal{G}_j|_{\mathcal{Y}_{S, (0, \infty)} \setminus \sum_{i \in I_j} \Gamma_i}$, the commutativity of the square in Definition 4.10 and Beauville–Laszlo let us glue id and $(\phi_j)_{\mathcal{Y}_{S, (0, \infty)} \setminus \sum_{i \in I_j} \Gamma_i}$ into an isomorphism of G -bundles

$$g \cdot \phi_j : (g \cdot \mathcal{G}_j)|_{\mathcal{Y}_{S, [0, \infty)} \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} (g \cdot \mathcal{G}_{j+1})|_{\mathcal{Y}_{S, [0, \infty)} \setminus \sum_{i \in I_j} \Gamma_i},$$

where $g \cdot \mathcal{G}_{k+1}$ denotes ${}^\tau(g \cdot \mathcal{G}_1)$. As \mathcal{G} is bounded by μ_\bullet , the analytic local G -shtuka $g \cdot \mathcal{G} := ((g \cdot \mathcal{G}_j)_{j=1}^k, (g \cdot \phi_j)_{j=1}^k)$ is too. Because $(g \cdot \mathcal{G}_j)|_{\mathcal{Y}_{S, [a, \infty)}}$ is canonically isomorphic to $\mathcal{G}_j|_{\mathcal{Y}_{S, [a, \infty)}}$, our δ induces a quasi-isogeny from $g \cdot \mathcal{G}$ to G . Since $(g \cdot \mathcal{G}_j)|_{\mathrm{Spec} R[[z]]}$ is canonically trivial, we have the trivial level- n structure $\mathrm{id} = (\mathrm{id})_{j=1}^k$ on $g \cdot \mathcal{G}$.

Altogether, we define the image of $(\mathcal{G}, \delta, (\psi^n)_{n \geq 0})$ under g to be $(g \cdot \mathcal{G}, \delta, (\mathrm{id})_{n \geq 0})$. When g lies in $G(\mathcal{O}_F)$, our $g \circ (\psi_j)_{R((z))}$ above extends to an isomorphism of G -bundles $g \circ \psi_j : \mathcal{G}_j|_{\mathrm{Spec} R[[z]]} \xrightarrow{\sim} G$, and tracing through our identifications shows that

this indeed recovers the action from 4.12. Finally, $\mathbf{1}_{K_n g K_n}$ is given by

$$\begin{array}{ccc} \mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} / (K_n \cap g^{-1} K_n g) & \xrightarrow{g} & \mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} / (g K_n g^{-1} \cap K_n) \\ \downarrow & & \downarrow \\ \mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} / K_n & & \mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} / K_n \end{array}$$

and identifying $\mathrm{LocSht}_{G, \mu_\bullet, \infty v}^{(I_1, \dots, I_k)} / K_n$ with $\mathrm{LocSht}_{G, \mu_\bullet, n v}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathrm{Spd} F_i}$. \square

4.14. Recall the following variant of the moduli of local shtukas, which is defined purely in terms of the Fargues–Fontaine curve. Let K be a compact open subgroup of $G(F)$.

Definition. Write $\mathcal{M}_{G, \mu_\bullet, K}^I |_{\prod_{i \in I} \mathrm{Spd} F_i}$ for the small v-sheaf over $\prod_{i \in I} \mathrm{Spd} F_i$ whose S -points parametrize data consisting of

- i) a G -bundle \mathcal{E} on X_S such that, for all geometric points \bar{s} of S , its pullback $\mathcal{E}_{\bar{s}}$ to $X_{\bar{s}}$ is trivial,
- ii) an isomorphism of G -bundles

$$\alpha : \mathcal{E} |_{X_S \setminus \sum_{i \in I} \Gamma_i} \xrightarrow{\sim} G$$

that is meromorphic along $\sum_{i \in I} \Gamma_i$ such that, for all geometric points \bar{s} of S , the relative position of $\alpha_{\bar{s}}$ at $\Gamma_{i, \bar{s}}$ is bounded by $\sum_{i'} \mu_{i'}$, where i' runs over elements of I satisfying $\Gamma_{i', \bar{s}} = \Gamma_{i, \bar{s}}$,

- iii) a \underline{K} -bundle \mathbb{P} on S whose pushforward along $\underline{K} \rightarrow G(F)$ equals the $\underline{G(F)}$ -bundle on S corresponding to \mathcal{E} via [11, Theorem III.2.4].

Write $f^{\mathcal{M}} : \mathcal{M}_{G, \mu_\bullet, K}^I |_{\prod_{i \in I} \mathrm{Spd} F_i} \rightarrow \prod_{i \in I} \mathrm{Spd} F_i$ for the structure morphism.

Recall that $\mathcal{M}_{G, \mu_\bullet, K}^I |_{\prod_{i \in I} \mathrm{Spd} F_i}$ is a locally spatial diamond.

4.15. The analytic moduli of local G -shtukas is related to the Fargues–Fontaine variant as follows.

Proposition. *We have a canonical morphism*

$$c : \mathrm{LocSht}_{G, \mu_\bullet, n v}^{(I)} |_{\prod_{i \in I} \mathrm{Spd} F_i} \rightarrow \mathcal{M}_{G, \mu_\bullet, K_n}^I |_{\prod_{i \in I} \mathrm{Spd} F_i}$$

of locally spatial diamonds over $\prod_{i \in I} \mathrm{Spd} F_i$.

Proof. Let $(\mathcal{G}, \delta, \psi)$ be an S -point of $\mathrm{LocSht}_{G, \mu_\bullet, n v}^{(I)}$. Theorem 3.12 and Proposition 3.8 show that $(\mathcal{G}_1 |_{\mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}}, (\phi_1)_{\tilde{\mathcal{R}}_R^{\mathrm{int}}})$ corresponds to a $\underline{G(\mathcal{O}_F)}$ -bundle on S , and Proposition 3.8 implies that ψ_1 corresponds to a reduction \mathbb{P} of this $\underline{G(\mathcal{O}_F)}$ -bundle to a $\underline{K_n}$ -bundle. Via continuation by Frobenius, $(\mathcal{G}_1 |_{\mathrm{Spec} \tilde{\mathcal{R}}_R^{\mathrm{int}}}, (\phi_1)_{\tilde{\mathcal{R}}_R^{\mathrm{int}}})$ also induces a τ - G -bundle (\mathcal{F}, v) over $\mathcal{Y}_{S, [0, \infty)}$ such that $(\mathcal{F} |_{\mathcal{Y}_{S, (0, \infty)}}, (v)_{\mathcal{Y}_{S, (0, \infty)}})$ corresponds to the pushforward of \mathbb{P} along $\underline{K_n} \rightarrow \underline{G(F)}$. Therefore the pullback of the G -bundle $\mathcal{E} := (\mathcal{F} |_{\mathcal{Y}_{S, (0, \infty)}}) / (v)_{\mathcal{Y}_{S, (0, \infty)}}^{\mathbb{Z}}$ from X_S to $X_{\bar{s}}$ is trivial for all geometric points \bar{s} of S , and the corresponding $\underline{G(F)}$ -bundle on S via [11, Theorem III.2.4] equals the pushforward of \mathbb{P} along $\underline{K_n} \rightarrow \underline{G(F)}$. Finally, continuation by Frobenius and Lemma 4.5 indicate that δ_1 induces an isomorphism of G -bundles $\alpha : \mathcal{E} |_{X_S \setminus \sum_{i \in I} \Gamma_i} \xrightarrow{\sim} G$ with the desired relative position bound, so altogether $(\mathcal{E}, \alpha, \mathbb{P})$ yields an S -point of $\mathcal{M}_{G, \mu_\bullet, K_n}^I$. \square

4.16. We will need the following results of Fargues–Scholze [11] on the intersection homology of the moduli of local shtukas. Recall the notation of 2.13, and let V be an object of $\text{Rep}_E({}^L G)^I$. Note that

$$\coprod_{\mu_\bullet} \mathcal{L}\text{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)}|_{(\text{Spd } \check{F})^I} \text{ and } \coprod_{\mu_\bullet} \mathcal{M}_{G, \mu_\bullet, K}^I|_{(\text{Spd } \check{F})^I}$$

naturally descend to small v-sheaves $\mathcal{L}\text{ocSht}_{G, V, nv}^{(I_1, \dots, I_k)}$ and $\mathcal{M}_{G, V, K}^I$ over $(\text{Spd } F)^I$, respectively, where μ_\bullet runs over highest weights appearing in $V_{\overline{\mathbb{Q}_\ell}}|_{\widehat{G}^I}$. Proposition 4.12 and [41, Proposition 13.4 (iv)] imply that $\mathcal{L}\text{ocSht}_{G, V, nv}^{(I_1, \dots, I_k)}$ is a locally spatial diamond, and we see that $\mathcal{M}_{G, V, K}^I$ is also a locally spatial diamond.

Let Λ be \mathcal{O}_E or E , and now let V be an object of $\text{Rep}_{\mathcal{O}_E}({}^L G)^I$. If $\Lambda = \mathcal{O}_E$, then by abuse of notation write V for V_E . Write $(\text{Spd } \check{F})^I$ for the I -th power of $\text{Spd } \check{F}$ over $\overline{\mathbb{F}_q}$, and write $'\mathcal{F}_{V, K, \Lambda}^I$ for the object of $D_{\blacksquare}(\mathcal{M}_{G, V, K}^I|_{(\text{Spd } \check{F})^I}, \Lambda)$ obtained from [11, Theorem VI.11.1] and V by first applying the double-dual embedding as in [11, p. 264] and then pulling back to $\mathcal{M}_{G, V, K}^I|_{(\text{Spd } \check{F})^I}$. Write $'\mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k)}$ for the pullback of $'\mathcal{F}_{V, K, \Lambda}^I$ under the composition

$$\mathcal{L}\text{ocSht}_{G, V, nv}^{(I_1, \dots, I_k)}|_{(\text{Spd } \check{F})^I} \rightarrow \mathcal{L}\text{ocSht}_{G, V, nv}^{(I)}|_{(\text{Spd } \check{F})^I} \xrightarrow{c} \mathcal{M}_{G, V, K_n}^I|_{(\text{Spd } \check{F})^I}.$$

Write W_F for the absolute Weil group of F .

Theorem. *Our c induces an isomorphism $f_{\natural}^{\mathcal{L}}(' \mathcal{F}_{V, nv, \Lambda}^{(I)}) \xrightarrow{\sim} f_{\natural}^{\mathcal{M}}(' \mathcal{F}_{V, K, \Lambda}^I)$. Consequently, the object $f_{\natural}^{\mathcal{L}}(' \mathcal{F}_{V, nv, \Lambda}^{(I)})$ of $D_{\blacksquare}((\text{Spd } \check{F})^I, \Lambda)$ naturally arises via pullback from $D(W_F^I, \Lambda)$.*

Proof. Using Theorem 3.12 and Proposition 3.8, the argument in the proof of [11, Proposition IX.3.2] yields the first claim. For the second claim, [11, Proposition VII.3.1 (iii)] enables us to identify $f_{\natural}^{\mathcal{M}}(' \mathcal{F}_{V, K, \Lambda}^I)$ with $i_1^* T_V(i_{1!}(c\text{-Ind}_{K_n}^{G(F)} \Lambda))$ as objects of $D(\Lambda)$, where $i_1 : [* / G(F)] \rightarrow \text{Bun}_G$ is the canonical open embedding, and T_V is the geometric Hecke operator associated with V . Therefore [11, Corollary IX.2.3] yields the desired result. \square

4.17. Finally, we define partial Frobenii for the analytic moduli of local G -shtukas and relate them to partial Frobenii on the Fargues–Fontaine variant as follows. Write $\mathcal{F}r^{(I_1, \dots, I_k)} : \mathcal{L}\text{ocSht}_{G, V, nv}^{(I_1, \dots, I_k)} \rightarrow \mathcal{L}\text{ocSht}_{G, V, nv}^{(I_2, \dots, I_k, I_1)}$ for the morphism that sends

$$\begin{array}{ccc} \mathcal{G}_1 - \xrightarrow{\phi_1} \dots - \xrightarrow{\phi_{k-1}} \mathcal{G}_k - \xrightarrow{\phi_k} \tau \mathcal{G}_1 & & \mathcal{G}_2 - \xrightarrow{\phi_2} \dots - \xrightarrow{\phi_k} \tau \mathcal{G}_1 - \xrightarrow{\tau \phi_1} \tau \mathcal{G}_2 \\ \downarrow \delta_1 & & \downarrow \delta_2 \\ \downarrow & & \downarrow \\ \mathcal{G} \text{ ===== } \dots \text{ ===== } \mathcal{G} \text{ ===== } \mathcal{G} & \text{ to } & \mathcal{G} \text{ ===== } \dots \text{ ===== } \mathcal{G} \text{ ===== } \mathcal{G} \\ \downarrow \delta_k & & \downarrow \tau \delta_1 \\ \downarrow & & \downarrow \\ \mathcal{G} \text{ ===== } \mathcal{G} & & \mathcal{G} \text{ ===== } \mathcal{G} \end{array}$$

Note that $\mathcal{M}_{G, V, K}^I$ naturally descends to a v-sheaf over $(\text{Div}_F^1)^I$, where Div_F^1 denotes the small v-sheaf over $\text{Spd } \mathbb{F}_q$ whose S -points parametrize degree-1 relative effective Cartier divisors of X_S . Write $\varphi_{I_1} : \mathcal{M}_{G, V, K}^I \rightarrow \mathcal{M}_{G, V, K}^I$ for the resulting endomorphism given by geometric q -Frobenius on the i -th factor of $(\text{Spd } F)^I$ for i in I_1 and the identity on all other factors.

Lemma. *We have a commutative diagram*

$$\begin{array}{ccccc}
 \mathcal{L}ocSht_{G,V,nv}^{(I_1, \dots, I_k)} & \longrightarrow & \mathcal{L}ocSht_{G,V,nv}^{(I)} & \xrightarrow{c} & \mathcal{M}_{G,V,K_n}^I \\
 \downarrow \mathcal{F}_\Gamma^{(I_1, \dots, I_k)} & & & & \downarrow \varphi_{I_1} \\
 \mathcal{L}ocSht_{G,V,nv}^{(I_2, \dots, I_k, I_1)} & \longrightarrow & \mathcal{L}ocSht_{G,V,nv}^{(I)} & \xrightarrow{c} & \mathcal{M}_{G,V,K_n}^I.
 \end{array}$$

Proof. This follows immediately from the proof of Proposition 4.15. \square

5. UNIFORMIZING THE MODULI SPACES OF GLOBAL SHUKAS

At this point, we shift focus from local to global considerations. Our goal in this section is to define the uniformization morphism, which is essential for our main results. First, we recall some facts about global shtukas and their moduli spaces. We then take formal completions at a fixed place and define the uniformization morphism on the level of formal stacks. By restricting to a Harder–Narasimhan truncation on the global moduli and using results from §2 on the local moduli, we can pass from formal stacks to formal schemes that are locally formally of finite type over \mathbb{D}^I . This enables us to upgrade the formal étaleness of our uniformization morphism to étaleness, as well as to avoid questions about analytifying stacks. Finally, we extend the uniformization theorem to the covering tower on generic fibers.

5.1. We start by switching our notation to a global context. Let C be a geometrically connected smooth proper curve over a finite field \mathbb{F}_q , and write F for $\mathbb{F}_q(C)$. Fix a separable closure \overline{F} of F , and write Γ_F for $\text{Gal}(\overline{F}/F)$. Write \mathbb{A} for the adèle ring of C , and write \mathbb{O} for its subring of integral adèles.

Let G be a parahoric group scheme over C as in [38, Definition 2.18], and write Z for the center of G . By [3, Proposition 2.2(b)], there exists an SL_h -bundle \mathcal{Y} on C and a closed embedding $\iota : G^{\text{ad}} \rightarrow \underline{\text{Aut}}(\mathcal{Y})$ of group schemes over C such that $\underline{\text{Aut}}(\mathcal{Y})/G^{\text{ad}}$ satisfies [3, (2.1)].

Let T be a maximal subtorus of G_F , and write $X_*^+(T)$ for the set of dominant cocharacters of $T_{\overline{F}}$ with respect to a fixed Borel subgroup $B \subseteq G_{\overline{F}}$ containing $T_{\overline{F}}$. Identify $X_*^+(T)$ with the set of conjugacy classes of cocharacters of $G_{\overline{F}}$. Let $\mu_\bullet = (\mu_i)_{i \in I}$ be in $X_*^+(T)$, and identify the field of definition of μ_i with $\mathbb{F}_q(C_i)$ for some finite generically étale cover $C_i \rightarrow C$. Write $\text{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i}$ for the closed affine Schubert variety as in 1.5.

5.2. Let us recall the definition of global G -shtukas. Let S be an affine scheme over C^I , and adopt the notation of 1.2. Write $\tau : S \rightarrow S$ for the absolute q -Frobenius endomorphism, and by abuse of notation, write $\tau : C \times S \rightarrow C \times S$ for the identity times τ .

Definition.

- a) A *global G -shtuka* over S consists of
- i) for all $1 \leq j \leq k$, a G -bundle \mathcal{G}_j on $C \times S$,
 - ii) for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\phi_j : \mathcal{G}_j|_{C \times S \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \mathcal{G}_{j+1}|_{C \times S \setminus \sum_{i \in I_j} \Gamma_i},$$

where \mathcal{G}_{k+1} denotes the G -bundle ${}^\tau \mathcal{G}_1$.

- b) Suppose that S lies over $\prod_{i \in I} C_i$, and let $\mathcal{G} = ((\mathcal{G}_j)_{j=1}^k, (\phi_j)_{j=1}^k)$ be a global G -shtuka over S . We say that \mathcal{G} is *bounded by μ_\bullet* if the S -point of

$$[L_I^+(G) \backslash \mathrm{Gr}_G^{(I_1, \dots, I_k)}]_{|\prod_{i \in I} C_i}$$

given by $((\mathcal{G}_j|_{\mathrm{Spec} \widehat{\mathcal{O}}_C(S)})_{j=1}^k, ((\phi_j)_{\widehat{\mathcal{O}}_C^\circ(S)})_{j=1}^k)$ lies in $[L_I^+(G) \backslash \mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}]_{|\prod_{i \in I} C_i}$.

- c) Let \mathcal{G} be a global G -shtuka over S . We say that \mathcal{G} has *Harder–Narasimhan polygon bounded by s* if the SL_r -bundle $\iota_*(\mathcal{G}_1^{\mathrm{ad}})$ has Harder–Narasimhan polygon bounded by $s2\rho^\vee$, where $2\rho^\vee$ denotes the sum of positive coroots in SL_h .

5.3. Next, we turn to level structures. Let N be a finite closed subscheme of C .

Definition. Suppose that S lies over $(C \setminus N)^I$, and let \mathcal{G} be a global G -shtuka over S . A *level- N structure* on \mathcal{G} consists of, for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\psi_j : \mathcal{G}_j|_{N \times S} \xrightarrow{\sim} G$$

such that the diagram

$$\begin{array}{ccc} \mathcal{G}_j|_{N \times S} & \xrightarrow{(\phi_j)_N} & \mathcal{G}_{j+1}|_{N \times S} \\ \downarrow \psi_j & & \downarrow \psi_{j+1} \\ G & \xlongequal{\quad} & G \end{array}$$

commutes, where \mathcal{G}_{k+1} denotes ${}^\tau \mathcal{G}_1$, and ψ_{k+1} denotes ${}^\tau \psi_1$.

Since S lies over $(C \setminus N)^I$, the $(\phi_j)_N$ are isomorphisms. Therefore ψ_1 uniquely determines ψ_j for $2 \leq j \leq k$.

5.4. We now recall the moduli of global G -shtukas and its associated structures. Write N_i for the preimage of N in C_i .

Definition. Write $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ for the stack over $\prod_{i \in I} C_i \setminus N_i$ whose S -points parametrize data consisting of

- i) a global G -shtuka \mathcal{G} over S bounded by μ_\bullet ,
- ii) a level- N structure $\psi = (\psi_j)_{j=1}^k$ on \mathcal{G} .

Write $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s}|_{\prod_{i \in I} C_i \setminus N_i}$ for the open substack of $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ whose S -points consist of the (\mathcal{G}, ψ) such that \mathcal{G} has Harder–Narasimhan polygon bounded by s .

Write $f^S : \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i} \rightarrow \prod_{i \in I} C_i \setminus N_i$ for the structure morphism.

Our $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ has an action of $Z(F) \backslash Z(\mathbb{A})$ by twisting. Since the image of Z in $\mathrm{Aut}(\mathcal{V})$ is trivial, $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s}|_{\prod_{i \in I} C_i \setminus N_i}$ is preserved by the $Z(F) \backslash Z(\mathbb{A})$ -action. Finally, note that $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ is the increasing union of the $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s}|_{\prod_{i \in I} C_i \setminus N_i}$.

5.5. For finite closed subschemes $N' \supseteq N$ of C , we have morphisms

$$\mathrm{Sht}_{G, \mu_\bullet, N'}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i} \rightarrow \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i}$$

given by pulling back ψ_j to $N \times S$ for all $1 \leq j \leq k$. Write $K_{N', N}$ for the kernel of $G(\mathcal{O}_{N'}) \rightarrow G(\mathcal{O}_N)$, and note that $K_{N', N}$ acts on $\mathrm{Sht}_{G, \mu_\bullet, N'}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i}$ over $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i}$ via postcomposition with ψ_j for all $1 \leq j \leq k$.

Proposition. *The morphism $\mathrm{Sht}_{G, \mu_\bullet, N'}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i} \rightarrow \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i}$ is finite Galois, where the Galois action is given by that of $K_{N', N}$.*

Proof. When $N = \emptyset$, the result follows from the proof of [43, Proposition 2.16 b)]. For general N , the result follows from the commutative triangle

$$\begin{array}{ccc} \mathrm{Sht}_{G, \mu_\bullet, N'}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i} & \longrightarrow & \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i} \\ & \searrow & \swarrow \\ & \mathrm{Sht}_{G, \mu_\bullet, \emptyset}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N'_i} & \end{array}$$

and compatibility of the $K_{N', N}$ -action with changing N' and N . \square

5.6. **Proposition.** *Our $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} C_i \setminus N_i}$ is a Deligne–Mumford stack that is separated and locally of finite type over $\prod_{i \in I} C_i \setminus N_i$. Moreover, for large enough $\deg N$, our $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} C_i \setminus N_i}$ is a scheme that is separated and locally of finite type over $\prod_{i \in I} C_i \setminus N_i$.*

Proof. The second claim follows from the proof of [32, Lemme 12.19]. Using Proposition 5.5, the first claim follows from the argument in [45, §5.1.5]. \square

5.7. Let \tilde{F} be the finite Galois extension of F such that $\mathrm{Gal}(\tilde{F}/F)$ equals the image of the Γ_F -action on $X_*^+(T)$, and identify \tilde{F} with $\mathbb{F}_q(\tilde{C})$ for some finite generically étale cover $\tilde{C} \rightarrow C$. Write \tilde{N} for the preimage of N in \tilde{C} . Write \hat{G} for the dual group of G_F over \mathcal{O}_E , and write ${}^L G$ for $\hat{G} \rtimes \mathrm{Gal}(\tilde{F}/F)$.

Let V be an object of $\mathrm{Rep}_E({}^L G)^I$. Note that $\prod_{\mu_\bullet} \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{(\tilde{C} \setminus \tilde{N})^I}$ and $\prod_{\mu_\bullet} \mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \Big|_{(\tilde{C} \setminus \tilde{N})^I}$ naturally descend to stacks

$$\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)} \quad \text{and} \quad \mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s}$$

over $(C \setminus N)^I$, respectively, where μ_\bullet runs over highest weights appearing in $V_{\overline{\mathbb{Q}}_l} \Big|_{\hat{G}^I}$. Proposition 5.6 and descent imply that $\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)}$ is a Deligne–Mumford stack that is separated and locally of finite type over $(C \setminus N)^I$, and for large enough $\deg N$, our $\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s}$ is a scheme that is separated and locally of finite type over $(C \setminus N)^I$.

5.8. Write K_N for the kernel of $G(\mathbb{O}) \rightarrow G(\mathcal{O}_N)$. For any g in $G(\mathbb{A})$, recall that we have a canonical finite étale correspondence $\mathbf{1}_{K_N g K_N}$ from $\mathrm{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} \mathbb{F}_q(C_i)}$ to itself [32, Construction 2.20]⁶. Note that $\mathbf{1}_{K_N g K_N}$ commutes with the $Z(F) \backslash Z(\mathbb{A})$ -action.

⁶Although [32, Construction 2.20] only addresses the case when G is split, it extends to the general case. Indeed, this is already implicitly used in [32, (12.16)].

5.9. Definition. Write $\mathrm{Fr}^{(I_1, \dots, I_k)} : \mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)} \rightarrow \mathrm{Sht}_{G, V, N}^{(I_2, \dots, I_k, I_1)}$ for the morphism given by

$$(\mathcal{G}_1 - \xrightarrow{\phi_1} \dots - \xrightarrow{\phi_{k-1}} \mathcal{G}_k - \xrightarrow{\phi_k} \tau \mathcal{G}_1) \longmapsto (\mathcal{G}_2 - \xrightarrow{\phi_2} \dots - \xrightarrow{\phi_k} \tau \mathcal{G}_1 - \xrightarrow{\tau \phi_1} \tau \mathcal{G}_2).$$

Note that $\mathrm{Fr}^{(I_1, \dots, I_k)}$ lies above the endomorphism of $(C \setminus N)^I$ given by geometric q -Frobenius on the i -th factor for i in I_1 and the identity on all other factors.

By [32, Lemme 3.1]⁷, there exists a non-negative integer $\kappa(V)$ such that

$$\begin{aligned} (\mathrm{Fr}^{(I_1, \dots, I_k)})^{-1}(\mathrm{Sht}_{G, V, N}^{(I_2, \dots, I_k, I_1), \leq s}) &\subseteq \mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s + \kappa(V)} \quad \text{and} \\ \mathrm{Fr}^{(I_1, \dots, I_k)}(\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s}) &\subseteq \mathrm{Sht}_{G, V, N}^{(I_2, \dots, I_k, I_1), \leq s + \kappa(V)}. \end{aligned}$$

5.10. At this point, we fix a place of F and begin exploring the interplay between the local and global situations. Let v be a closed point of C , write r for the degree of v , and write \mathcal{O}_v for $\widehat{\mathcal{O}}_{C, v}$. Choose a uniformizer z of \mathcal{O}_v , which yields an identification $\mathcal{O}_v = \mathbb{F}_{q^r}[[z]]$. Write F_v for the fraction field of \mathcal{O}_v , and write \mathbb{D} for the formal scheme $\mathrm{Spf} \mathcal{O}_v$.

Fix a separable closure \overline{F}_v of F_v , and fix an embedding $\overline{F} \rightarrow \overline{F}_v$. By abuse of notation, write G for the pullback of G to \mathcal{O}_v . Using T_{F_v} for our maximal subtorus of G_{F_v} and $B_{\overline{F}_v}$ for our Borel subgroup of $G_{\overline{F}_v}$, we can identify F_i from 1.5 with the closure of $\mathbb{F}_q(C_i)$ in \overline{F}_v as well as identify \mathbb{D}_i from 1.5 with the formal completion of C_i at the closed point v_i of C_i above v induced by $\overline{F} \rightarrow \overline{F}_v$.

5.11. The following two lemmas explain how to resolve the clash between our local and global base fields. Write \mathbb{D}^I for the I -th power of \mathbb{D} over \mathbb{F}_{q^r} . Adopt the notation of 1.3, and let $S = \mathrm{Spec} R$ be an affine scheme over \mathbb{D}^I .

Lemma. *We have a natural isomorphism of affine formal schemes*

$$\coprod_d \mathbb{D} \times_{\mathbb{F}_{q^r, d}} S \xrightarrow{\sim} \mathbb{D} \times S,$$

where $\mathbb{D} \times_{\mathbb{F}_{q^r, d}} S$ denotes the product of $S \rightarrow \mathrm{Spec} \mathbb{F}_{q^r} \xrightarrow{\tau^d} \mathrm{Spec} \mathbb{F}_{q^r}$ and \mathbb{D} over \mathbb{F}_{q^r} , and d runs over \mathbb{Z}/r . Under this identification, $\tau : \mathbb{D} \times S \rightarrow \mathbb{D} \times S$ on the right-hand side corresponds to the disjoint union of $\tau : \mathbb{D} \times_{\mathbb{F}_{q^r, d}} S \rightarrow \mathbb{D} \times_{\mathbb{F}_{q^r, d-1}} S$ on the left-hand side.

Proof. Take $\mathbb{D} \times_{\mathbb{F}_{q^r, d}} S \rightarrow \mathbb{D} \times S$ to be the natural morphism. Since \mathbb{F}_{q^r} is finite Galois over \mathbb{F}_q with $\mathrm{Gal}(\mathbb{F}_{q^r}/\mathbb{F}_q) = \tau^{\mathbb{Z}/r}$, the induced morphism above is an isomorphism. The last statement follows immediately. \square

5.12. Lemma. *A local G -shtuka over S is equivalent to data consisting of*

- i)* for all $1 \leq j \leq k$, a G -bundle \mathcal{H}_j on $\mathbb{D} \times S$,
- ii)* for all $1 \leq j \leq k$, an isomorphism of G -bundles

$$\chi_j : \mathcal{H}_j|_{\mathbb{D} \times S \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \mathcal{H}_{j+1}|_{\mathbb{D} \times S \setminus \sum_{i \in I_j} \Gamma_i},$$

where \mathcal{H}_{k+1} denotes the G -bundle $\tau^* \mathcal{H}_k$.

⁷While [32, Lemme 3.1] only treats the case of split G , it extends to the general case. Indeed, this is already implicitly used in [32, (12.15)].

Proof. Let \mathcal{G} be a local G -shtuka over S , and for all $1 \leq j \leq k$, view \mathcal{G}_j as a G -bundle on $\mathbb{D} \times_{\mathbb{F}_{q^r}} S$. Using Lemma 5.11, we can form \mathcal{H}_j by taking $\tau^d \mathcal{G}_1$ on $\mathbb{D} \times_{\mathbb{F}_{q^r}, d} S$ for $1 \leq d \leq r-1$ and \mathcal{G}_j on $\mathbb{D} \times_{\mathbb{F}_{q^r}} S$. Note that $\tau \mathcal{H}_1$ is given by $\tau^d \mathcal{G}_1$ on $\mathbb{D} \times_{\mathbb{F}_{q^r}, d} S$ for all $1 \leq d \leq r$. Therefore we can form χ_j by taking id on $\mathbb{D} \times_{\mathbb{F}_{q^r}, d} S$ for $1 \leq d \leq r-1$ and ϕ_j on $\mathbb{D} \times_{\mathbb{F}_{q^r}} S$.

Conversely, let $\mathcal{H} := ((\mathcal{H}_j)_{j=1}^k, (\chi_j)_{j=1}^k)$ be as above. Write $(-)|_d$ for restrictions to $\mathbb{D} \times_{\mathbb{F}_{q^r}, d} S$. Since Γ_i lies in $\mathbb{D} \times_{\mathbb{F}_{q^r}} S$ for all i in I , our $\chi_j|_d$ is an isomorphism for all $1 \leq j \leq k$ and $1 \leq d \leq r-1$. By repeatedly using Lemma 5.11, this identifies $\mathcal{H}_j|_d$ with $\tau^d \mathcal{H}_1|_r$. Hence this also identifies $\mathcal{H}_{k+1}|_r$ with $\tau^r \mathcal{H}_1|_r$, so altogether we see that $\mathcal{H}|_r$ yields a local G -shtuka over S . \square

5.13. In our study of the uniformization morphism, we start by defining it on the level of formal stacks. Write $\prod_{i \in I} \mathbb{D}_i$ for the product of the \mathbb{D}_i over \mathbb{F}_{q^r} , and write $\prod_{i \in I} v_i$ for the product of the v_i over \mathbb{F}_{q^r} . Assume that N and v are disjoint, and write $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$ for the formal completion of $\text{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ along $\prod_{i \in I} v_i$ in $\prod_{i \in I} C_i \setminus N_i$.

Proposition. *We have a canonical morphism*

$$\widehat{\Theta} : \text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$$

of stacks over $\prod_{i \in I} \mathbb{D}_i$ that is formally étale.

This result generalizes cases of [3, Theorem 5.3].

Proof. First, we define $\widehat{\Theta}$. Let (\mathcal{G}, δ) be an S -point of $\text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$, and let $((\mathcal{H}_j)_{j=1}^k, (\chi_j)_{j=1}^k)$ be the data corresponding to \mathcal{G} as in Lemma 5.12. For all $1 \leq j \leq k$, Lemma 5.11 shows that taking δ_j on $\mathbb{D} \times_{\mathbb{F}_{q^r}} S$ and $\tau^d \delta_1$ on $\mathbb{D} \times_{\mathbb{F}_{q^r}, d} S$ for $1 \leq d \leq r-1$ yields an isomorphism of G -bundles

$$\epsilon_j : \mathcal{H}_j|_{\mathbb{D} \times S \setminus v \times S} \xrightarrow{\sim} G.$$

Beauville–Laszlo lets us use ϵ_j to glue \mathcal{H}_j and $G|_{C \times S \setminus v \times S}$ into a G -bundle \mathcal{G}_j^\ominus on $C \times S$. Because the square in Definition 2.2.b) commutes, Beauville–Laszlo also lets us glue χ_j and id into an isomorphism of G -bundles

$$\phi_j^\ominus : \mathcal{G}_j^\ominus|_{C \times S \setminus \sum_{i \in I_j} \Gamma_i} \xrightarrow{\sim} \mathcal{G}_{j+1}^\ominus|_{C \times S \setminus \sum_{i \in I_j} \Gamma_i},$$

where we use Lemma 1.3 to identify $R[[z]]$ with $\widehat{\mathcal{O}}_C(S)$, and $\mathcal{G}_{k+1}^\ominus$ denotes the G -bundle $\tau \mathcal{G}_1^\ominus$. As \mathcal{G} is bounded by μ_\bullet , the global G -shtuka $\mathcal{G}^\ominus := ((\mathcal{G}_j^\ominus)_{j=1}^k, (\phi_j^\ominus)_{j=1}^k)$ is too. Because N and v are disjoint, $\mathcal{G}_j^\ominus|_{N \times S}$ and $\phi_j^\ominus|_{N \times S}$ are canonically trivial, so we have the trivial level- N structure $\text{id} = (\text{id})_{j=1}^k$ on \mathcal{G}^\ominus . Altogether, we define $\widehat{\Theta}(\mathcal{G}, \delta)$ to be the S -point $(\mathcal{G}^\ominus, \text{id})$ of $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathbb{D}_i}$.

To see that $\widehat{\Theta}$ is formally étale, let J be an ideal of R satisfying $J^n = 0$, and write $\overline{S} \rightarrow S$ for the associated closed embedding. For any commutative square

$$\begin{array}{ccc} \overline{S} & \longrightarrow & \mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i} \\ \downarrow & \nearrow & \downarrow \widehat{\Theta} \\ S & \longrightarrow & \widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}, \end{array}$$

write $(\overline{\mathcal{G}}, \overline{\delta})$ for the \overline{S} -point of $\mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}$, and write (\mathcal{F}, ψ) for the S -point of $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}$. The restriction of \mathcal{F} to $\mathbb{D} \times S$ yields data as in Lemma 5.12, which corresponds to a local G -shtuka \mathcal{G} over S . As \mathcal{F} is bounded by μ_\bullet , our \mathcal{G} is too. Because the pullback of \mathcal{F} to \overline{S} is $\widehat{\Theta}(\overline{\mathcal{G}}, \overline{\delta})$, we see that the pullback of \mathcal{G} to \overline{S} is $\overline{\mathcal{G}}$. Therefore Proposition 2.3 yields a unique quasi-isogeny δ from \mathcal{G} to G whose pullback to \overline{S} is $\overline{\delta}$.

Consider the S -point of $\mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}$ given by (\mathcal{G}, δ) . The top triangle commutes by construction, and the bottom triangle commutes by the uniqueness of Beauville–Laszlo gluing. Finally, the uniqueness of Proposition 2.3 and Beauville–Laszlo gluing also imply that (\mathcal{G}, δ) is the unique such morphism, as desired. \square

5.14. By restricting to a Harder–Narasimhan truncation and letting the (tame) level be large enough, we can pass from formal stacks to formal schemes. Maintain the assumptions of 5.13, Write $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ for the formal completion of

$$\text{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} C_i \setminus N_i}$$

along $\prod_{i \in I} v_i$ in $\prod_{i \in I} C_i \setminus N_i$, and write $\mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ for the preimage of $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ under $\widehat{\Theta}$.

Proposition. *For large enough deg N , the restriction*

$$\widehat{\Theta} : \mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$$

is an étale morphism of formal schemes.

Proof. Proposition 5.13 shows that the restriction

$$\widehat{\Theta} : \mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i} \rightarrow \widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$$

is formally étale. Because $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ is an open substack of

$$\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i},$$

we see that $\mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ is an open subsheaf of $\mathcal{L}\text{ocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}$, so Theorem 2.12 implies that $\mathcal{L}\text{ocSht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k)} \big|_{\prod_{i \in I} \mathbb{D}_i}$ is a formal scheme that is locally of finite type over $\prod_{i \in I} \mathbb{D}_i$. For large enough deg N , Proposition 5.6 implies that $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \big|_{\prod_{i \in I} \mathbb{D}_i}$ is also a formal scheme that is formally of finite type over $\prod_{i \in I} \mathbb{D}_i$. Hence the above restriction is formally of finite type, so it is an étale morphism of formal schemes. \square

5.15. To add level at v , we need to pass to generic fibers as follows. Maintain the assumptions of 5.14, and assume that $\deg N$ is large enough as in Proposition 5.14. Proposition 5.6 shows that $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} C_i \setminus N_i}$ is separated over $\prod_{i \in I} C_i \setminus N_i$, so the natural morphism of adic spaces

$$\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \mathbb{D}_i} \rightarrow (\text{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathbb{D}_i}$$

is an open embedding [26, (4.6.iv.c)]. Write $\prod_{i \in I} \text{Spa } F_i$ for the product of the $\text{Spa } F_i$ over \mathbb{F}_{q^r} . For any non-negative integer n , write $\widehat{\text{Sht}}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spa } F_i}$ for the preimage of $\widehat{\text{Sht}}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spa } F_i}$ in $(\text{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \text{Spa } F_i}$.

Write $\prod_{i \in I} \text{Spd } F_i$ for the product of the $\text{Spd } F_i$ over \mathbb{F}_{q^r} . Write

$$\text{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spd } F_i}$$

for the preimage of $(\text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spa } F_i})^\diamond$ in $\text{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \text{Spd } F_i}$, where we use Theorem 4.9 to identify $(\text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i})^\diamond$ with

$$\text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} |_{\prod_{i \in I} \mathbb{D}_i^\diamond}.$$

5.16. We can now define the uniformization morphism on generic fibers. Maintain the assumptions of 5.15, and let $S = \text{Spa}(R, R^+)$ be an affinoid perfectoid space over $\prod_{i \in I} \text{Spa } F_i$.

Theorem. *We have a canonical morphism*

$$\Theta_n : \text{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spd } F_i} \rightarrow (\widehat{\text{Sht}}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spa } F_i})^\diamond$$

of locally spatial diamonds over $\prod_{i \in I} \text{Spd } F_i$ that is étale.

Proof. First, we define Θ_n . By Theorem 4.9, an S -point of

$$\text{LocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spd } F_i}$$

corresponds to a cover $(S_\alpha)_\alpha$ of S by rational open subspaces $S_\alpha = \text{Spa}(R_\alpha, R_\alpha^+)$ with pairwise intersections $S_{\alpha\beta} = \text{Spa}(R_{\alpha\beta}, R_{\alpha\beta}^+)$, a family $(\mathcal{G}^\alpha, \delta^\alpha)$ of $\text{Spf } R_\alpha^+$ -points of $\text{LocSht}_{G, \mu_\bullet}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \mathbb{D}_i}$ that agree on $\text{Spf } R_{\alpha\beta}^+$, and a level- n structure ψ on the analytic local G -shtuka over S obtained from gluing the $(\mathcal{G}^\alpha)^{\text{an}}$.

Proposition 5.6 indicates that $\text{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} C_i \setminus N_i}$ is locally of finite type over $\prod_{i \in I} C_i \setminus N_i$, so for all α , our $\Theta(\mathcal{G}^\alpha, \delta^\alpha)$ yields an R_α^+ -point of

$$\text{Sht}_{G, \mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} C_i \setminus N_i}.$$

Write $\mathcal{G}^{\alpha, \Theta}$ for the resulting global G -shtuka over $\text{Spec } R_\alpha$, which is bounded by μ_\bullet and has Harder–Narasimhan polygon bounded by m . Note that the pullback ψ^α of ψ to S_α is precisely a level- nv structure on $\mathcal{G}^{\alpha, \Theta}$, so we can form a level- $(nv + N)$ structure $\psi^{\alpha, \Theta}$ on $\mathcal{G}^{\alpha, \Theta}$ by taking ψ^α on nv and id on N . Then $(\mathcal{G}^{\alpha, \Theta}, \psi^{\alpha, \Theta})$ induces an S_α -point of $\widehat{\text{Sht}}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s} |_{\prod_{i \in I} \text{Spa } F_i}$, and because the $\mathcal{G}^{\alpha, \Theta}$ and $\psi^{\alpha, \Theta}$ agree on $\text{Spec } R_{\alpha\beta}$, the resulting family glues into an S -point. We define this S -point to be the value of Θ_n .

To see that Θ_n is étale, note that we have a commutative square

$$\begin{array}{ccc} \mathcal{L}\text{ocSht}_{G,\mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \text{Spd } F_i} & \longrightarrow & (\mathcal{L}\text{ocSht}_{G,\mu}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \text{Spa } F_i})^\diamond \\ \downarrow \Theta_n & & \downarrow \widehat{\Theta}^\diamond \\ (\widehat{\text{Sht}}_{G,\mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \text{Spa } F_i})^\diamond & \longrightarrow & (\widehat{\text{Sht}}_{G,\mu_\bullet, N}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \text{Spa } F_i})^\diamond. \end{array}$$

Theorem 4.9 and Proposition 4.12 imply that the top arrow is étale, and Proposition 5.5 and [41, Lemma 15.6] imply that the bottom arrow is étale. Proposition 5.14 and [41, Lemma 15.6] show that $\widehat{\Theta}^\diamond$ is étale, so the 2-out-of-3 property [41, Proposition 11.30] concludes that Θ_n is étale. \square

5.17. As before, we reindex everything in terms of representations of the dual group. Maintain the assumptions of 5.15. Let \widetilde{F}_v be the extension of F_v as in 2.13, and identify \widetilde{F}_v with the completion of \widetilde{F} at the place \widetilde{v} of \widetilde{F} above v induced by $\overline{F} \rightarrow \overline{F}_v$. Identify \widehat{G} with the dual group of G_{F_v} over \mathcal{O}_E , and write ${}^L G_v$ for $\widehat{G} \rtimes \text{Gal}(\widetilde{F}_v/F_v)$. Note that we have a natural inclusion ${}^L G_v \rightarrow {}^L G$.

Let V be an object of $\text{Rep}_E({}^L G_v)^I$. Write $\widehat{\text{Sht}}_{G,V,N}^{(I_1, \dots, I_k)}$ and $\widehat{\text{Sht}}_{G,V,N}^{(I_1, \dots, I_k), \leq s}$ for the formal completions of $\text{Sht}_{G,V,N}^{(I_1, \dots, I_k)}$ and $\text{Sht}_{G,V,N}^{(I_1, \dots, I_k), \leq s}$, respectively, along v^I in $(C \setminus N)^I$. Proposition 5.13 and descent yield a canonical morphism

$$\widehat{\Theta} : \mathcal{L}\text{ocSht}_{G,V}^{(I_1, \dots, I_k)} \rightarrow \widehat{\text{Sht}}_{G,V,N}^{(I_1, \dots, I_k)}$$

that is formally étale. Write $\mathcal{L}\text{ocSht}_{G,V}^{(I_1, \dots, I_k), \leq s}$ for the preimage of $\widehat{\text{Sht}}_{G,V,N}^{(I_1, \dots, I_k), \leq s}$ under $\widehat{\Theta}$.

Write $\widehat{\text{Sht}}_{G,V, nv+N}^{(I_1, \dots, I_k), \leq s}$ for the preimage of $\widehat{\text{Sht}}_{G,V,N}^{(I_1, \dots, I_k), \leq s}$ in $(\text{Sht}_{G,V, nv+N}^{(I_1, \dots, I_k)})_{(\text{Spa } F_v)^I}$, and write $\mathcal{L}\text{ocSht}_{G,V, nv}^{(I_1, \dots, I_k), \leq s}$ for the preimage of $(\mathcal{L}\text{ocSht}_{G,V}^{(I_1, \dots, I_k), \leq s})_{(\text{Spa } F_v)^I}^\diamond$ in $\mathcal{L}\text{ocSht}_{G,V, nv}^{(I_1, \dots, I_k)}$, where we use Theorem 4.9 to identify $(\mathcal{L}\text{ocSht}_{G,V}^{(I_1, \dots, I_k), \leq s})_{(\text{Spa } F_v)^I}^\diamond$ with $\mathcal{L}\text{ocSht}_{G,V, 0v}^{(I_1, \dots, I_k)}$. Theorem 5.16 and Galois descent yield a canonical morphism

$$\Theta_n : \mathcal{L}\text{ocSht}_{G,V, nv}^{(I_1, \dots, I_k), \leq s} \rightarrow (\widehat{\text{Sht}}_{G,V, nv+N}^{(I_1, \dots, I_k), \leq s})^\diamond$$

of locally spatial diamonds over $(\text{Spd } F_v)^I$ that is étale.

5.18. We conclude by showing that the uniformization morphism is compatible with partial Frobenii. Maintain the assumptions of 5.15.

Lemma. *Our $\mathcal{F}_\Gamma^{(I_1, \dots, I_k)}$ restricts to a morphism*

$$\mathcal{F}_\Gamma^{(I_1, \dots, I_k)} : \mathcal{L}\text{ocSht}_{G,V, nv}^{(I_1, \dots, I_k), \leq s} \rightarrow \mathcal{L}\text{ocSht}_{G,V, nv}^{(I_1, \dots, I_k), \leq s+r\kappa(V)}.$$

After enlarging $\text{deg } N$, we can also form the r -fold composition

$$(\text{Fr}^{(I_1, \dots, I_k)})_{\tau^{r-1}(\text{Spa } F_v)^{I_1} \times (\text{Spa } F_v)^{I \setminus I_1}} \circ \dots \circ (\text{Fr}^{(I_1, \dots, I_k)})_{(\text{Spa } F_v)^I},$$

which yields a morphism

$$(\text{Fr}^{(I_1, \dots, I_k)})_{(\text{Spa } F_v)^I}^r : (\text{Sht}_{G,V, nv+N}^{(I_1, \dots, I_k), \leq s})_{(\text{Spa } F_v)^I} \rightarrow (\text{Sht}_{G,V, nv+N}^{(I_1, \dots, I_k), \leq s+r\kappa(V)})_{(\text{Spa } F_v)^I}.$$

Finally, we have $\Theta_n \circ \mathcal{F}_\Gamma^{(I_1, \dots, I_k)} = (\text{Fr}^{(I_1, \dots, I_k)})_{(\text{Spa } F_v)^I}^{r, \diamond} \circ \Theta_n$.

Proof. Write $\widehat{\text{Sht}}_{G,V,N}^{(I_1,\dots,I_k)}|_{\tau\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}}$ for the formal completion of $\text{Sht}_{G,V,N}^{(I_1,\dots,I_k)}$ along $\tau(v)^{I_1}\times v^{I\setminus I_1}$ in $(C\setminus N)^I$. We see from 5.9 that $\text{Fr}^{(I_1,\dots,I_k)}$ induces a morphism

$$\widehat{\text{Fr}}^{(I_1,\dots,I_k)} : \widehat{\text{Sht}}_{G,V,N}^{(I_1,\dots,I_k)} \rightarrow \widehat{\text{Sht}}_{G,V,N}^{(I_2,\dots,I_k,I_1)}|_{\tau\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}}.$$

If $r = 1$, then stop here. Otherwise, the relative effective Cartier divisors on $C \times S$ corresponding to S -points of $\tau\mathbb{D}$ and \mathbb{D} are disjoint, so the right-hand side is naturally isomorphic to $\widehat{\text{Sht}}_{G,V,N}^{(I_1,\dots,I_k)}|_{\tau\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}}$. By forming $\widehat{\text{Fr}}^{(I_1,\dots,I_k)}|_{\tau\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}}$ and repeating this $r - 1$ more times, we obtain a morphism

$$\widehat{\text{Fr}}^{(I_1,\dots,I_k)}|_{\tau^{r-1}\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}} \circ \dots \circ \widehat{\text{Fr}}^{(I_1,\dots,I_k)} : \widehat{\text{Sht}}_{G,V,N}^{(I_1,\dots,I_k)} \rightarrow \widehat{\text{Sht}}_{G,V,N}^{(I_2,\dots,I_k,I_1)}.$$

Tracing through our identifications shows that

$$\widehat{\Theta} \circ \widehat{\mathfrak{F}r}^{(I_1,\dots,I_k)} = \widehat{\text{Fr}}^{(I_1,\dots,I_k)}|_{\tau^{r-1}\mathbb{D}^{I_1}\times\mathbb{D}^{I\setminus I_1}} \circ \dots \circ \widehat{\text{Fr}}^{(I_1,\dots,I_k)} \circ \widehat{\Theta},$$

so 5.9 implies that $\widehat{\mathfrak{F}r}^{(I_1,\dots,I_k)}$ restricts to a morphism

$$\mathfrak{LocSht}_{G,V}^{(I_1,\dots,I_k),\leq s} \rightarrow \mathfrak{LocSht}_{G,V}^{(I_1,\dots,I_k),\leq s+r\kappa(V)}.$$

Pulling back to $\text{Spa } F_v$ and using Theorem 4.9 yields the desired result. \square

6. LOCAL-GLOBAL COMPATIBILITY

Our goal in this section is to prove Theorem A. First, we recall the coefficient sheaves used in the cohomology of the global and local moduli problems. We show that they are compatible under the uniformization morphism from §5. Next, we recall smoothness theorems for our cohomology sheaves, which are due to Xue [44] in the global case and Fargues–Scholze [11] in the local case.

These smoothness theorems yield global and local excursion operators. Using the uniformization morphism, we prove that the global and local excursion operators are compatible. From this, we deduce that the Bernstein center elements constructed by Genestier–Lafforgue [14] agree with those constructed by Fargues–Scholze [11], as well as deduce Theorem A.

6.1. For the cohomology of the moduli of global G -shtukas, we use the following coefficient sheaves obtained from geometric Satake. For large enough e , recall from 1.5 that the natural $L_I^+(G)$ -action on $\text{Gr}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i}$ factors through $L_I^e(G)$. Write $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}$ for the $L_I^e(G)$ -bundle on $\text{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i\setminus N_i}$ whose fiber over (\mathcal{G}, ψ) parametrizes trivializations of the G -bundle ${}^\tau\mathcal{G}_1|_{e\sum_{i\in I}\Gamma_i}$. Note that we have a natural $L_I^e(G)$ -equivariant morphism $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)} \rightarrow \text{Gr}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i}$, which is smooth by [32, p. 867]. Write $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k),\leq s}$ for the restriction of $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}$ to $\text{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k),\leq s}|_{\prod_{i\in I} C_i\setminus N_i}$.

Write V_{μ_\bullet} for the highest weight representation of \widehat{G}^I corresponding to μ_\bullet , and write $\mathcal{S}_{\mu_\bullet,E}^{(I_1,\dots,I_k)}$ for the corresponding object of $D(\text{Gr}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i}, E)$ under geometric Satake. Write $\mathcal{F}_{\mu_\bullet,N,E}^{(I_1,\dots,I_k)}$ for the object of $D(\text{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i\setminus N_i}, E)$ obtained by first pulling back $\mathcal{S}_{\mu_\bullet,E}^{(I_1,\dots,I_k)}$ to $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}$ and then using $L_I^e(G)$ -equivariance to descend along $A_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)} \rightarrow \text{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}|_{\prod_{i\in I} C_i\setminus N_i}$. Finally, write $\mathcal{F}_{\mu_\bullet,N,E}^{(I_1,\dots,I_k),\leq s}$ for the restriction of $\mathcal{F}_{\mu_\bullet,N,E}^{(I_1,\dots,I_k)}$ to $\text{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k),\leq s}|_{\prod_{i\in I} C_i\setminus N_i}$.

6.2. We will also take cohomology after quotienting by a lattice Ξ of $Z(F)\backslash Z(\mathbb{A})$, where by a *lattice* we mean a discrete torsionfree cocompact subgroup. We proceed as follows. Note that $L_I^+(Z)$ acts trivially on $\mathrm{Gr}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} C_i}$, so the natural $L_I^+(G)$ -action factors through $L_I^+(G^{\mathrm{ad}})$. For large enough e , 1.5 indicates that this factors through $L_I^e(G^{\mathrm{ad}})$. Now $L_I^e(Z)$ acts trivially on the objects of

$$D(\mathrm{Gr}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} C_i}, E)$$

obtained from geometric Satake [32, Théorème 12.16], so these objects are $L_I^e(G^{\mathrm{ad}})$ -equivariant. Adapting the construction in 6.1 yields an object $\mathcal{F}_{\Xi,\mu_\bullet,N,E}^{(I_1,\dots,I_k)}$ of

$$D(\mathrm{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}/\Xi|_{\prod_{i \in I} C_i \setminus N_i}, E),$$

and we see that the pullback of $\mathcal{F}_{\Xi,\mu_\bullet,N,E}^{(I_1,\dots,I_k)}$ to $\mathrm{Sht}_{G,\mu_\bullet,N}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} C_i \setminus N_i}$ equals $\mathcal{F}_{\Xi,\mu_\bullet,N,E}^{(I_1,\dots,I_k)}$.

6.3. Next, we describe the coefficient sheaves used for the homology of the moduli of local G -shtukas. Recall $\mathcal{L}_I^e(G)$ and $\mathcal{L}_I^+(G)$ from Definition 4.3. For large enough e , 1.5 and Lemma 4.5 indicate that the natural $\mathcal{L}_I^+(G)$ -action on $\mathcal{G}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathbb{D}_i^\diamond}$ factors through $\mathcal{L}_I^e(G)$. Write $\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}$ for the $\mathcal{L}_I^e(G)$ -bundle on

$$\mathrm{LocSht}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}$$

whose fiber over $(\mathcal{G}, \delta, \psi)$ parametrizes trivializations of the G -bundle ${}^{\tau^r} \mathcal{G}_1|_e \sum_{i \in I} \Gamma_i$. Note that we have a natural $\mathcal{L}_I^e(G)$ -equivariant morphism

$$\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)} \rightarrow \mathcal{G}_{G,\mu_\bullet}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathrm{Spd} F_i}.$$

Recall Λ from 4.16, and write $\prod_{i \in I} \mathrm{Spd} \check{F}_i$ for the product of the $\mathrm{Spd} \check{F}_i$ over $\overline{\mathbb{F}}_q$. Write $'\mathcal{F}_{\mu_\bullet,nv,\Lambda}^{(I_1,\dots,I_k)}$ for the object of $D_\blacksquare(\mathrm{LocSht}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i}, \Lambda)$ obtained from [11, Theorem VI.11.1] and V_{μ_\bullet} by first applying the double-dual embedding as in [11, p. 264], then pulling back to $\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i}$, and finally using $\mathcal{L}_I^e(G)$ -equivariance and [41, Proposition 17.3] to descend along

$$\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)} \rightarrow \mathrm{LocSht}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i}.$$

Write $\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k), \leq s}$ for the restriction of $\mathcal{A}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k)}$ to $\mathrm{LocSht}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k), \leq s}|_{\prod_{i \in I} \mathrm{Spd} F_i}$, and write $'\mathcal{F}_{\mu_\bullet,nv,\Lambda}^{(I_1,\dots,I_k), \leq s}$ for the restriction of $'\mathcal{F}_{\mu_\bullet,nv,\Lambda}^{(I_1,\dots,I_k)}$ to

$$\mathrm{LocSht}_{G,\mu_\bullet,nv}^{(I_1,\dots,I_k), \leq s}|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i}.$$

6.4. Our local and global coefficient sheaves are compatible under Θ_n in the following sense. Adapt the assumptions of 5.15, and write $\prod_{i \in I} \mathrm{Spa} \check{F}_i$ for the product of the $\mathrm{Spa} \check{F}_i$ over $\overline{\mathbb{F}}_q$. Write $(\mathcal{F}_{\mu_\bullet,nv+N,E}^{(I_1,\dots,I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}$ for the object of

$$D((\mathrm{Sht}_{G,\mu_\bullet,nv+N}^{(I_1,\dots,I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}, E)$$

obtained by analytifying $\mathcal{F}_{\mu_\bullet,nv+N,E}^{(I_1,\dots,I_k), \leq s}$ as in [27, (3.2.8)]. Because

$$(\mathrm{Sht}_{G,\mu_\bullet,nv+N}^{(I_1,\dots,I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}$$

is an analytic adic space, [41, Lemma 15.6] and [41, Remark 14.14] indicate that $(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}$ yields an object $(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ of

$$D_{\text{ét}}((\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond, E).$$

Lemma. $(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ is universally locally acyclic over $\prod_{i \in I} \mathrm{Spa} \check{F}_i$.

Moreover, its image $'(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ in $D_\bullet((\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond, E)$ under the double-dual embedding as in [11, p. 260] satisfies

$$\Theta_n^* ['(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond] = ' \mathcal{F}_{\mu_\bullet, nv, E}^{(I_1, \dots, I_k), \leq s}.$$

Proof. We start by rewriting $(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ as follows. Since

$$(\mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}$$

is an analytic adic space, [41, Lemma 15.6] and [41, Remark 14.14] indicate that $(\mathcal{S}_{\mu_\bullet, E}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ yields an object of $D_{\text{ét}}((\mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond, E)$. By first pulling back $(\mathcal{S}_{\mu_\bullet, E}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ to $(A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ and then using $\mathcal{L}_I^e(G)$ -equivariance and [41, Proposition 17.3] to descend along

$$(A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond \rightarrow (\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond,$$

where we use Lemma 4.5 to identify $(L_I^e(G))_{\mathbb{D}_I}^\diamond$ with $\mathcal{L}_I^e(G)$, we see that the resulting object of $D_{\text{ét}}((\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond, E)$ equals $(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$.

Let us prove the first claim. By using the explicit description in [11, Proposition VI.7.9] and the fiberwise criterion for perversity [11, Corollary VI.7.6], we see that $(\mathcal{S}_{\mu_\bullet, E}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ equals the object obtained from [11, Theorem VI.11.1] and V_{μ_\bullet} , where we use Lemma 4.5 to identify $(\mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ with $\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)}|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i}$. Hence $(\mathcal{S}_{\mu_\bullet, E}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ is universally locally acyclic over $\prod_{i \in I} \mathrm{Spa} \check{F}_i$. Now 6.1 and [41, Proposition 24.4] show that

$$(A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond \rightarrow (\mathrm{Gr}_{G, \mu_\bullet}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$$

is ℓ -cohomologically smooth, so [11, Proposition IV.2.13 (i)] implies that the pull-back of $(\mathcal{S}_{\mu_\bullet, E}^{(I_1, \dots, I_k)})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ to $(A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$ remains universally locally acyclic over $\prod_{i \in I} \mathrm{Spa} \check{F}_i$. Applying [41, Proposition 24.4] again shows that

$$(A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond \rightarrow (\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$$

is ℓ -cohomologically smooth, so [11, Proposition IV.2.13 (ii)] implies that

$$(\mathcal{F}_{\mu_\bullet, nv+N, E}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}^\diamond$$

is universally locally acyclic over $\prod_{i \in I} \mathrm{Spa} \check{F}_i$, as desired.

For the second claim, note that Θ_n naturally induces a morphism

$$\mathcal{A}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} \rightarrow (A_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})_{\prod_{i \in I} \mathrm{Spa} F_i}^\diamond$$

such that the diagram

$$\begin{array}{ccc}
\mathcal{G}_{G, \mu_\bullet}^{(I_1, \dots, I_k)} \Big|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i} & \xlongequal{\quad} & (\mathrm{G}\Gamma_{G, \mu_\bullet}^{(I_1, \dots, I_k)})^\diamond \Big|_{\prod_{i \in I} \mathrm{Spa} \check{F}_i} \\
\uparrow & & \uparrow \\
\mathcal{A}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i} & \longrightarrow & (\mathcal{A}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})^\diamond \Big|_{\prod_{i \in I} \mathrm{Spa} \check{F}_i} \\
\downarrow & & \downarrow \\
\mathcal{L}\mathrm{ocSht}_{G, \mu_\bullet, nv}^{(I_1, \dots, I_k), \leq s} \Big|_{\prod_{i \in I} \mathrm{Spd} \check{F}_i} & \xrightarrow{\Theta_n} & (\mathrm{Sht}_{G, \mu_\bullet, nv+N}^{(I_1, \dots, I_k), \leq s})^\diamond \Big|_{\prod_{i \in I} \mathrm{Spa} \check{F}_i}
\end{array}$$

commutes. Therefore the above discussion yields the desired result. \square

6.5. We now consider the cohomology of the moduli of global G -shtukas. Let V be an object of $\mathrm{Rep}_E({}^L G)^I$. Note that the $\mathcal{F}_{\mu_\bullet, N, E}^{(I_1, \dots, I_k)}$ and $\mathcal{F}_{\mu_\bullet, N, E}^{(I_1, \dots, I_k), \leq s}$ naturally descend to objects $\mathcal{F}_{V, N, E}^{(I_1, \dots, I_k)}$ and $\mathcal{F}_{V, N, E}^{(I_1, \dots, I_k), \leq s}$ of $D(\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)}, E)$ and $D(\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s}, E)$, respectively, where μ_\bullet runs over highest weights appearing in $V_{\overline{\mathbb{Q}}_\ell} \Big|_{\widehat{G}^I}$ with multiplicity.

Recall that $f_!^S \mathcal{F}_{V, N, E}^{(I_1, \dots, I_k), \leq s}$ is independent of the ordered partition I_1, \dots, I_k [32, p. 868], so we write it as $\mathcal{H}_{V, N, E}^{I, \leq s}$. The same holds for $f_!^S \mathcal{F}_{V, N, E}^{(I_1, \dots, I_k)}$, so we write it as $\mathcal{H}_{V, N, E}^I$. Because $\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)}$ is the increasing union of the $\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k), \leq s}$, we have $\mathcal{H}_{V, N, E}^I = \varinjlim_s \mathcal{H}_{V, N, E}^{I, \leq s}$. Note that 5.8 yields an action of $C_c(K_N \backslash G(\mathbb{A})/K_N, E)$ on $\mathcal{H}_{V, N, E}^I$.

6.6. Recall the following smoothness result of Xue [44]. Write $\overline{\eta}$ for $\mathrm{Spec} \overline{F}$, and write Δ for diagonal morphisms. Write W_F for the absolute Weil group of F , and write $\mathrm{val}_F : W_F \rightarrow \mathbb{Z}$ for the homomorphism that sends geometric q -Frobenii to 1. Write $U \subseteq C$ for the largest open subspace where G_U is reductive.

Theorem. *The cohomology sheaves of $\mathcal{H}_{V, N, E}^I \Big|_{(U \setminus N)^I}$ are ind-smooth, and the cohomology sheaves of $\mathcal{H}_{V, N, E}^I \Big|_{\Delta(\overline{\eta})}$ have a natural action of W_F^I . For any $\gamma_\bullet = (\gamma_i)_{i \in I}$ in W_F^I , the γ_\bullet -action sends the image of the cohomology groups of $\mathcal{H}_{V, N, E}^{I, \leq s} \Big|_{\Delta(\overline{\eta})}$ to the image of the cohomology groups of $\mathcal{H}_{V, N, E}^{I, \leq s'} \Big|_{\Delta(\overline{\eta})}$ for $s' \geq s + \sum_{i \in I} \max\{0, \mathrm{val}_F(\gamma_i)\}$.*

Proof. The first claim follows from the proof of [44, Theorem 6.0.12], and the W_F^I -action follows from the proof of [44, Proposition 6.0.10]. The last claim follows from 5.9. \square

6.7. Let us record the analogous results after quotienting by Ξ . Let V be an object of $\mathrm{Rep}_E({}^L G)^I$, and note that the $\mathcal{F}_{\Xi, \mu_\bullet, N, E}^{(I_1, \dots, I_k)}$ naturally descend to an object $\mathcal{F}_{\Xi, V, N, E}^{(I_1, \dots, I_k)}$ of $D(\mathrm{Sht}_{G, V, N}^{(I_1, \dots, I_k)}/\Xi, E)$, where μ_\bullet runs over highest weights appearing in $V_{\overline{\mathbb{Q}}_\ell} \Big|_{\widehat{G}^I}$ with multiplicity.

Recall that $f_!^S \mathcal{F}_{\Xi, V, N, E}^{(I_1, \dots, I_k)}$ is independent of the ordered partition I_1, \dots, I_k [32, p. 868], so we write it as $\mathcal{H}_{\Xi, V, N, E}^I$. Note that 5.8 yields an action of

$$C_c(K_N \backslash G(\mathbb{A})/K_N, E)$$

on $\mathcal{H}_{\Xi, V, N, E}^I$. Recall that the cohomology sheaves of $\mathcal{H}_{\Xi, V, N, E}^I|_{(U \setminus N)^I}$ are ind-smooth [44, Theorem 6.0.12], and the cohomology sheaves of $\mathcal{H}_{\Xi, V, N, E}^I|_{\Delta(\bar{\eta})}$ have a natural action of W_F^I [44, Proposition 6.0.10].

6.8. Next, we consider the homology of the moduli of local G -shtukas. Let V be an object of $\text{Rep}_E({}^L G_v)^I$. Note that the $'\mathcal{F}_{\mu_\bullet, nv, \Lambda}^{(I_1, \dots, I_k)}$ and $'\mathcal{F}_{\mu_\bullet, nv, \Lambda}^{(I_1, \dots, I_k), \leq s}$ naturally descend to objects $'\mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k)}$ and $'\mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k), \leq s}$ of $D_\blacksquare(\text{LocSht}_{G, V, nv}^{(I_1, \dots, I_k)}|_{(\text{Spd } \check{F}_v)^I}, \Lambda)$ and $D_\blacksquare(\text{LocSht}_{G, V, nv}^{(I_1, \dots, I_k), \leq s}|_{(\text{Spd } \check{F}_v)^I}, \Lambda)$, respectively, where μ_\bullet runs over highest weights appearing in $V_{\bar{\mathbb{Q}}_v}^I|_{\hat{G}^I}$ with multiplicity.

Recall the notation of 4.16. Since the square

$$\begin{array}{ccc} \text{LocSht}_{G, V, nv}^{(I)}|_{(\text{Spd } \check{F}_v)^I} & \xrightarrow{c} & \mathcal{M}_{G, V, K_n}^I|_{(\text{Spd } \check{F}_v)^I} \\ \uparrow & & \downarrow \\ \text{LocSht}_{G, V, nv}^{(I_1, \dots, I_k)}|_{(\text{Spd } \check{F}_v)^I} & \longrightarrow & [\mathcal{L}_I^e(G) \backslash \mathcal{G}_{G, V}^{(I)}|_{(\text{Spd } \check{F}_v)^I}] \end{array}$$

commutes, where $\mathcal{G}_{G, V}^{(I)}$ denotes the natural descent of $\prod_{\mu_\bullet} \mathcal{G}_{G, \mu_\bullet}^{(I)}|_{(\mathbb{D}^I)^\diamond}$ to $(\mathbb{D}^I)^\diamond$, the $'\mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k)}$ defined in 4.16 agrees with the $'\mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k)}$ defined here.

The smallness of convolution implies that $f_{\mathfrak{h}}^{\mathcal{M}}(' \mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k), \leq s})$ is independent of the ordered partition I_1, \dots, I_k , so we write it as $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I, \leq s}$. The same holds for $f_{\mathfrak{h}}^{\mathcal{M}}(' \mathcal{F}_{V, nv, \Lambda}^{(I_1, \dots, I_k)})$, so we write it as $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I}$. Because $\text{LocSht}_{G, V, nv}^{(I_1, \dots, I_k)}$ is the increasing union of the $\text{LocSht}_{G, V, nv}^{(I_1, \dots, I_k), \leq s}$, we have $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I} = \varinjlim_s \mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I, \leq s}$. Note that Proposition 4.13 yields an action of $C_c(K_n \backslash G(F_v)/K_n, E)$ on $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I}$.

Write \mathbb{C}_v for the completion of \bar{F}_v , and write $\bar{\eta}_v$ for $\text{Spd } \mathbb{C}_v$. Theorem 4.16 yields a natural action of $W_{F_v}^I$ on the cohomology groups of $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I}|_{\Delta(\bar{\eta}_v)}$. For any γ_\bullet in W_{F_v} , Lemma 5.18 and Lemma 4.17 imply that the γ_\bullet -action sends the image of the cohomology groups of $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I, \leq s}$ to the image of the cohomology groups of $\mathcal{H}_{V, nv, \Lambda}^{\text{loc}, I, \leq s'}$ for $s' \geq s + \sum_{i \in I} \max\{0, \text{val}_F(\gamma_i)\}$.

6.9. Let us recall some facts about excursion algebras. For any abstract group W , finite group Q with a pinned action on \hat{G} , and group homomorphism $W \rightarrow Q$, write $\text{Exc}(W, \hat{G})$ for the excursion algebra over \mathcal{O}_E as in [11, Definition VIII.3.4]. Recall that $\text{Exc}(W, \hat{G})$ is flat over \mathcal{O}_E and has canonical generators $S_{I, V, x, \xi, \gamma_\bullet}$ subject to explicit relations, where I runs over finite sets, V runs over objects of $\text{Rep}_{\mathcal{O}_E}((\hat{G} \rtimes Q)^I)$, x runs over morphisms $\mathbf{1} \rightarrow V|_{\Delta(\hat{G})}$, ξ runs over morphisms $V|_{\Delta(\hat{G})} \rightarrow \mathbf{1}$, and γ_\bullet runs through W^I .

Proposition. *Let L be an algebraically closed field over \mathcal{O}_E . We have a unique bijection*

$$\left\{ \begin{array}{l} \mathcal{O}_E\text{-algebra homomorphisms} \\ \chi : \text{Exc}(W, \hat{G}) \rightarrow L \end{array} \right\} \xrightarrow{\sim} \left\{ \begin{array}{l} \text{semisimple homomorphisms} \\ \rho : W \rightarrow \hat{G}(L) \rtimes Q \text{ over } Q \end{array} \right\} / \hat{G}(L)\text{-conj.}$$

such that $\chi(S_{I, V, x, \xi, \gamma_\bullet})$ equals the composition

$$L \xrightarrow{x} V(L) \xrightarrow{(\rho(\gamma_i))_{i \in I}} V(L) \xrightarrow{\xi} L.$$

Proof. This follows immediately from [11, Corollary VII.4.3]. \square

6.10. The following theorem summarizes the work of V. Lafforgue [32] and Xue [44] on global excursion operators. Write $\text{Bun}_{G,N}(\mathbb{F}_q)$ for the groupoid of G -bundles on C equipped with a trivialization along N .

Theorem. *There exists a unique E -algebra homomorphism*

$$\text{Exc}(W_F, \widehat{G})_E \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E))$$

that sends $S_{I,V,x,\xi,\gamma_\bullet}$ to the composition

$$\begin{array}{ccccc} C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E) & \xlongequal{\quad} & \mathcal{H}_{\mathbf{1},N,E}^{*,0}|_{\overline{\eta}} & \xrightarrow{x} & \mathcal{H}_{V|\Delta(\widehat{G}),N,E}^{*,0}|_{\overline{\eta}} & \xlongequal{\quad} & \mathcal{H}_{V,N,E}^{I,0}|_{\Delta(\overline{\eta})} \\ & & & & & & \downarrow \gamma_\bullet \\ C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E) & \xlongequal{\quad} & \mathcal{H}_{\mathbf{1},N,E}^{*,0}|_{\overline{\eta}} & \xleftarrow{\xi} & \mathcal{H}_{V|\Delta(\widehat{G}),N,E}^{*,0}|_{\overline{\eta}} & \xlongequal{\quad} & \mathcal{H}_{V,N,E}^{I,0}|_{\Delta(\overline{\eta})}. \end{array}$$

Moreover, the image of $\text{Exc}(W_F, \widehat{G})_E$ in $\text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E))$ preserves the kernel of the surjective $C_c(K_N \backslash G(\mathbb{A})/K_N, E)$ -equivariant map

$$C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E) \rightarrow C_c(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi, E),$$

so we obtain an E -algebra homomorphism

$$\text{Exc}(W_F, \widehat{G})_E \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi, E)).$$

Proof. Arguing as in [32, p. 870] shows that the images of the $S_{I,V,x,\xi,\gamma_\bullet}$ satisfy the explicit relations, so we get the desired E -algebra homomorphism

$$\text{Exc}(W_F, \widehat{G})_E \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E)).$$

Next, because $\text{Sht}_{G,V,N}^{(I_1, \dots, I_k)} \rightarrow \text{Sht}_{G,V,N}^{(I_1, \dots, I_k)}/\Xi$ is étale, 6.2 yields a natural $!$ -pushforward morphism $\mathcal{H}_{V,N,E}^I \rightarrow \mathcal{H}_{\Xi,V,N,E}^I$, which induces a morphism from the composition diagram above to the analogous composition diagram for $\mathcal{H}_{\Xi,V,N,E}^I$. Note that, when $I = *$ and $V = \mathbf{1}$, the natural $!$ -pushforward morphism recovers

$$C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E) \rightarrow C_c(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi, E)$$

on fibers. Thus the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E))$ satisfies the desired property. \square

6.11. We now elaborate on variants of Theorem 6.10. Recall that

$$\text{Bun}_{G,N}(\mathbb{F}_q) \cong \coprod_{\alpha} G_{\alpha}(F) \backslash G_{\alpha}(\mathbb{A})/K_N$$

as groupoids [32, Remarque 12.2], where α runs over G -bundles on $\text{Spec } F$ whose pullback to $\text{Spec } F_c$ is trivial for all closed points c of C , and G_{α} denotes the inner twist of G_F over F associated with α . Hence $C_c(G(F) \backslash G(\mathbb{A})/K_N, E)$ and $C_c(G(F) \Xi \backslash G(\mathbb{A})/K_N, E)$ are $C_c(K_N \backslash G(\mathbb{A})/K_N, E)$ -stable direct summands of

$$C_c(\text{Bun}_{G,N}(\mathbb{F}_q), E) \text{ and } C_c(\text{Bun}_{G,N}(\mathbb{F}_q)/\Xi, E),$$

respectively, so Theorem 6.10 induces E -algebra homomorphisms

$$\text{Exc}(W_F, \widehat{G})_E \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(G(F) \backslash G(\mathbb{A})/K_N, E)),$$

$$\text{Exc}(W_F, \widehat{G})_E \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(G(F) \Xi \backslash G(\mathbb{A})/K_N, E)).$$

6.12. For us, the most convenient interpretation of Fargues–Scholze [11] is the following theorem. Write $\mathfrak{z}_{K_n}(G(F_v), \Lambda)$ for the center of $C_c(K_n \backslash G(F_v)/K_n, \Lambda)$.

Theorem. *There exists a unique Λ -algebra homomorphism*

$$\mathrm{Exc}(W_{F_v}, \widehat{G})_\Lambda \rightarrow \mathfrak{z}_{K_n}(G(F_v), \Lambda)$$

that sends $S_{I,V,x,\xi,\gamma_\bullet}$ to the composition

$$\begin{array}{ccccc} C_c(G(F_v)/K_n, \Lambda) & \xlongequal{\quad} & \mathcal{H}_{\mathbf{1},nv,\Lambda}^{\mathrm{loc},*,0}|_{\overline{\eta}_v} & \xrightarrow{x} & \mathcal{H}_{V|_{\Delta(\widehat{G})},nv,\Lambda}^{\mathrm{loc},*,0}|_{\overline{\eta}_v} & \xlongequal{\quad} & \mathcal{H}_{V,nv,\Lambda}^{\mathrm{loc},I,0}|_{\Delta(\overline{\eta}_v)} \\ & & & & & & \downarrow \gamma_\bullet \\ C_c(G(F_v)/K_n, \Lambda) & \xlongequal{\quad} & \mathcal{H}_{\mathbf{1},nv,\Lambda}^{\mathrm{loc},*,0}|_{\overline{\eta}_v} & \xleftarrow{\xi} & \mathcal{H}_{V|_{\Delta(\widehat{G})},nv,\Lambda}^{\mathrm{loc},*,0}|_{\overline{\eta}_v} & \xlongequal{\quad} & \mathcal{H}_{V,nv,\Lambda}^{\mathrm{loc},I,0}|_{\Delta(\overline{\eta}_v)}. \end{array}$$

Proof. This follows from [11, Corollary IX.2.4] and [11, Theorem VIII.4.1]. \square

6.13. We now prove local-global compatibility on the level of algebras over E . Write \mathbb{A}^v for the away-from- v adeles, write K_N^v for $\mathbb{A}^v \cap K_N$, and let n be the multiplicity of v in N . So $K_N = K_n K_N^v$.

Theorem. *The square*

$$\begin{array}{ccc} \mathrm{Exc}(W_{F_v}, \widehat{G})_E & \xrightarrow{\quad} & \mathfrak{z}_{K_n}(G(F_v), E) \\ \downarrow & & \downarrow \\ \mathrm{Exc}(W_F, \widehat{G})_E & \xrightarrow{\quad} & \mathrm{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, E)}(C_c(G(F) \backslash G(\mathbb{A})/K_N, E)) \end{array}$$

commutes.

Proof. It suffices to check commutativity on the canonical generators $S_{I,V,x,\xi,\gamma_\bullet}$ of $\mathrm{Exc}(W_{F_v}, \widehat{G})_E$, where I is a finite set, V is an object of $\mathrm{Rep}_E((\widehat{G} \rtimes \mathrm{Gal}(\widetilde{F}/F))^I)$, x is a morphism $\mathbf{1} \rightarrow V|_{\Delta(\widehat{G})}$, ξ is a morphism $V|_{\Delta(\widehat{G})} \rightarrow \mathbf{1}$, and γ_\bullet is in $W_{F_v}^I$. This amounts to computing certain actions on $C_c(G(F) \backslash G(\mathbb{A})/K_N, E)$, which we check on the basis given by $\mathbf{1}_{G(F)gK_N}$ for g in $G(\mathbb{A})$. Since the $C_c(K_n \backslash G(F_v)/K_n, E)$ -action commutes with the $C_c(K_N^v \backslash G(\mathbb{A}^v)/K_N^v, E)$ -action, we can assume that the away-from- v components of g equal 1.

Then $\mathbf{1}_{G(F)gK_N}$ equals the image of $\mathbf{1}_{g_v K_n}$ under the natural pushforward map

$$C_c(G(F_v)/K_n, E) \rightarrow C_c(G(F) \backslash G(\mathbb{A})/K_N, E).$$

Because this map commutes with the $C_c(K_n \backslash G(F_v)/K_n, E)$ -action, it also commutes with the action of the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\mathfrak{z}_{K_n}(G(F_v), E)$. Hence we can compute the latter for $\mathbf{1}_{G(F)gK_N}$ by computing it for $\mathbf{1}_{g_v K_n}$.

Fix s such that $\mathbf{1}_{g_v K_n}$ lies in the image of $\mathcal{H}_{\mathbf{1},nv,E}^{*,\leq s,0}$ in $\mathcal{H}_{\mathbf{1},nv,E}^{*,0} = C_c(G(F_v)/K_n, E)$. By Theorem 6.12 and 6.8, the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\mathfrak{z}_{K_n}(G(F_v), E)$ acts on $\mathbf{1}_{g_v K_n}$ via the composition

$$\begin{aligned} & \mathcal{H}_{\mathbf{1},nv,E}^{\mathrm{loc},*,\leq s,0}|_{\overline{\eta}_v} \xrightarrow{x} \mathcal{H}_{V|_{\Delta(\widehat{G})},nv,E}^{\mathrm{loc},*,\leq s,0}|_{\overline{\eta}_v} = \mathcal{H}_{V,nv,E}^{\mathrm{loc},I,\leq s,0}|_{\Delta(\overline{\eta}_v)} \\ (\star) \quad & \xrightarrow{\gamma_\bullet} \mathcal{H}_{V,nv,E}^{\mathrm{loc},I,\leq s',0}|_{\Delta(\overline{\eta}_v)} = \mathcal{H}_{V|_{\Delta(\widehat{G})},nv,E}^{\mathrm{loc},*,\leq s',0}|_{\overline{\eta}_v} \xrightarrow{\xi} \mathcal{H}_{\mathbf{1},nv,E}^{\mathrm{loc},*,\leq s',0}|_{\overline{\eta}_v} \end{aligned}$$

for large enough s' . By enlarging the away-from- v part of N and using the action of $C_c(K_N^v \backslash G(\mathbb{A}^v)/K_N^v, E)$ as before, we can assume that $\deg N$ is large enough. Then Lemma 6.4 shows that Θ_n yields a natural \natural -pushforward morphism

$$\mathcal{H}_{V, nv, \bar{E}}^{\text{loc}, I, \leq s} |_{\Delta(\bar{\eta}_v)} \rightarrow \mathcal{H}_{V, N, E}^{I, \leq s} |_{\Delta(\bar{\eta})},$$

where we use Lemma 6.4, [11, Proposition VII.5.2], and [27, (5.7.2)] to identify

$$(f^S)_{\Delta(\bar{\eta}_v)}^{\diamond} |_{\natural} [({}'\mathcal{F}_{V, N, E}^{(I_1, \dots, I_k), \leq s})_{\Delta(\bar{\eta}_v)}^{\diamond}] = \mathcal{H}_{V, N, E}^{I, \leq s} |_{\Delta(\bar{\eta})}.$$

Lemma 5.18 and Lemma 4.17 imply that $\mathcal{H}_{V, nv, \bar{E}}^{\text{loc}, I, \leq s} |_{\Delta(\bar{\eta}_v)} \rightarrow \mathcal{H}_{V, N, E}^{I, \leq s} |_{\Delta(\bar{\eta})}$ induces a morphism from the composition diagram in Equation (\star) to the composition diagram

$$\begin{aligned} \mathcal{H}_{\mathbf{1}, N, E}^{*, \leq s, 0} |_{\bar{\eta}} &\xrightarrow{x} \mathcal{H}_{V |_{\Delta(\hat{G})}, N, E}^{*, \leq s, 0} |_{\bar{\eta}} = \mathcal{H}_{V, N, E}^{I, \leq s, 0} |_{\Delta(\bar{\eta})} \\ (\star\star) \quad &\xrightarrow{\gamma_{\bullet}} \mathcal{H}_{V, N, E}^{I, \leq s', 0} |_{\Delta(\bar{\eta})} = \mathcal{H}_{V |_{\Delta(\hat{G})}, N, E}^{*, \leq s', 0} |_{\bar{\eta}} \xrightarrow{\xi} \mathcal{H}_{\mathbf{1}, N, E}^{*, \leq s', 0} |_{\bar{\eta}}. \end{aligned}$$

When $I = *$ and $V = \mathbf{1}$, the natural \natural -pushforward morphism recovers

$$C_c(G(F_v)/K_n, E) \rightarrow C_c(G(F) \backslash G(\mathbb{A})/K_N, E)$$

on fibers, so we see that the image of $S_{I, V, x, \xi, \gamma_{\bullet}}$ in $\mathfrak{z}_{K_n}(G(F_v), E)$ acts on $\mathbf{1}_{G(F)gK_N}$ via Equation $(\star\star)$. But Theorem 6.10 and 6.5 indicate that this is precisely how the image of $S_{I, V, x, \gamma_{\bullet}}$ in $\text{Exc}(W_F, \hat{G})_E$ acts on $\mathbf{1}_{G(F)gK_N}$, as desired. \square

6.14. Let us recall the elements of the Bernstein center constructed by Genestier–Lafforgue [14]. Write \mathfrak{m}_E for the maximal ideal of \mathcal{O}_E , and let c be a non-negative integer. Write $\mathfrak{z}_{K_n}(G(F_v), \mathcal{O}_E/\lambda^c)$ for the center of $C_c(K_n \backslash G(F_v)/K_n, \mathcal{O}_E/\mathfrak{m}_E^c)$. For any finite set I , algebraic function f on $\hat{G} \backslash ({}^L G)^I / \hat{G}$, element γ_{\bullet} of $W_{F_v}^I$, and positive integer n , write $\mathfrak{z}_{n, c, I, f, \gamma_{\bullet}}^{\text{GL}}$ for the element of $\mathfrak{z}_{K_n}(G(F_v), \mathcal{O}_E/\mathfrak{m}_E^c)$ constructed in [14, Théorème 1.1]⁸.

6.15. We prove that the elements of the Bernstein center constructed by Fargues–Scholze coincide with those constructed by Genestier–Lafforgue. Recall that the image of $\text{Exc}(W_F, \hat{G})$ in $\text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)}(C_{\text{cusp}}(G(F) \Xi \backslash G(\mathbb{A})/K_N, E))$ preserves $C_{\text{cusp}}(G(F) \Xi \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)$ [32, Proposition 13.1], so 6.11 induces an \mathcal{O}_E -algebra homomorphism

$$\text{Exc}(W_F, \hat{G}) \rightarrow \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)}(C_{\text{cusp}}(G(F) \Xi \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)).$$

For any object V of $\text{Rep}_{\mathcal{O}_E}({}^L G)^I$, morphism $x : \mathbf{1} \rightarrow V |_{\Delta(\hat{G})}$, and morphism $\xi : V |_{\Delta(\hat{G})} \rightarrow \mathbf{1}$, write f for the algebraic function on $\hat{G} \backslash ({}^L G)^I / \hat{G}$ given by $g_{\bullet} \mapsto \xi(g_{\bullet} \cdot x)$.

Theorem. *The square*

$$\begin{array}{ccc} \text{Exc}(W_{F_v}, \hat{G}) & \longrightarrow & \mathfrak{z}_{K_n}(G(F_v), \mathcal{O}_E) \\ \downarrow & & \downarrow \\ \text{Exc}(W_F, \hat{G}) & \longrightarrow & \text{End}_{C_c(K_N \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)}(C_{\text{cusp}}(G(F) \Xi \backslash G(\mathbb{A})/K_N, \mathcal{O}_E)) \end{array}$$

⁸While [14, Théorème 1.1] is stated for split G , the proof adapts for all G . Indeed, this is implicitly used in [14, Théorème 8.1].

commutes. Consequently, the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\mathfrak{z}_{K_n}(G(F_v), \mathcal{O}_E/\mathfrak{m}_E^c)$ equals $\mathfrak{z}_{n,c,I,f,\gamma_\bullet}^{\text{GL}}$.

Proof. Since Theorem 6.12 is compatible with changing Λ , the first claim follows immediately from 6.11 and Theorem 6.13. From here, tensoring with $\mathcal{O}_E/\mathfrak{m}_E^c$ shows that the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\mathfrak{z}_{K_n}(G(F_v), \mathcal{O}_E/\mathfrak{m}_E^c)$ has the same action on

$$C_{\text{cusp}}(G(F)\Xi\backslash G(\mathbb{A})/K_N, \mathcal{O}_E/\mathfrak{m}_E^c)$$

as the image of $S_{I,V,x,\xi,\gamma_\bullet}$ in $\text{Exc}(W_F, \widehat{G})$ does. Now $\mathfrak{z}_{n,c,I,f,\gamma_\bullet}^{\text{GL}}$ enjoys the same property by [14, Proposition 1.3], so they must be equal by [14, Lemma 1.4]. \square

6.16. We conclude this section by proving Theorem A. For us, *cuspidal automorphic representations* of $G(\mathbb{A})$ are irreducible summands of $C_{\text{cusp}}^\infty(G(F)\Xi\backslash G(\mathbb{A}), \overline{\mathbb{Q}}_\ell)$ for some lattice Ξ of $Z(F)\backslash Z(\mathbb{A})$.

Theorem. *The square*

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{cuspidal automorphic} \\ \text{representations of } G(\mathbb{A}) \end{array} \right\} & \xrightarrow{\text{GLC}_G} & \left\{ \begin{array}{l} L\text{-parameters} \\ \text{for } G \text{ over } F \end{array} \right\} \\ \downarrow (-)_v & & \downarrow (-)|_{W_{F_v}}^{\text{ss}} \\ \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } G(F_v) \end{array} \right\} & \xrightarrow{\text{LLC}_{G_{F_v}}^{\text{ss}}} & \left\{ \begin{array}{l} \text{semisimple } L\text{-parameters} \\ \text{for } G_{F_v} \text{ over } F_v \end{array} \right\} \end{array}$$

commutes.

Proof. Let Π be an irreducible summand of $C_{\text{cusp}}^\infty(G(F)\Xi\backslash G(\mathbb{A})/K_N, \overline{\mathbb{Q}}_\ell)$, and let N be large enough such that Π^{K_N} is nonzero. Schur's lemma shows that Π^{K_N} induces an $\overline{\mathbb{Q}}_\ell$ -algebra homomorphism

$$\chi_\Pi : \text{End}_{C_c(K_N\backslash G(\mathbb{A})/K_N, \overline{\mathbb{Q}}_\ell)}(C_{\text{cusp}}(G(F)\Xi\backslash G(\mathbb{A})/K_N, \overline{\mathbb{Q}}_\ell)) \rightarrow \overline{\mathbb{Q}}_\ell.$$

Adapt the notation of 6.13, and note that $\Pi_v^{K_n}$ induces an $\overline{\mathbb{Q}}_\ell$ -algebra homomorphism $\chi_{\Pi_v} : \mathfrak{z}_{K_n}(G(F_v), \overline{\mathbb{Q}}_\ell) \rightarrow \overline{\mathbb{Q}}_\ell$.

By Theorem 6.15, the square

$$\begin{array}{ccc} \text{Exc}(W_{F_v}, \widehat{G})_{\overline{\mathbb{Q}}_\ell} & \xrightarrow{\hspace{10em}} & \mathfrak{z}_{K_n}(G(F_v), \overline{\mathbb{Q}}_\ell) \\ \downarrow & & \downarrow \\ \text{Exc}(W_F, \widehat{G})_{\overline{\mathbb{Q}}_\ell} & \xrightarrow{\hspace{10em}} & \text{End}_{C_c(K_N\backslash G(\mathbb{A})/K_N, \overline{\mathbb{Q}}_\ell)}(C_{\text{cusp}}(G(F)\Xi\backslash G(\mathbb{A})/K_N, \overline{\mathbb{Q}}_\ell)) \end{array}$$

commutes. The composition of χ_Π with the bottom arrow corresponds to $\text{GLC}_G(\Pi)$ under Proposition 6.9, and further composition with the left arrow corresponds to $\text{GLC}_G(\Pi)|_{W_{F_v}}^{\text{ss}}$ under Proposition 6.9. On the other hand, the composition of χ_Π with the right arrow equals χ_{Π_v} , and further composition with the top arrow corresponds to $\text{LLC}_{G_{F_v}}^{\text{ss}}(\Pi_v)$ under Proposition 6.9. Hence commutativity of the square yields the desired result. \square

7. APPLICATIONS

We revert our notation to the local context: let F be a local field of characteristic $p > 0$, let G be a connected reductive group over F , and write C for its radical. Our goal in this section is to prove Theorem B, Theorem C, and Theorem D. The proofs all proceed by carefully embedding local representations into global ones.

7.1. We now prove Theorem B. Fix an isomorphism $\overline{\mathbb{Q}}_\ell \cong \mathbb{C}$.

Theorem. *If $G_{\overline{F}}^{\text{ad}}$ has any B_n or C_n factors, assume that $p \geq 3$. If $G_{\overline{F}}^{\text{ad}}$ has any F_4 or G_2 factors, assume that $p \geq 5$. Then LLC_G^{ss} uniquely lifts to a family of maps*

$$\text{LLC}_G : \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } G(F) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} L\text{-parameters} \\ \text{for } G \text{ over } F \end{array} \right\},$$

where G runs over connected reductive groups over F , that is compatible with twisting by characters, compatible with parabolic induction for essentially L^2 representations as in [29, Conjecture 4.1 (5)], and whose value on L^2 representations with finite order central character is pure.

Proof. By compatibility with parabolic induction for essentially L^2 representations, LLC_G is determined by its values on essentially L^2 representations π . By compatibility with twisting by characters, we can assume that π also has finite order central character $\omega_\pi : C(F) \rightarrow \overline{\mathbb{Q}}_\ell^\times$. There exists at most one pure L -parameter for G over F whose semisimplification equals $\text{LLC}_G^{\text{ss}}(\pi)$ [12, Lemma 3.5.(b)], so we just need to construct it.

By [13, Lemma 3.2], there exists a global field \mathbf{F} of characteristic p , a place v of \mathbf{F} , a connected reductive group \mathbf{G} over \mathbf{F} , and an isomorphism $\mathbf{F}_v \cong F$ such that

- $\mathbf{G}_{\mathbf{F}_v}$ is identified with G as group schemes over $\mathbf{F}_v \cong F$,
- the radical \mathbf{C} of \mathbf{G} has \mathbf{F} -split rank equal to the F -split rank of C .

Write $\mathbb{A}_{\mathbf{F}}$ for the adèle ring of \mathbf{F} . Let $v' \neq v$ be a place of \mathbf{F} where $\mathbf{G}_{\mathbf{F}_{v'}}$ is unramified, write $\mathbb{F}_{q'}$ for the residue field of $\mathbf{F}_{v'}$, and identify $\mathbf{F}_{v'}$ with $\mathbb{F}_{q'}(\langle \frac{1}{z} \rangle)$.

By [39, Lemma 2.1], $\mathbf{G}_{\mathbf{F}_{v'}}$ is the pullback of a connected reductive group \mathbf{H} over $\mathbb{F}_{q'}$. Let ϕ be a generic character for \mathbf{H} as in [24, Section 1.3], write Π' for the cuspidal automorphic representation of $\mathbf{H}(\mathbb{A}_{\mathbb{F}_{q'}(z)})$ associated with the automorphic sheaf A_ϕ as in [24, Definition 2.6], and write ρ' for the L -parameter for \mathbf{H} over $\mathbb{F}_{q'}(z)$ associated with the ${}^L\mathbf{H}$ -local system $\text{Kl}_{L\mathbf{H}}(\phi)$ as in [24, Theorem 1(1)]. Since A_ϕ is a Hecke eigensheaf with eigenvalue $\text{Kl}_{L\mathbf{H}}(\phi)$, we see that Π' and ρ' are associated via the Satake isomorphism at cofinitely many places of $\mathbb{F}_{q'}(z)$. Now $\rho'|_{W_{\mathbb{F}_{q'}(\langle \frac{1}{z} \rangle)}}$ is irreducible by [24, Theorem 2], so [32, Théorème 12.3] and [7, Proposition 6.4]⁹ imply that $\rho' = \text{GLC}_{\mathbf{H}}(\Pi')$. From here, Theorem 6.16 shows that $\text{LLC}_{\mathbf{H}_{\mathbb{F}_{q'}(\langle \frac{1}{z} \rangle)}}^{\text{ss}}(\Pi'_\infty) = \rho'|_{W_{\mathbb{F}_{q'}(\langle \frac{1}{z} \rangle)}}^{\text{ss}} = \rho'|_{W_{\mathbb{F}_{q'}(\langle \frac{1}{z} \rangle)}}$.

By [13, p. 2829], there exists a finite order character $\omega : \mathbf{C}(\mathbf{F}) \backslash \mathbf{C}(\mathbb{A}_{\mathbf{F}}) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ such that ω_v is identified with ω_π and $\omega_{v'}$ is identified with an unramified twist of $\omega_{\Pi'_\infty}$. Note that $\ker \omega$ contains a lattice Ξ of $\mathbf{C}(\mathbf{F}) \backslash \mathbf{C}(\mathbb{A}_{\mathbf{F}})$. Then [12, Lemma A.1] and [13, Lemma 8.1] yield an irreducible summand Π of $C_{\text{cusp}}^\infty(\mathbf{G}(\mathbf{F})\Xi \backslash \mathbf{G}(\mathbb{A}_{\mathbf{F}}), \overline{\mathbb{Q}}_\ell)$ such that

- Π_v has the same cuspidal support as π ,

⁹While [7] only considers split G , [7, Proposition 6.4] immediately extends to general G .

- $\Pi_{v'}$ is isomorphic to an unramified twist of Π'_∞ via $\mathbf{F}_{v'} \cong \mathbb{F}_{q'}(\!(\frac{1}{z})\!)$.

Theorem 6.16 and [11, p. 326] indicate that $\mathrm{GLC}_{\mathbf{G}}(\Pi)|_{W_{\mathbf{F}_{v'}}}^{\mathrm{ss}}$ equals an unramified twist of $\mathrm{LLC}_{\mathbf{H}_{\mathbb{F}_{q'}(\!(\frac{1}{z})\!)}}^{\mathrm{ss}}(\Pi'_\infty)$. This shows that $\mathrm{GLC}_{\mathbf{G}}(\Pi)$ is irreducible, so [32, Lemme 16.2] and [39, Lemma 11.4] imply that $\mathrm{GLC}_{\mathbf{G}}(\Pi)$ is pure.¹⁰ Hence $\mathrm{GLC}_{\mathbf{G}}(\Pi)|_{W_{\mathbf{F}_v}}$ is pure as in [12, Definition 3.3.(b)]. Finally, Theorem 6.16 and [11, Corollary IV.7.3] show that $\mathrm{GLC}_{\mathbf{G}}(\Pi)|_{W_{\mathbf{F}_v}}^{\mathrm{ss}} = \mathrm{LLC}_{\mathbf{G}}^{\mathrm{ss}}(\Pi_v) = \mathrm{LLC}_{\mathbf{G}}^{\mathrm{ss}}(\pi)$, so $\mathrm{GLC}_{\mathbf{G}}(\Pi)|_{W_{\mathbf{F}_v}}$ is the unique pure L -parameter for G over F whose semisimplification equals $\mathrm{LLC}_{\mathbf{G}}^{\mathrm{ss}}(\pi)$. \square

7.2. We give the following abstract proof of Theorem C.

Theorem. *There exists at most one family of maps*

$$\mathcal{LLC}_{\mathbf{G}}^{\mathrm{ss}} : \left\{ \begin{array}{l} \text{irreducible smooth} \\ \text{representations of } G(F) \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{semisimple } L\text{-parameters} \\ \text{for } G \text{ over } F \end{array} \right\},$$

where G runs over connected reductive groups over F , that is compatible with twisting by characters as in [20, Property 2.8], compatible with parabolic induction as in [20, Property 2.13], and satisfies the conclusion of Theorem 6.16.

Consequently, the Genestier–Lafforgue correspondence agrees with the Fargues–Scholze correspondence.

Proof. By compatibility with parabolic induction, $\mathcal{LLC}_{\mathbf{G}}^{\mathrm{ss}}$ is determined by its values on cuspidal representations π . By compatibility with twisting by characters, we can assume that π also has finite order central character $\omega_\pi : C(F) \rightarrow \overline{\mathbb{Q}_\ell}^\times$.

Let \mathbf{F} , v , \mathbf{G} , and \mathbf{C} be as in the proof of Theorem 7.1. By [13, Lemma 3.3], there exists a finite order character $\omega : \mathbf{C}(\mathbf{F}) \backslash \mathbf{C}(\mathbb{A}_{\mathbf{F}}) \rightarrow \overline{\mathbb{Q}_\ell}^\times$ such that ω_v is identified with ω_π . Note that $\ker \omega$ contains a lattice Ξ of $\mathbf{C}(\mathbf{F}) \backslash \mathbf{C}(\mathbb{A}_{\mathbf{F}})$. Poincaré series yield an irreducible summand Π of $C_{\mathrm{cusp}}^\infty(\mathbf{G}(\mathbf{F})\Xi \backslash \mathbf{G}(\mathbb{A}), \overline{\mathbb{Q}_\ell})$ such that Π_v is identified with π [14, Theorem 1.1], so the conclusion of Theorem 6.16 uniquely determines $\mathcal{LLC}_{\mathbf{G}}^{\mathrm{ss}}(\pi)$ as $\mathrm{GLC}_{\mathbf{G}}(\Pi)|_{W_{\mathbf{F}_v}}^{\mathrm{ss}}$.

The Fargues–Scholze correspondence satisfies the aforementioned properties by [11, p. 326], [11, Corollary IX.7.3], and Theorem 6.16. The Genestier–Lafforgue correspondence also satisfies these properties by [14, Théorème 8.1], so the above shows that it agrees with the Fargues–Scholze correspondence. \square

7.3. Finally, we prove Theorem D. Let D be a central simple algebra over F of degree n .

Theorem. *The triangle*

$$\begin{array}{ccc} \left\{ \begin{array}{l} \text{irreducible essentially } L^2 \\ \text{representations of } D^\times \end{array} \right\} & \xrightarrow{\mathrm{JL}} & \left\{ \begin{array}{l} \text{irreducible essentially } L^2 \\ \text{representations of } \mathrm{GL}_n(F) \end{array} \right\} \\ & \searrow \mathrm{LLC}_{D^\times}^{\mathrm{ss}} & \swarrow \mathrm{LLC}_{\mathrm{GL}_n}^{\mathrm{ss}} \\ & \left\{ \begin{array}{l} n\text{-dimensional semisimple} \\ \text{representations of } W_F \end{array} \right\} & \end{array}$$

commutes, where JL denotes the local Jacquet–Langlands correspondence as in [5, (th. 1.1)].

¹⁰Now [32, Lemme 16.2] and [39, Lemma 11.4] are stated for split G , but they hold in general.

Proof. Because both JL [5, (th. 1.1)] and LLC_G^{ss} [11, p. 326] are compatible with twisting by characters, it suffices to check commutativity on L^2 representations π with finite order central character $\omega_\pi : F^\times \rightarrow \overline{\mathbb{Q}_\ell}^\times$.

Let \mathbf{F} be a global field of characteristic p along with a place v of \mathbf{F} and an isomorphism $\mathbf{F}_v \cong F$, and let \mathbf{D} be a central division algebra over \mathbf{F} such that $\mathbf{D}_{\mathbf{F}_v}$ is identified with D as central simple algebras over $\mathbf{F}_v \cong F$. Using the pseudo-coefficient for $\text{JL}(\pi)$ constructed in [6, Section 5], the proof of [34, (15.10)] yields a lattice Ξ of $\mathbf{F}^\times \backslash \mathbb{A}_{\mathbf{F}}^\times$ and an irreducible summand $\tilde{\Pi}$ of $C_{\text{cusp}}^\infty(\text{GL}_n(\mathbf{F})\Xi \backslash \text{GL}_n(\mathbb{A}_{\mathbf{F}}), \overline{\mathbb{Q}_\ell})$ such that

- $\tilde{\Pi}_v$ is isomorphic to $\text{JL}(\pi)$,
- for all places $v' \neq v$ of \mathbf{F} where $\mathbf{D}_{\mathbf{F}_{v'}}$ is ramified, $\tilde{\Pi}_{v'}$ is cuspidal.

Therefore we can apply the global Jacquet–Langlands correspondence [6, Theorem 3.2] to $\tilde{\Pi}$, which yields an irreducible summand Π of $C_{\text{cusp}}^\infty(\mathbf{D}^\times \Xi \backslash (\mathbf{D} \otimes_{\mathbf{F}} \mathbb{A}_{\mathbf{F}})^\times, \overline{\mathbb{Q}_\ell})$ such that

- Π_v is isomorphic to π ,
- for all places w of \mathbf{F} where $\mathbf{D}_{\mathbf{F}_w}$ is split, Π_w is isomorphic to $\tilde{\Pi}_w$.

Then [32, Théorème 12.3] and the Chebotarev density theorem imply that

$$\text{GLC}_{\mathbf{D}^\times}(\Pi) = \text{GLC}_{\text{GL}_n}(\tilde{\Pi}),$$

so Theorem 6.16 enables us to conclude that

$$\text{LLC}_{\mathbf{D}^\times}^{\text{ss}}(\pi) = \text{GLC}_{\mathbf{D}^\times}(\Pi)|_{W_{\mathbf{F}_v}}^{\text{ss}} = \text{GLC}_{\text{GL}_n}(\tilde{\Pi})|_{W_{\mathbf{F}_v}}^{\text{ss}} = \text{LLC}_{\text{GL}_n}^{\text{ss}}(\text{JL}(\pi)). \quad \square$$

REFERENCES

- [1] J. Anshütz. Extending torsors on the punctured $\text{Spec}(A_{\text{inf}})$. *J. Reine Angew. Math.*, 783:227–268, 2022.
- [2] E. Arasteh Rad and U. Hartl. Local \mathbb{P} -shtukas and their relation to global \mathfrak{G} -shtukas. *Münster J. Math.*, 7(2):623–670, 2014.
- [3] E. Arasteh Rad and U. Hartl. Uniformizing the Moduli Stacks of Global G -Shtukas. *International Mathematics Research Notices*, 11 2019.
- [4] J. Arthur. Unipotent automorphic representations: conjectures. Number 171-172, pages 13–71. 1989. Orbites unipotentes et représentations, II.
- [5] A. I. Badulescu. Correspondance de Jacquet-Langlands pour les corps locaux de caractéristique non nulle. *Ann. Sci. École Norm. Sup. (4)*, 35(5):695–747, 2002.
- [6] A. I. Badulescu and P. Roche. Global Jacquet-Langlands correspondence for division algebras in characteristic p . *Int. Math. Res. Not. IMRN*, (7):2172–2206, 2017.
- [7] G. Böckle, M. Harris, C. Khare, and J. A. Thorne. \hat{G} -local systems on smooth projective curves are potentially automorphic. *Acta Math.*, 223(1):1–111, 2019.
- [8] F. Bruhat and J. Tits. Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d’une donnée radicielle valuée. *Inst. Hautes Études Sci. Publ. Math.*, (60):197–376, 1984.
- [9] H. Carayol. Sur les représentations l -adiques associées aux formes modulaires de Hilbert. *Ann. Sci. École Norm. Sup. (4)*, 19(3):409–468, 1986.
- [10] P. Deligne. Letter to Piatetski-Shapiro, 1973. <http://publications.ias.edu/sites/default/files/deligne73.pdf>.
- [11] L. Fargues and P. Scholze. Geometrization of the local Langlands correspondence. *arXiv e-prints*, page arXiv:2102.13459, Feb. 2021, 2102.13459.
- [12] W. T. Gan, M. Harris, W. Sawin, and R. Beuzart-Plessis. Local parameters of supercuspidal representations. *arXiv e-prints*, page arXiv:2109.07737, Sept. 2021, 2109.07737.
- [13] W. T. Gan and L. Lomeli. Globalization of supercuspidal representations over function fields and applications. *J. Eur. Math. Soc. (JEMS)*, 20(11):2813–2858, 2018.

- [14] A. Genestier and V. Lafforgue. Chtoucas restreints pour les groupes réductifs et paramétrisation de Langlands locale. *arXiv e-prints*, page arXiv:1709.00978, Sept. 2017, 1709.00978.
- [15] I. Gleason. Specialization maps for Scholze’s category of diamonds. *arXiv e-prints*, page arXiv:2012.05483, Dec. 2020, 2012.05483.
- [16] B. H. Gross and M. Reeder. Arithmetic invariants of discrete Langlands parameters. *Duke Math. J.*, 154(3):431–508, 2010.
- [17] A. Grothendieck and J. A. Dieudonné. *Éléments de géométrie algébrique. I*, volume 166 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1971.
- [18] T. J. Haines and T. Richarz. The test function conjecture for local models of Weil-restricted groups. *Compos. Math.*, 156(7):1348–1404, 2020.
- [19] D. Hansen, T. Kaletha, and J. Weinstein. On the Kottwitz conjecture for local shtuka spaces. *Forum Math. Pi*, 10:Paper No. e13, 79, 2022.
- [20] M. Harris. Local Langlands correspondences. *arXiv e-prints*, page arXiv:2205.03848, May 2022, 2205.03848.
- [21] M. Harris and R. Taylor. *The geometry and cohomology of some simple Shimura varieties*, volume 151 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001. With an appendix by Vladimir G. Berkovich.
- [22] U. Hartl. Period spaces for Hodge structures in equal characteristic. *Ann. of Math. (2)*, 173(3):1241–1358, 2011.
- [23] U. Hartl and E. Viehmann. The Newton stratification on deformations of local G -shtukas. *J. Reine Angew. Math.*, 656:87–129, 2011.
- [24] J. Heinloth, B.-C. Ngô, and Z. Yun. Kloosterman sheaves for reductive groups. *Ann. of Math. (2)*, 177(1):241–310, 2013.
- [25] R. Howe and I. I. Piatetski-Shapiro. A counterexample to the “generalized Ramanujan conjecture” for (quasi-) split groups. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, Proc. Sympos. Pure Math., XXXIII, pages 315–322. Amer. Math. Soc., Providence, R.I., 1979.
- [26] R. Huber. A generalization of formal schemes and rigid analytic varieties. *Math. Z.*, 217(4):513–551, 1994.
- [27] R. Huber. *Étale cohomology of rigid analytic varieties and adic spaces*. Aspects of Mathematics, E30. Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [28] T. Kaletha. Representations of reductive groups over local fields. *arXiv e-prints*, page arXiv:2201.07741, Jan. 2022, 2201.07741.
- [29] T. Kaletha and O. Taïbi. The local Langlands conjecture. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, R.I. To appear.
- [30] K. S. Kedlaya. Sheaves, stacks, and shtukas. In *Perfectoid spaces*, volume 242 of *Mathematical Surveys and Monographs*, pages 58–205. Amer. Math. Soc., Providence, RI, 2019.
- [31] K. S. Kedlaya and R. Liu. Relative p -adic Hodge theory: foundations. *Astérisque*, (371):239, 2015.
- [32] V. Lafforgue. Chtoucas pour les groupes réductifs et paramétrisation de Langlands globale. *J. Amer. Math. Soc.*, 31(3):719–891, 2018.
- [33] E. Landvogt. Some functorial properties of the Bruhat-Tits building. *J. Reine Angew. Math.*, 518:213–241, 2000.
- [34] G. Laumon, M. Rapoport, and U. Stuhler. \mathcal{S} -elliptic sheaves and the Langlands correspondence. *Invent. Math.*, 113(2):217–338, 1993.
- [35] S. D. Li-Huerta. The local Langlands correspondence for GL_n over function fields. *Algebra Number Theory*, 16(9):2119–2214, 2022.
- [36] M. Rapoport and T. Zink. *Period spaces for p -divisible groups*, volume 141 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1996.
- [37] M. Rapoport and T. Zink. A finiteness theorem in the Bruhat-Tits building: an application of Landvogt’s embedding theorem. *Indag. Math. (N.S.)*, 10(3):449–458, 1999.
- [38] T. Richarz. Affine Grassmannians and geometric Satake equivalences. *Int. Math. Res. Not. IMRN*, (12):3717–3767, 2016.
- [39] W. Sawin and N. Templier. On the Ramanujan conjecture for automorphic forms over function fields I. Geometry. *J. Amer. Math. Soc.*, 34(3):653–746, 2021.

- [40] P. Scholze. The local Langlands correspondence for GL_n over p -adic fields. *Invent. Math.*, 192(3):663–715, 2013.
- [41] P. Scholze. Etale cohomology of diamonds. *arXiv e-prints*, page arXiv:1709.07343, Sept. 2017, 1709.07343.
- [42] P. Scholze and J. Weinstein. *Berkeley Lectures on p -adic Geometry*, volume 207 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2020.
- [43] Y. Varshavsky. Moduli spaces of principal F -bundles. *Selecta Math. (N.S.)*, 10(1):131–166, 2004.
- [44] C. Xue. Smoothness of cohomology sheaves of stacks of shtukas. *arXiv e-prints*, Dec. 2020, 2012.12833.
- [45] Z. Yun and W. Zhang. Shtukas and the Taylor expansion of L -functions. *Ann. of Math. (2)*, 186(3):767–911, 2017.
- [46] X. Zhu. On the coherence conjecture of Pappas and Rapoport. *Ann. of Math. (2)*, 180(1):1–85, 2014.
- [47] X. Zhu. An introduction to affine Grassmannians and the geometric Satake equivalence. In *Geometry of moduli spaces and representation theory*, volume 24 of *IAS/Park City Math. Ser.*, pages 59–154. Amer. Math. Soc., Providence, RI, 2017.

Email address: `sli@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, 1 OXFORD STREET, CAMBRIDGE, MA 02138